Alterations in T lymphocytes and T-lymphocyte subpopulations in patients with syphilis

JØRGEN R JENSEN AND ELLIS FROM

From the Department of Dermatology and Venereology, University of Aarhus, Marselisborg Hospital, Aarhus, Denmark

SUMMARY The distribution of T-lymphocyte subpopulations was studied in 34 patients with primary or secondary syphilis before and after treatment. An absolute and relative T lymphopenia was found in all patients. In primary syphilis the concentration of helper cells—T cells with Fc receptors for IgM (T_H)—was low whereas in secondary syphilis the suppressor cell concentration—T cells with Fc receptors for IgG (T_S)—was reduced. Using lymphocytes from healthy subjects this could be imitated in vitro by the addition of serum from patients with secondary syphilis. In many autoimmune diseases a low concentration of T_S cells may be a primary factor in the production of autoantibodies. The occurrence of similar changes in patients with secondary syphilis, however, indicates that such fluctuations in the T-cell subpopulations may take place during a strong immune response.

Introduction

In syphilis the appearance of heterologous antibodies is diagnostic of the disease and indicates polyclonal activation of B lymphocytes. At the same time various cell-mediated immune functions seem to be reduced.¹⁻⁶ These findings indicate that patients with syphilis have an imbalance between B-cell activation and T cells with immunoregulatory functions.

In the present investigation of patients with syphilis, characteristic changes were observed in the subpopulations of T lymphocytes. These changes were related to the presence of antibodies and could be reproduced in vitro. The implication of these findings is discussed in relation to similar observations in patients with autoimmune diseases.

Patients and methods

STUDY POPULATION

Thirty-four patients with syphilis were studied. Their ages ranged from 16 to 44 years (mean 27.2 years). Seventeen had primary and 17 secondary syphilis. All had positive serological test results for syphilis. After syphilis had been diagnosed clinically, blood was collected for immunological investigations. The patients were then treated with procaine penicillin

retard (PAM) 600 000 IU daily for either 10 days (primary syphilis) or 14 days (secondary syphilis). The immunological investigations were repeated one and two months after treatment.

Ninety-eight controls were studied, ranging in age from 16 to 60 years (mean 29 years); these were staff members and other individuals who had been screened for venereal disease and were without actual disease at the time of the study.

ISOLATION OF MONONUCLEAR CELLS

Lymphocytes from heparinised blood were isolated on Ficoll-Hypaque gradients and phagocytic cells removed by the carbonyl iron incubation method. After separation, lymphocytes were washed three times in Hank's balanced salt solution (HBSS) with 2% fetal calf serum (FCS) and adjusted to 1×10^6 /ml.

ENUMERATION OF T LYMPHOCYTES

One hundred microlitres of lymphocytes (1×10^6 /ml) and 100 μ l of 2-aminoethylisothiouonium bromide hydrobromide-treated sheep erythrocytes (E-AET)⁷ were incubated at 37°C for 15 minutes in RPMI-1640 with 20% FCS in a ratio of 1:100. The mixture was spun at $150 \times g$ for five minutes and incubated at 4°C for at least one hour. After 10 μ l of 0.1% acridine orange had been added the pellet was carefully resuspended, and 200 lymphocytes were counted with a fluorescence microscope. Lymphocytes with three or more erythrocytes bound to their surface were counted as rosette-forming cells.

Address for reprints: Dr J R Jensen, Department of Dermatology and Venereology, Marselisborg Hospital, 8000 Aarhus C, Denmark

Accepted for publication 14 July 1981

The subfraction of T lymphocytes with high affinity to sheep erythrocytes form rosettes with untreated sheep erythrocytes (E-active or E-a rosettes). One hundred microlitres of lymphocytes suspended in RPMI-1640 without FCS were incubated with 100 μ l untreated sheep erythrocytes in a ratio of 1:10 for five minutes at 37°C. The mixture was spun for five minutes at 150 $\times$ g, incubated at 4°C for five minutes, and resuspended for rosette enumeration.

PURIFICATION OF T LYMPHOCYTES

Five millilitres of E-AET and 5 ml of lymphocytes were mixed and incubated as described above. After resuspension, Ficoll-Hypaque flotation was performed (600 $\times$ g for 15 minutes). The pellet of E-AET-rosetting T lymphocytes was resuspended in 0.83% TRIS NH_4Cl (10 minutes) to lyse the erythrocytes. After three washes with HBSS without serum the T lymphocytes were adjusted to a concentration of 1×10^6 /ml. Purity, as determined by E-AET rosetting of the T-lymphocyte suspensions, was between 91% and 98%. Lymphocytes with surface immunoglobulin (B lymphocytes) were not found in the purified T-lymphocyte suspensions.

PREPARATION OF ANTI-ORBC ANTIBODIES

Rabbits were immunised by weekly subcutaneous injections of 1 ml of 10% ox red cells (ORBC) in Freund's incomplete adjuvant. Bleedings were performed weekly starting one week after the first immunisation.⁸ Serum was inactivated at 56°C for 30 minutes and IgG was purified by DEAE-cellulose ion-exchange chromatography. IgM was obtained by gel filtration (Ultrogel AcA 22, LKB). Both the IgG and IgM suspensions gave single peaks in a crossed immunoelectrophoresis against swine anti-rabbit immunoglobulins. Purity of the IgM suspension was further indicated by the lack of rosetting capacity of monocytes with ORBC coated with IgM antibodies; all monocytes with ORBC coated with IgG antibody, however, formed rosettes.

ERYTHROCYTE-ANTIBODY (EA) SUSPENSIONS

ORBC-antibody complexes were prepared from ox erythrocytes in Alsevers solution, washed five times in HBSS, and resuspended to a concentration of 2%. Subagglutinating concentrations of IgG or IgM anti-ORBC antibody were mixed with equal volumes of 2% ORBC and kept for 90 minutes at room temperature. Three washes of the ORBC-antibody complexes were done in HBSS with 2% FCS. Finally, 10% FCS was added. These erythrocyte-antibody (EA) suspensions were kept at 4°C and used during the next two days.

ENUMERATION OF T-LYMPHOCYTE SUBPOPULATIONS

Purified T lymphocytes were incubated in RPMI-1640 with 20% FCS at 37°C in a humidified CO_2 atmosphere. After 24 hours' (day 1) and 48 hours' (day 2) incubation appropriate amounts of T lymphocytes were washed three times in HBSS with 2% FCS. The T lymphocytes (0.01 ml) were mixed with ORBC coated with subagglutinating doses of either IgG (0.01 ml) or IgM (0.01 ml), spun immediately at 150 $\times$ g for five minutes, and incubated at 4°C for three hours. The percentage of cells forming rosettes was counted as described above.

SERUM ASSAYS

Immunoglobulins in serum from each of three patients with secondary syphilis were removed by ammonium sulphate precipitation. Sera from three controls were treated similarly. T lymphocytes from the three controls were cultured for 24 hours and incubated for 60 minutes at 37°C with autologous serum, autologous serum without immunoglobulins, syphilitic serum, and syphilitic serum without immunoglobulins. T lymphocytes without addition of serum acted as controls. Serum was not inactivated by heat. Cell numbers and cell deaths were controlled before and after serum treatment. Three washes with HBSS with 2% FCS were done before rosette enumeration.

STATISTICAL ANALYSIS

Student's *t* test (two-tailed) was used.

Results

The total number of T lymphocytes (E-AET rosette-forming lymphocytes) was significantly reduced during both primary and secondary syphilis (table I). This was due to lymphopenia, since the percentage of T lymphocytes was within the normal range although slightly reduced (table II).

The subfraction of T lymphocytes with high affinity for sheep erythrocytes (E-a rosette-forming lymphocytes) was also reduced (table I). In patients with primary syphilis both an absolute and relative reduction of E-a T lymphocytes was found, whereas in patients with secondary syphilis only the absolute number of E-a T cells was reduced (table II).

During primary syphilis the percentage of T lymphocytes with Fc receptors for IgM (T_μ) was significantly reduced. T lymphocytes with Fc receptors for IgG (T_γ) were slightly increased ($t_{98}=1.50$, $0.1 < P < 0.2$; fig 1). In patients with secondary syphilis this was almost reversed. The

TABLE I Total number of T lymphocytes in peripheral blood from patients with syphilis and controls

	T cells per mm ³ (mean ± SEM) (n):			
	Before treatment	One month after treatment	Two months after treatment	Controls
E-AET				
Primary syphilis	1479 ± 213(9)*	2052 ± 293(6)	2045 ± 228(7)	2089 ± 73·11(87)
Secondary syphilis	1579 ± 183(13)†	2204 ± 224(10)	2651 ± 352(9)	
E-a				
Primary syphilis	829 ± 63(9)‡	1175 ± 157(7)	1061 ± 229(6)	1179 ± 47·24(84)
Secondary syphilis	882 ± 83(12)§	1302 ± 130(12)	1477 ± 245(8)	

SEM = standard error of mean; n = No of patients or controls; E-AET = 2-aminoethylisothiuronium bromide hydrobromide-treated sheep erythrocytes; E-a = untreated sheep erythrocytes
*Significantly reduced (*t*₉₄ = 2·71, *P* < 0·01)
†Significantly reduced (*t*₉₈ = 2·58, *P* < 0·02)
‡Significantly reduced (*t*₉₁ = 4·45, *P* < 0·005)
§Significantly reduced (*t*₉₄ = 3·12; *P* < 0·005)

TABLE II Percentage of T lymphocytes in peripheral blood from patients with syphilis and controls

	Percentage of T cells (mean ± SEM) (n):			
	Before treatment	One month after treatment	Two months after treatment	Controls
E-AET				
Primary syphilis	77·4 ± 2·5(15)	76·7 ± 1·2(10)	78·7 ± 1·4 (7)	80·5 ± 0·55(98)
Secondary syphilis	77·6 ± 2·0(16)	81·4 ± 1·2(14)	78·3 ± 1·5(11)	
E-a				
Primary syphilis	40·4 ± 1·8(14)*	41·7 ± 1·7(9)†	38·3 ± 3·3(7)‡	45·9 ± 0·93(95)
Secondary syphilis	44·1 ± 2·4(15)	46·7 ± 3·8(14)	41·0 ± 2·1(10)	

SEM = standard error of mean; n = No of patients or controls; E-AET = 2-aminoethylisothiuronium bromide hydrobromide-treated sheep erythrocytes; E-a = untreated sheep erythrocytes
*Significantly reduced (*t*₁₀₇ = 2·7, *P* < 0·01)
†Significantly reduced (*t*₁₀₂ = 2·15, *P* < 0·05)
‡Significantly reduced (*t*₁₀₀ = 2·23, *P* < 0·05)

FIG 1 The percentage of Ty and Tu rosettes in patients with primary syphilis before treatment (A, n = 14), one month after (B, n = 7), and two months after treatment (C, n = 7) in T cells cultured for 24 hours and 48 hours compared with controls (n = 84). Results represent mean ± SEM (Student's *t* test).

FIG 2 The percentage of Ty and Tu rosettes in patients with secondary syphilis before treatment (A, n = 9), one month after (B, n = 11), and two months after treatment (C, n = 12) in T cells cultured for 24 hours and 48 hours compared with controls (n = 84). Results represent mean ± SEM (Student's *t* test).

percentage of T μ lymphocytes was normal whereas that of T γ cells was significantly reduced (fig 2).

The reduced percentage of T γ cells during the secondary stage was investigated in vitro by adding serum from each of three patients with secondary syphilis to T lymphocytes from three healthy donors. Complement was not inactivated by heat. The expression of the Fc-IgG receptor on the T lymphocytes was significantly reduced without selective killing of T γ cells ($t_2=6.48$, $P<0.025$). When the immunoglobulin in serum was removed through ammonium sulphate precipitation this effect disappeared (table III). No reduction of the expression of the Fc-IgM receptor was observed in similar experiments using serum from a patient with primary syphilis (results are not shown).

Discussion

The importance of cell-mediated immunity (CMI) in syphilis is suggested by T-lymphocyte depletion in the paracortical areas of lymph nodes and the periarteriolar area of the spleen in children with fatal congenital syphilis.^{9,10} In rabbits, the infection takes a severe course if T lymphocytes are destroyed by cyclophosphamide treatment.¹¹ In patients with syphilis evidence has been found for a depression in CMI, which involves cutaneous reactivity,¹ low T-cell numbers,⁶ low in-vitro reactivity to *Treponema pallidum*,^{1,3-5} and immunosuppressive factors in serum.^{1,2}

This study demonstrates that the total number of T lymphocytes are reduced in patients with early syphilis. The T lymphopenia may be due to an unequal distribution of lymphocytes in the lymphoid system, although a depletion of lymphocytes is also seen in histological examinations of enlarged lymph nodes during early syphilis.¹² Another possibility is

that cytotoxic antibodies to T lymphocytes may occur, although lymphocytotoxic antibodies were shown to be active only at 15°C.¹³ Also, the T lymphopenia disappears rapidly after treatment and the serum assays do not show any enhanced cell death. A third possibility is that *T pallidum* itself is directly toxic to T lymphocytes.

It has been proposed that T lymphocytes with Fc receptors may function as helper (T μ) or suppressor (T γ) cells.¹⁴ In primary syphilis the T μ -cell concentration was low, consistent with compartmentalisation of helper cells—that is, an unequal distribution of lymphocytes in the body during an immune response.¹⁵ Compartmentalisation of helper cells occurs during early antibody production.^{16,17} The T γ -cell concentration was increased, perhaps as a result of suppressor-cell activation.¹⁸

In patients with secondary syphilis the T μ -cell concentration was normal until two months after treatment when a fall occurred. The reason for this fall is obscure. During active secondary syphilis the T γ -cell concentration was significantly reduced. Explanations for this may be similar to those for the T lymphopenia. Furthermore, during secondary syphilis immune complex formation takes place,¹⁹ and it is well known that immune complexes interact in vitro with T γ cells leading to an irreversible loss of the Fc receptor for IgG.²⁰ This mechanism may explain the reduction of the T γ cells in peripheral blood as indicated by the results from the serum assays (table III). No similar mechanism seemed to operate in primary syphilis.

It has been found that T γ cells activated by pokeweed mitogen and immune complexes release immunosuppressive factors.²¹ *T pallidum* may give an in-vivo stimulation, which in conjunction with immune complexes stimulates T γ cells to release immunosuppressive factors. This may explain the

TABLE III Effect of secondary syphilitic serum on homologous T lymphocyte subpopulations in three controls

Donor serum (40%)	Subpopulations	Percentage of T lymphocyte subpopulations in controls:		
		Control 1	Control 2	Control 3
Autologous serum	T γ	9.0	15.0	13.0
	T μ	31.5	53.0	40.5
Autologous serum without Ig	T γ	9.5	18.0	16.0
	T μ	29.5	36.5	39.0
Secondary syphilitic serum	T γ^*	3.0	4.0	4.0
	T μ	33.0	50.5	43.5
Secondary syphilitic serum without Ig	T γ	8.5	17.0	14.0
	T μ	33.0	37.0	43.0
Controls without serum	T γ	11.0	20.0	17.5
	T μ	33.0	36.0	40.5

*Significant reduction (78%; $t_2=6.48$, $P<0.025$)

reduced cutaneous reactivity in syphilitic patients¹ and the depressed in-vitro functions of lymphocytes from peripheral blood.^{1,3-5}

Reduction of the Ty-cell concentration has been observed in various autoimmune diseases and in other diseases with increased antibody production.²²⁻²⁸ The increased antibody production may be caused by direct polyclonal B-cell activation or indirectly by a defective suppressor-cell mechanism, which may be a primary disorder or secondary to the occurrence of lymphocytotoxic antibodies or immune complexes. In systemic lupus erythematosus (SLE) the suppressor-cell defect is mainly a quantitative rather than a qualitative disorder.²⁹ Immunosuppressive factors, which depress non-specific suppressor-cell activity, have also been found in SLE.^{30,31}

The changes observed in the subpopulations of T lymphocytes occurred during an infectious disease in young subjects who were not suspected of having any immune deficiencies. These changes are similar to those found in various diseases with an imbalance between an increased B-cell activity and signs of reduced immunoregulatory T cells.²²⁻²⁸ Our findings seem to show that the changes in immunoregulatory T cells may be secondary to a strong immune reaction—that is, treponemal infection.

This work was supported by the Danish Medical Research Council, grants Nos 512-15133, 512-20160, and 512-20207. We wish to thank Mrs A Jespersen and J Schjøth for technical assistance and Dr K Thestrup-Pedersen for his constant encouragement and for reviewing the manuscript.

References

- From E, Thestrup-Pedersen K, Thulin H. Reactivity of lymphocytes from patients with syphilis towards *T pallidum* antigen in the leucocyte migration and lymphocyte transformation tests. *Br J Vener Dis* 1976; **52**:224-9.
- Levene GM, Turk JI, Wright DJM, Grimble AGS. Reduced lymphocyte transformation due to a plasma factor in patients with active syphilis. *Lancet* 1969; **ii**:246-7.
- Fulford KWM, Brostoff J. Leucocyte migration and cell-mediated immunity in syphilis. *Br J Vener Dis* 1972; **48**:483-8.
- Pavia CS, Baseman JB, Folds JD. Selective response of lymphocytes from *Treponema pallidum*-infected rabbits to mitogens and *Treponema reuteri*. *Infect Immun* 1977; **15**:417-22.
- Pavia CS, Folds JD, Baseman JB. Selective in-vitro response of thymus-derived lymphocytes from *Treponema pallidum*-infected rabbits. *Infect Immun* 1977; **18**:603-11.
- Bos JD, Hamerlinck F, Cormane RH. T lymphoid cells in primary syphilis. Quantitative studies. *Br J Vener Dis* 1980; **56**:74-6.
- Kaplan ME, Clark C. An improved rosetting assay for the detection of human T lymphocytes. *J Immunol Methods* 1974; **5**:131-5.
- Gupta S, Good RA. Subpopulations of human T lymphocytes I. Studies in immunodeficient patients. *Clin Exp Immunol* 1977; **30**:222-8.
- Levene GM, Wright DJM, Turk JL. Cell-mediated immunity and lymphocyte transformation in syphilis. *Proc R Soc Med* 1971; **64**:426-8.
- Wright DJM, Grimble AS. Why is the infectious stage of syphilis prolonged? *Br J Vener Dis* 1974; **50**:45-9.
- Pacha J, Metzger M, Smogór W, Michalska E, Podwinska J, Ruczkowska J. Effect of immunosuppressive agents on the course of experimental syphilis in rabbits. *Arch Immunol Ther Ex (Warsz)* 1979; **27**:45-51.
- Turner DR, Wright DJM. Lymphadenopathy in early syphilis. *J Pathol* 1973; **110**:305-8.
- de Jong NHL, Koehorst JAM, van der Sluis JJ, Boer AM. Autolymphocytotoxins in syphilis. *Br J Vener Dis* 1980; **56**:297-301.
- Moretta L, Webb SR, Grossi CE, Lydyard PM, Cooper MD. Functional analysis of two human T-cell subpopulations, helper and suppression of B-cell responses by T cells bearing receptors for IgM and IgG. *J Exp Med* 1977; **146**:184-200.
- Schlossman SF, Levin HA, Rocklin RE, David JR. The compartmentalization of antigen-reactive lymphocytes in desensitized guinea pigs. *J Exp Med* 1971; **134**:741-50.
- Lagrange PH, Mackaness GB, Miller TE. Influence of dose and route of antigen injection on the immunological induction of T cells. *J Exp Med* 1974; **139**:528-42.
- Hahn H, Kaufmann SHE, Falkenberg F, Chahinin M, Horn W. Peritoneal exudate T lymphocytes with specificity to sheep red blood cells. II Inflammatory helper T cells and effector T cells in mice with delayed-type hypersensitivity and in suppressed mice. *Immunology* 1979; **38**:51-5.
- Ramshaw IA, Bretscher PA, Parish CR. Regulation of the immune response. I Suppression of delayed type hypersensitivity by T cells from mice expressing humoral immunity. *Europ J Immunol* 1979; **6**:674-9.
- Sølling J, Sølling K, Jakobsen KU, From E. Circulating immune complexes in syphilis. *Acta Derm Venereol (Stockh)* 1978; **58**:263-7.
- Pichler WJ Lum L, Border S. Fc receptors on human T lymphocytes. I Transition of Ty to T_μ cells. *J Immunol* 1978; **121**:1540-8.
- Moretta L, Mingari MC, Moretta A, Cooper MD. Human T-lymphocyte subpopulations: studies of the mechanism by which T cells bearing Fc receptors for IgG suppress T-dependent B cell differentiation induced by PWM. *J Immunol* 1979; **122**:984-90.
- Fauci AS, Steinberg AD, Haynes BF, Whalen G. Immunoregulatory aberrations in systemic lupus erythematosus. *J Immunol* 1978; **121**:1473-9.
- Alarcón-Segovia D, Ruiz-Argüelles A. Decreased circulation thymus-derived cells with receptors for the Fc portion of immunoglobulin G in systemic lupus erythematosus. *J Clin Invest* 1978; **62**:1390-4.
- Reinherz EL, Rubinstein A, Geha RS, Strelkauskas AJ, Rosen FS, Schlossman SF. Abnormalities of immunoregulatory T cells in disorders of immune function. *N Engl J Med* 1979; **301**:1018-22.
- Sandilans GP, Galbraith I, Reid FM, Mills PR, MacSween RMN. Immune-complex inhibition of lymphocyte Fc receptors in primary biliary cirrhosis. A possible immunomodulatory mechanism. *Lancet* 1980; **ii**:9-13.
- Reinherz EL, Weiner HL, Hauser SL, Cohen JA, Distaso JA, Schlossman SF. Loss of suppressor T cells in active multiple sclerosis. Analysis with monoclonal antibodies. *N Engl J Med* 1980; **303**:125-9.
- Bach M-A, Phan-Dinh-Tuy F, Tournier E, Chatenoud L, Bach J-F. Deficit of suppressor T cells in active multiple sclerosis. *Lancet* 1980; **ii**:1221-3.
- Jensen JR, Cramers M, Thestrup-Pedersen K. Subpopulations of T lymphocytes and non-specific suppressor-cell activity in patients with atopic dermatitis. *Clin Exp Immunol* 1981; **45**:118-23.
- Moretta M, Mingari MC, Santoli D, Perlman P, Moretta L. Human T-lymphocyte subpopulations: alterations in systemic lupus erythematosus. *Scand J Immunol* 1979; **10**:223-8.
- Sagawa A, Abdou NI. Suppressor-cell antibody in systemic lupus erythematosus. Possible mechanism for suppressor-cell dysfunction. *J Clin Invest* 1979; **63**:536-9.
- Sakane T, Steinberg AD, Reeves JP, Green I. Studies of immune functions of patients with systemic lupus erythematosus. Complement-dependent immunoglobulin M anti-thymus-derived cell antibodies preferentially inactivated suppressor cells. *J Clin Invest* 1979; **63**:954-65.