unusual. The authors prefer benzyl benzoate to other treatments but insist that all members of the family and all sexual contacts should be treated even in the absence of overt lesions. Factors which encourage the spread of scabies are mass movements of population, depression and promiscuity. It is thought that pruritus and papulovesicular lesions are due to allergic hypersensitivity as the small number of mites does not account for the extent of lesions seen. Postscabietic pruritus and eosinophilia also support this view. One explanation for the present spread of scabies might be due to the previous low degree of infestation and absence of an immune response. This would fit in with the present spread of the infestation in children. Infestation does not lead to true immunity but is thought to produce a reactivity of the body giving rise to early symptoms when infected. G. W. Csonka

Torsion of the Testis and Its Appendages CHAPMAN, R. H., and WALTON, A. J. (1972) Brit. med. J., 1, 164 15 refs

Prophylatic Treatment with Penicillin in Venereal Diseases [in German] LUGER, A. (1971) Hautarzt, 22, 135

Behcet's Disease O'DUFFY, J. D., CARNEY, J. A., and DEODHAR, S. (1971) Ann. intern. Med., 75, 561

Genito-urinary Tuberculosis Gow, J. G. (1971) Practitioner, 207, 609 1 fig., 4 refs Cell-dependent Differences in the Production of Infectious Herpes Simplex Virus at a Supraoptimal Temperature CROUCH, N. A., and RAPP, F. (1972) J. Virol., 9, 223 1 fig., 22 refs

Herpes Simplex Virus Infection in the Newborn PETTAY, O., LEINIKKI, P., DONNER, M., and LAPINTEIMU, K., (1972) Arch. Dis. Childh., 47, 97 2 figs, 18 refs

Herpes Antibody and Cervical Carcinoma in situ CATALANO, L. W., and JOHNSON, L. D., (1971) J. Amer. med. Ass., 217, 447 10 refs

Condylomata Acuminata (1972) Brit. med. J., 2, 179

Correction

In the paper by A. Forsgren, which appeared in the June issue of the journal (Brit. J. vener. Dis. (1972), 48, 205), in the Table on p. 205, col. 1, row 11, for 36°C, please read 56°C.