

Subject index

- access**, Maintaining patient access to GUM clinics: is it compatible with appointments?, 11
Treating sexually transmitted infections in primary care: a missed opportunity?, 134
- accident and emergency department**, Urine based screening for asymptomatic/undiagnosed genital chlamydial infection in young people visiting the accident and emergency department is feasible, acceptable, and can be epidemiologically helpful, 229
- adolescence**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, 31
- adolescent health**, Adolescent women underestimate their susceptibility to sexually transmitted infections, 408
- adolescents**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, 469
- adult respiratory distress syndrome**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, 337
- Africa**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, 98
Male circumcision: an acceptable strategy for HIV prevention in Botswana, 214
Sexually transmitted infections in Africa: single dose treatment is now affordable, 432
- age**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, 286
- AIDS**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, 179
Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, 45
HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, 442
Male circumcision: an acceptable strategy for HIV prevention in Botswana, 214
Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, 62
Television coverage of AIDS and transmission of the AIDS virus, 170
- AIDS/STD**, Effects of two telephone survey methods on the level of reported risk behaviours, 325
- amphotericin B**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, 265
- antenatal syphilis screening**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, 375
- antibiotics**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, 399
- antibodies**, Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, 351
- antiretroviral therapy**, Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, 7
Lymphocytic interstitial pneumonitis in HIV infected adults, 88
- appointments**, Maintaining patient access to GUM clinics: is it compatible with appointments?, 11
- attitude**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, 35
- Australia**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, 31
Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, 332
- azithromycin**, Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, 453
- bacterial vaginosis**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, 224
- behaviour**, Determinants of disclosure of genital herpes to partners, 42
- Belgium**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, 157
- bisexuality**, Sexual behaviour of lesbians and bisexual women, 147
- black township**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, 442
- Botswana**, Male circumcision: an acceptable strategy for HIV prevention in Botswana, 214
- Brazil**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, 448
Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, 286
- Britain**, Male circumcision in Britain: findings from a national probability sample survey, 499
- Cambodia**, Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, e2
- Candida albicans**, Vulvovaginal carriage of yeasts other than *Candida albicans*, 249
- Candida glabrata**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, 265
Vulvovaginal carriage of yeasts other than *Candida albicans*, 249
- Castleman's disease**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, 179
- cutaneous larva migrans**, Cutaneous larva migrans of the penis, 500
- ceftriaxone**, Treatment of neurosyphilis with ceftriaxone, 415
- cervical cancer**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, 191
- cervical screening**, Awareness of human papillomavirus among women attending a well woman clinic, 320
- cervix**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, 496
- cervix uterus**, Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, 474
- chancere**, Perforating chancre: any cause-effect relation with HIV infection?, 262
- chancroid**, Chancroid: clinical manifestations, diagnosis, and management, 68
Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, 294
- chaperone**, Chaperoning in genitourinary medicine clinics, 422
- chaperones**, Chaperoning in genitourinary medicine: supporting patients and protecting doctors, 498
- chlamydia**, Chlamydia testing before termination of pregnancy, 352
Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, 224
LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, 94
Recent pilot studies of chlamydia screening, 352
Repeat chlamydia screening by mail, San Francisco, 28
Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, 372
Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, 62
Validity and utility of screening tests for STIs, 356
- chlamydia screening**, Urine based screening for asymptomatic/undiagnosed genital chlamydial infection in young people visiting the accident and emergency department is feasible, acceptable, and can be epidemiologically helpful, 229

- Chlamydia trachomatis**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
- Are all genital *Chlamydia trachomatis* infections pathogenic? A study in men, **349**
- Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
- Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
- Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- Spending money to save money, **4**
- chlamydial infection**, Chlamydia screening in the United Kingdom, **3**
- First, do not harm: also an issue in NAA assay diagnostics for chlamydial infection, **76**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- circumcision**, Circumcision practice patterns in South Korea: community based survey, **65**
- HIV and circumcision: new factors to consider, **495**
- Issues associated with the introduction of circumcision into a non-circumcising society, **427**
- Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
- clinical practice variations**, Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- cognitions**, Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- cohort study**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- Colombia**, Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- computerised questionnaire**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- condom negotiation skills**, Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- condoms**, Don't overlook condoms for HIV prevention, **268**
- congenital syphilis**, Congenital syphilis in the Russian Federation: magnitude, determinants, and consequences, **106**
- continuing medical education**, Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- cost effectiveness**, A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
- cost effectiveness analysis**, Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Spending money to save money, **4**
- Costa Rica**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- cost-effectiveness analysis**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- counselling**, Genital herpes serotyping: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- couple transmission**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- cranial tuberculomas**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- Cryptosporidium**, Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- cytokines**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- cytolethal distending toxin**, Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, **294**
- DALYs**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- dark ground microscopy**, Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- delay**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
- demographic risk factors**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- diagnosis**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- diagnostics**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- doctors**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- drug resistance**, New drugs for treating drug resistant HIV-1, **176**
- dual neoplastic diseases**, Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- education**, Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- education initiatives**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- EIA amplicon detection**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
- England**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- environmental contamination**, Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
- epidemiology**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- Estonia**, Seroprevalence study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- ethnicity**, Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- Europe**, A survey of STI policies and programmes in Europe, **171**
- family planning**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- female bar workers**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- female circumcision**, Female genital mutilation in the Sudan: survey of the attitude of Khartoum university students towards this practice, **220**
- female genital mutilation**, Female genital mutilation in the Sudan: survey of the attitude of Khartoum university students towards this practice, **220**
- female sex workers**, Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- females**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**

- flucytosine**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
- follow up**, Patients lost to follow up: experience of an HIV clinic, **349**
- foreskin**, Circumcision practice patterns in South Korea: community based survey, **65**
- general practice**, Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- genital chlamydia infection**, Who is being tested for genital chlamydia in primary care?, **234**
- genital diseases**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**
- genital herpes**, Determinants of disclosure of genital herpes to partners, **42**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Genital herpes serotyping: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- Is HSV serology useful for the management of first episode genital herpes?, **276**
- genital infection**, Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
- genital ulcer disease**, Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- genital warts**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- genital warts, circumcision, treatment**, Circumcision in genital warts—let us not forget!, **265**
- genitourinary diseases**, Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- genitourinary infection**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- genitourinary infections**, Self obtained samples for genitourinary infections, **171**
- genitourinary medicine**, Chaperoning in genitourinary medicine: supporting patients and protecting doctors, **498**
- genitourinary medicine clinics**, Chaperoning in genitourinary medicine clinics, **74**
- Raising awareness of UK GUM clinic activities, **264**
- genitourinary medicine education**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- genotypes**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- Germany**, Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, **351**
- glucocorticoids**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- gold mining**, Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- gonorrhoea**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
- Gonorrhoea reinfection in heterosexual STD clinic attendees: longitudinal analysis of risks for first reinfection, **124**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- Pharyngeal gonorrhoea: the forgotten reservoir, **345**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- The fall and rise of gonorrhoea incidence in Israel: an international phenomenon?, **116**
- Greater Manchester**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- GUM clinics**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
- GUM services**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- gynaecological examination**, The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- HAART**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- Polymorphic immune restoration syndrome after effective HAART, **171**
- Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- Haemophilus ducreyi**, Chancroid: clinical manifestations, diagnosis, and management, **68**
- healthcare settings**, Chlamydia screening in the United Kingdom, **3**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- hepatitis B**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- hepatitis C**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- hepatitis C testing**, Hepatitis C testing in HIV infected patients, **76**
- hepatitis C virus**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**
- Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- herpes simplex virus**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- herpes simplex virus serology**, Is HSV serology useful for the management of first episode genital herpes?, **276**
- herpes simplex virus seroprevalence**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- HIV**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals, **137**
- Don't overlook condoms for HIV prevention, **268**
- Enhanced risk of HIV sexual transmission during structured treatment interruption, **74**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Hepatitis C testing in HIV infected patients, **76**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**

- HIV and circumcision: new factors to consider, **495**
- HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Lower motor neuron syndrome and HIV infection, **351**
- Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
- New drugs for treating drug resistant HIV-1, **176**
- Perforating chancres: any cause-effect relation with HIV infection?, **262**
- Polymyositis masquerading as mitochondrial toxicity, **417**
- Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- Superior mesenteric artery syndrome in an HIV positive patient, **262**
- The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, **453**
- Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- View 1: Assisted conception in couples with HIV infection, **185**
- Commissioning HIV services, **84**
- View 2: Assisted conception in couples with HIV infection, **189**
- HIV clinic**, Patients lost to follow up: experience of an HIV clinic, **349**
- HIV infection**, Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
- Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- What's new for injection drug users with HIV infection?, **80**
- HIV prevention**,
- HIV testing attitudes**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- HIV transmission risks**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- HIV treatment adherence**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- HIV/AIDS epidemic**, The HIV/AIDS epidemic in Ukraine: stable or still exploding?, **263**
- HIV/AIDS**, Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- homosexual men**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- homosexuality**, Sexual behaviour of lesbians and bisexual women, **147**
- hospital clinic**, Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- human immunodeficiency virus**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- human papillomavirus**, Awareness of human papillomavirus among women attending a well woman clinic, **320**
- Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
- Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- human parvovirus B19**, Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- hygiene**, Circumcision practice patterns in South Korea: community based survey, **65**
- immune reconstitution**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- income inequality**, Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- India**, Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
- indigenous people**, Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, **332**
- infertility**, View 1: Assisted conception in couples with HIV infection, **185**
- View 2: Assisted conception in couples with HIV infection, **189**
- injecting drug use**, Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- injection drug users**, What's new for injection drug users with HIV infection?, **80**
- internet**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- intimate examination**, Chaperoning in genitourinary medicine clinics, **422**
- Israel**, The fall and rise of gonorrhoea incidence in Israel: an international phenomenon?, **116**
- Haemophilus ducreyi**, Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, **294**
- Sexually Transmitted Infections**, STI case management at a South African teaching hospital, **498**
- Taking up the reins, **2**
- Jacobson JM, editor Immunotherapy for Infectious Diseases**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
- journalology**, Taking up the reins, **2**
- Kaposi's sarcoma**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
- knowledge**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- lactic acidosis**, Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
- lesbians**, Sexual behaviour of lesbians and bisexual women, **147**
- leucocyte esterase dipstick**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- ligase chain reaction**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- lipodystrophy**, A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals, **137**

- lymphocytes**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- lymphogranuloma venereum**, Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, **453**
- male circumcision**, Male circumcision in Britain: findings from a national probability sample survey, **499**
- male clients**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- maternal syphilis**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- men**, Are all genital *Chlamydia trachomatis* infections pathogenic? A study in men, **349**
- Mexico**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- microscopy**, Poor sensitivity and consistency of microscopy in the diagnosis of low grade non-gonococcal urethritis, **487**
- middle aged men**, Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- middle aged women**, Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- mitochondrial toxicity**, Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
Polymyositis masquerading as mitochondrial toxicity, **417**
- mixed infection**, Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
- molecular typing**, Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
- monosymptomatic hypochondriacal psychosis**, Monosymptomatic hypochondriacal psychosis, **425**
- Morocco**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- mother to child transmission**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- motor neuron**, Lower motor neuron syndrome and HIV infection, **351**
- Mozambique**, Geographical focusing: an intervention to address increased risk for sexually transmitted diseases during repatriation and resettlement in post-war Mozambique, **77**
- MSM**, Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- MSSVD**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- multidrug resistance**, Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- muscle biopsy**, Polymyositis masquerading as mitochondrial toxicity, **417**
- Mycoplasma genitalium**, Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
- Mycoplasma infections**, Tetracycline treatment does not eradicate *Mycoplasma genitalium*, **318**
- myopathy**, Polymyositis masquerading as mitochondrial toxicity, **417**
- Neisseria gonorrhoeae**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
- neurology**, Lower motor neuron syndrome and HIV infection, **351**
- neurosyphilis**, Treatment of neurosyphilis with ceftriaxone, **415**
- non-gonococcal urethritis**, Poor sensitivity and consistency of microscopy in the diagnosis of low grade non-gonococcal urethritis, **487**
- non-paying partners**, Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- nucleic acid amplification assay**, First, do not harm: also an issue in NAA assay diagnostics for chlamydial infection, **76**
- nurses**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- opportunistic testing**, Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- outreach programmes**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- outreach service**, Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- paediatric AIDS**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- Papanicolaou smear**, Was the Papanicolaou smear responsible for the decline of *Trichomonas vaginalis*?, **263**
- papillomavirus**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, **191**
- partner notification**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
Partner notification in primary care, **264**
- pelvic inflammatory disease**, Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
Diagnosis of pelvic inflammatory disease: time for a rethink, **491**
Spending money to save money, **4**
- penis**, Cutaneous larva migrans of the penis, **500**
- perianal**, Perianal verrucous epidermal naevus mimicking perianal warts, **424**
- perianal verrucous epidermal naevus**, Perianal verrucous epidermal naevus mimicking perianal warts, **424**
- perinatal mortality**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- perinatal transmission**, Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- Peru**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- pharmacists**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- phimosi**, Circumcision practice patterns in South Korea: community based survey, **65**
- pneumonitis**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- podophyllin**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- podophyllotoxin**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- point of care diagnosis**, Validity and utility of screening tests for STIs, **356**
- polymerase chain reaction**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
- polymorphic immune restoration syndrome**, Polymorphic immune restoration syndrome after effective HAART, **171**
- polymyositis**, Polymyositis masquerading as mitochondrial toxicity, **417**

- pontine myelinolysis**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
- population surveys**, LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- post-risk behaviours**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- postmenopausal woman**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- PPNG**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
- pregnancy**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- pretest counselling**, Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- primary care**, Partner notification in primary care, **264**
Surveillance of sexually transmitted infections in primary care, **174**
Treating sexually transmitted infections in primary care: a missed opportunity?, **134**
Who is being tested for genital chlamydia in primary care?, **234**
- priority setting**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- private practice**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- psychology**, Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- psychosocial impact**, Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- public health**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
Validity and utility of screening tests for STIs, **356**
- public understanding**, Awareness of human papillomavirus among women attending a well woman clinic, **320**
- qualitative assessment**, Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- randomised controlled trial**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- recurrent infection**, Repeat chlamydia screening by mail, San Francisco, **28**
- reinfection**, Gonorrhoea reinfection in heterosexual STD clinic attendees: longitudinal analysis of risks for first reinfection, **124**
- reproductive technology**, View I: Assisted conception in couples with HIV infection, **185**
View 2: Assisted conception in couples with HIV infection, **189**
- reproductive tract infections**, Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- risk behaviour**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- risk behaviours**, Avoiding risky sex partners: perception of partners' risks v partners' self reported risks, **197**
- risk factors**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- risk taking**, National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- Russian Federation**, Congenital syphilis in the Russian Federation: magnitude, determinants, and consequences, **106**
- sampling**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- screening**, A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
Chlamydia screening in the United Kingdom, **3**
- Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- Recent pilot studies of chlamydia screening, **352**
- Repeat chlamydia screening by mail, San Francisco, **28**
- Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- Spending money to save money, **4**
- The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- Validity and utility of screening tests for STIs, **356**
- Who is being tested for genital chlamydia in primary care?, **234**
- self collected urine samples**, Improving response rates for self collected urine samples, **346**
- self obtained samples**, Self obtained samples for genitourinary infections, **171**
- sensitivity**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- seroepidemiological study**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- serology**, Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- seroprevalence**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
- serotesting**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- sex**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
Sexual behaviour of lesbians and bisexual women, **147**
Who is being tested for genital chlamydia in primary care?, **234**
- sex behaviour**, Sexual behaviour of lesbians and bisexual women, **147**
- sex differences**, Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- sex on premises venues**, A novel research approach in sex on premises venues (SOPV): objective measure of sexual behaviour and low level intrusion to patrons, **78**
- sex resort**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- sex workers**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- sexual abuse**, Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- sexual assault**, Follow up of patients who have been recently sexually assaulted, **349**

- sexual behaviour**, A novel research approach in sex on premises venues (SOPV): objective measure of sexual behaviour and low level intrusion to patrons, **78**
- Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- sexual health**, Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- Sexual health—health of the nation, **85**
- sexual health services**, Staff training in integrated sexual health services, **354**
- sexual risk**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- sexual risk factors**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- sexual transmission**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- sexually transmitted disease**, Chancroid: clinical manifestations, diagnosis, and management, **68**
- sexually transmitted diseases**, Avoiding risky sex partners: perception of partners' risks v partners' self reported risks, **197**
- Determinants of disclosure of genital herpes to partners, **42**
- Geographical focusing: an intervention to address increased risk for sexually transmitted diseases during repatriation and resettlement in post-war Mozambique, **77**
- Tetracycline treatment does not eradicate *Mycoplasma genitalium*, **318**
- Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
- Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- sexually transmitted infection**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Sexually Transmitted Infections**, Taking up the reins, **2**
- sexually transmitted infections**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, **332**
- Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Sexual health—health of the nation, **85**
- Sexually transmitted infections in Africa: single dose treatment is now affordable, **432**
- STI services in the United Kingdom: a way forward, **439**
- STI services in the United Kingdom: how shall we cope?, **435**
- Surveillance of sexually transmitted infections in primary care, **174**
- Treating sexually transmitted infections in primary care: a missed opportunity?, **134**
- Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- Validity and utility of screening tests for STIs, **356**
- Sexually Transmitted Infections Foundation**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- shared care**, Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- South Africa**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- STI case management at a South African teaching hospital, **498**
- specificity**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- Sri Lanka**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- staff training**, Staff training in integrated sexual health services, **354**
- STD**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- STD clinic**, Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- STD clinics**, Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- STD intervention trials**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- STD prevention**, Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- STI policies**, A survey of STI policies and programmes in Europe, **171**
- STI prevalence**, Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- STIs**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- Chaperoning in genitourinary medicine clinics, **422**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- streptococcal infections**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**
- superior mesenteric artery syndrome**, Superior mesenteric artery syndrome in an HIV positive patient, **262**
- survey**, A survey of STI policies and programmes in Europe, **171**
- survey method**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- susceptibility**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
- syndromic diagnosis**, Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- syphilis**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- Perforating chancre: any cause-effect relation with HIV infection?, **262**
- Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- Syphilis prevalence has rapidly decreased in South Korea, **323**
- Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, **351**
- Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- Tanzania**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- telephone survey**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- television**, Television coverage of AIDS and transmission of the AIDS virus, **170**
- termination of pregnancy**, Chlamydia testing before termination of pregnancy, **352**
- test of cure**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**

- tinidazole**, Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
- treatment interruption**, Enhanced risk of HIV sexual transmission during structured treatment interruption, **74**
- Trichomonas vaginalis***, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
- Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
- Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
- Was the Papanicolaou smear responsible for the decline of *Trichomonas vaginalis*?, **263**
- trichomoniasis**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- tuberculosis**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- tuberculous meningitis**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- type specific serology**, The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- Ukraine**, The HIV/AIDS epidemic in Ukraine: stable or still exploding?, **263**
- United Kingdom**, Sexual health—health of the nation, **85**
- STI services in the United Kingdom: a way forward, **439**
- STI services in the United Kingdom: how shall we cope?, **435**
- urine**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- urine testing**, Chlamydia screening in the United Kingdom, **3**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- vaccination**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- vaginitis**, Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
- varicella zoster virus**, Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
- VDRL test**, Syphilis prevalence has rapidly decreased in South Korea, **323**
- virus-like particles**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, **191**
- vulvovaginal yeasts**, Vulvovaginal carriage of yeasts other than *Candida albicans*, **249**
- women**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
- Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area