

The serious threat of multidrug-resistant and untreatable gonorrhoea: the pressing need for global action to control the spread of antimicrobial resistance, and mitigate the impact on sexual and reproductive health



EDITOR'S
CHOICE

Francis Ndowa,^{1,2} Manjula Lusti-Narasimhan,³
Magnus Unemo⁴

STIs remain a major cause of morbidity and mortality worldwide. The WHO estimated that >498 million new cases of syphilis, gonorrhoea, chlamydial infection and trichomoniasis occurred globally in 2008 in adults (15–49 years of age). More than 106 million were gonococcal infections.¹ Accordingly, gonorrhoea remains a major global public health problem with serious health, social and economic consequences. Inadequate control and treatment can result in severe complications, such as pelvic inflammatory disease, leading to infertility or ectopic pregnancy, and neonatal eye infections which can cause blindness. Up to 10% of women who remain untreated (or inadequately treated) for chlamydial and gonococcal infections may become infertile. Urogenital gonorrhoea is also asymptomatic in at least 50% of women, which leaves the infections undetected and untreated with the risk of severe complications. On a global scale, up to 4000 newborn babies, annually, may become

blind because of gonococcal and chlamydial ophthalmia neonatorum.² In men, gonococcal infections cause symptomatic urethritis which, if untreated, can result in epididymitis in 10%–30% of cases,^{w1} that may lead to infertility. Gonorrhoea also significantly enhances the transmission of HIV. The burden of gonococcal infections is highest in countries that are least able to manage them because of several factors, including stigma, inadequate resources and lack of appropriate diagnostics, surveillance and antimicrobial treatment. The problem is further compounded by the ability of *Neisseria gonorrhoeae* to develop antimicrobial resistance (AMR) within a relatively short time span.^{3 w2 w3}

Historically, since the advent of the antibiotic era, gonococcal infections have been easy to treat. However, the susceptibility of *N gonorrhoeae* to sulphonamides and penicillins from the mid-1930s and mid-1940s, respectively, gradually eroded as the organism developed a number of mechanisms for antimicrobial resistance. The mid-1980s and early 1990s were promising with the introduction of effective single-dose treatments that could be administered orally, such as fluoroquinolones (mainly ciprofloxacin and ofloxacin) and oral third-generation cephalosporins (cefixime being the most potent one). The macrolide azithromycin was even more exciting as it could treat both gonococcal and chlamydial infections with a single oral dose regimen. However, resistance to fluoroquinolones emerged only after about a decade, and these

were abandoned as first-line empiric treatment for gonorrhoea in the Asia-Pacific region already in the mid-to-late 1990s and, subsequently, in the USA, Europe and parts of Africa.^{3 w2 w3–w6} Azithromycin-resistant *N gonorrhoeae* emerged in the mid-to-late 1990s,^{4 w7} and, subsequently, also high-level azithromycin resistance has been described from several countries.^{5 w8–w11} Thus, since the early 2000s, third-generation cephalosporins (mainly cefixime and ceftriaxone) have been the sole class of antimicrobials recommended as first-line empiric treatment.^{3 w2 w3}

Worryingly, resistance and treatment failures to cefixime have been verified in Japan⁶ and recently in Europe.^{7–9} The recent report of a strain of *N gonorrhoeae* in Japan that was highly resistant to the parenteral ceftriaxone, and associated with a probable treatment failure with ceftriaxone,¹⁰ the last remaining option for empiric treatment, sounded alarm bells of significant future challenges to the treatment and control of gonococcal infections and their complications. This was also followed by the identification of a highly ceftriaxone-resistant strain in France¹¹ and in Spain.¹² Furthermore, many regions, worldwide, describe a decreasing susceptibility to ceftriaxone, and reports of ceftriaxone treatment failure of gonococcal pharyngeal infections have been published.^{3 13 w2 w3} Given the ability of *N gonorrhoeae* to develop AMR within a relatively short time span compounded with its ability to retain the resistance to previous antibiotics, even after their use has been discontinued, the threat of a widespread ceftriaxone resistance and untreatable gonorrhoea in certain circumstances is real.^{3 10 11 w2 w3}

The global trend of AMR in *N gonorrhoeae*, and decreased susceptibility and resistance to the third-generation cephalosporins, in the age of easy international travel dictated that a global approach needed to be elaborated to respond to the emerging menace of untreatable gonorrhoea. This was especially important because gonococcal AMR data were not available in many settings, and the true global magnitude of the problem is unknown. Therefore, the WHO Global Gonococcal Antimicrobial Surveillance Programme (WHO Global GASP), initially established in the 1990s, was revisited and revamped in 2009 as a network of collaborators in order to enable a global, coordinated response. The WHO Global GASP network consists of the WHO Headquarters, Geneva, as the managing focal centre working with interregional collaborators in Africa, Asia, the Americas, Europe and the

¹Department of Reproductive Health and Research, World Health Organization, Geneva, Switzerland; ²STI Specialist and Consultant, Harare, Zimbabwe;

³Department of Reproductive Health and Research, World Health Organization, Geneva, Switzerland; ⁴World Health Organization Collaborating Centre for Gonorrhoea and other STIs, National Reference Laboratory for Pathogenic *Neisseria*, Örebro University Hospital, Örebro, Sweden

Correspondence to Dr Manjula Lusti-Narasimhan, Department of Reproductive Health and Research, World Health Organization, 20 Avenue Appia, CH-1211 Geneva 27, Switzerland; lustinarasimhanm@who.int

Western Pacific, and aims to encourage and recruit national laboratories to monitor gonococcal AMR data, with particular attention to third-generation cephalosporins, and identify treatment failures using recommended treatment.

To facilitate the functioning of the GASP network and engage the support of governments and the international community and donors in this venture, the WHO published the 'Global Action Plan to Control the Spread and Impact of Antimicrobial Resistance in *N gonorrhoeae*' in June 2012.¹⁴ The Global Action Plan aims to facilitate early detection of strains resistant to recommended treatment, combined with a public health response to mitigate the impact of cephalosporin-resistant *N gonorrhoeae* on sexual and reproductive health morbidity. The WHO Global Action Plan calls for the public health approach to the control of gonococcal infections and the emergence of multidrug-resistant and untreatable gonorrhoea. Some of the main components and actions of the Global Action Plan are:

- ▶ Advocacy for increased awareness on correct use of antibiotics among healthcare providers and the consumers, particularly in key populations, including men-who-have-sex-with-men and sex workers.
- ▶ Effective prevention, diagnosis and control of gonorrhoea, using prevention messages, interventions, recommended diagnosis and treatment regimens.
- ▶ Systematic monitoring and early detection of treatment failures with recommended treatment (cefixime and ceftriaxone) by developing a standard case definition of treatment failure, and protocols for verification, reporting and management of failure.
- ▶ Effective drug regulations and prescription policies.
- ▶ Strengthened AMR surveillance, especially in countries with a high burden of gonococcal infections (and/or gonococcal AMR), other STIs and HIV.
- ▶ Capacity building to establish regional networks of laboratories to perform quality-assured gonococcal culture and antimicrobial susceptibility testing.
- ▶ Research to identify new molecular methods for detecting and monitoring antimicrobial resistance.
- ▶ Research (basic science, in vitro and clinical studies) to identify alternative

effective treatment strategies and/or antimicrobials (or other effective compounds) for gonorrhoea.

This Global Action Plan will be implemented within the WHO's 'Global Strategy for the prevention and control of sexually transmitted infections: 2006–2015' which was adopted by the WHO Member States at the 59th World Health Assembly in May 2006.² To ensure harmonisation and sustainability, the WHO GASP network will be linked to, and operate within, the WHO Policy Package to Combat AMR which was launched on World Health Day 2011 in Geneva.¹⁵

The implementation of the WHO GASP activities both at the global and national levels can only be achieved with the formation of partnerships for funding and training among different but related programmes, and adequate funding for surveillance of STIs, in general, and quality-assured gonococcal AMR testing, in particular. Several region-specific and even country-specific action plans to control the spread of gonococcal antimicrobial resistance, are also under preparation, these plans are also crucial for the global response to the serious threat of untreatable gonorrhoea. Ultimately, it is essential to identify new strategies and/or antimicrobials (or other effective compounds) for effective treatment of gonorrhoea. It is imperative that this research is adequately supported and funded by national and international partners.

Acknowledgements We are grateful to all the WHO colleagues, the WHO GASP members and other international experts for their valuable contribution in developing the WHO Global Action Plan to Control the Spread and Impact of Antimicrobial Resistance in *N gonorrhoeae*.

▶ Additional references are published online only. To view these references please visit the journal online (<http://sti.bmj.com/content/88/5.toc>).

Contributors FN, MLN and MU collaboratively wrote the present Editorial.

Competing interests None.

Provenance and peer review Commissioned; internally peer reviewed.

Accepted 10 June 2012

Sex Transm Infect 2012;**88**:317–318.
doi:10.1136/sextrans-2012-050674

REFERENCES

1. **World Health Organization.** *Global Prevalence and Incidence Of Selected Curable Sexually Transmitted*

Infections—2008: Overview and Estimates. Geneva, 2012. http://www.who.int/reproductivehealth/publications/rtis/2008_STI_estimates.pdf (accessed 6 Jul 2012).

2. **World Health Organization.** *Global Strategy For The Prevention And Control of Sexually Transmitted Infections: 2006–2015.* Geneva, 2007. ISBN: 978 92 4 156347 5. <http://www.who.int/reproductivehealth/publications/rtis/9789241563475> (accessed 28 May 2012).
3. **Tapsall JW, Ndowa F, Lewis DA, et al.** Meeting the public health challenge of multidrug- and extensively drug-resistant *Neisseria gonorrhoeae*. *Expert Rev Anti Infect Ther* 2009;**7**:821–34.
4. **Dillon JA, Ruben M, Li H, et al.** Challenges in the control of gonorrhoea in South America and the Caribbean: monitoring the development of resistance to antibiotics. *Sex Transm Dis* 2006;**33**:87–95.
5. **Palmer HM, Young H, Winter A, et al.** Emergence and spread of azithromycin-resistant *Neisseria gonorrhoeae* in Scotland. *J Antimicrob Chemother* 2008;**62**:490–4.
6. **Yokoi S, Deguchi T, Ozawa T, et al.** Threat to cefixime treatment of gonorrhoea. *Emerg Infect Dis* 2007;**13**:1275–7.
7. **Ison CA, Hussey J, Sankar KN, et al.** Gonorrhoea treatment failures to cefixime and azithromycin in England, 2010. *Euro Surveill* 2011;**16**:19833.
8. **Unemo M, Golparian D, Syversen G, et al.** Two cases of verified clinical failures using internationally recommended first-line cefixime for gonorrhoea treatment, Norway, 2010. *Euro Surveill* 2010;**15**:19721.
9. **Unemo M, Golparian D, Sary A, et al.** First *Neisseria gonorrhoeae* strain with resistance to cefixime causing gonorrhoea treatment failure in Austria, 2011. *Euro Surveill* 2011;**16**:19998.
10. **Ohnishi M, Golparian D, Shimuta K, et al.** Is *Neisseria gonorrhoeae* initiating a future era of untreatable gonorrhoea? Detailed characterization of the first strain with high-level resistance to ceftriaxone. *Antimicrob Agents Chemother* 2011;**55**:3538–45.
11. **Unemo M, Golparian D, Nicholas R, et al.** High-level cefixime- and ceftriaxone-resistant *N gonorrhoeae* in France: novel penA mosaic allele in a successful international clone causes treatment failure. *Antimicrob Agents Chemother* 2012;**56**:1273–80.
12. **Cámara J, Serra J, Ayats J, et al.** Molecular characterization of two high-level ceftriaxone-resistant *Neisseria gonorrhoeae* isolates detected in Catalonia, Spain. *J Antimicrob Chemother.* Published Online First: 7 May 2012. doi:10.1093/jac/dks162
13. **Unemo M, Golparian D, Hestner A.** Ceftriaxone treatment failure of pharyngeal gonorrhoea verified by international recommendations, Sweden, July 2010. *Euro Surveill* 2011;**16**:19792.
14. **World health Organization.** Global action plan to control the spread and impact of antimicrobial resistance in *Neisseria gonorrhoeae*. 2012. ISBN: 978 92 4 150350 1. <http://www.who.int/reproductivehealth/publications/rtis/9789241503501> (accessed 28 May 2012).
15. **World Health Day.** *Policy Briefs/WHO Policy Package to Combat Antimicrobial Resistance.* 2011. <http://www.who.int/world-health-day/2011/policybriefs/en/index.html> (accessed 23 May 2012).

STI

The serious threat of multidrug-resistant and untreatable gonorrhoea: the pressing need for global action to control the spread of antimicrobial resistance, and mitigate the impact on sexual and reproductive health

Francis Ndowa, Manjula Lusti-Narasimhan and Magnus Unemo

Sex Transm Infect 2012 88: 317-318
doi: 10.1136/sextrans-2012-050674

Updated information and services can be found at:
<http://sti.bmj.com/content/88/5/317>

These include:

Supplementary Material

Supplementary material can be found at:
<http://sti.bmj.com/content/suppl/2012/07/13/sextrans-2012-050674.DC1>
<http://sti.bmj.com/content/suppl/2012/10/29/sextrans-2012-050674.DC2>

References

This article cites 10 articles, 3 of which you can access for free at:
<http://sti.bmj.com/content/88/5/317#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections
[Editor's choice](#) (79)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>