

The model is used to assess the 20 year impact (on HIV incidence/prevalence) of scaling up early ART to PWID, compared to a baseline of current ART access (20% of HIV+ individuals with CD4 < 350 cells/ μ L are recruited onto ART annually). An uncertainty analysis was undertaken using posterior model fits to consider which PWID behavioural/epidemiological factors affect impact.

Results Only small reductions in PWID HIV incidence/prevalence (< 10% over 20 years) will occur with current ART provision in Manipur. If current ART recruitment rates are maintained, but all HIV+ PWID become eligible for ART, then a median 12%/11% relative decrease in HIV incidence/prevalence occurs over 20 years compared to baseline. This increases to 22%/20% or 33%/28% if 40% or 80% of HIV+ PWID are recruited per year, respectively. If the ART LTFU amongst PWID is halved to 7.5% per year then impact increases by half. Uncertainty analyses suggest the impact achieved through scaling-up ART is highly dependent on baseline HIV prevalence and the cofactor increase in HIV infectivity during early infection, with less impact occurring for higher HIV prevalences and cofactors. The injecting cessation rate and level of mixing between risk groups are also important.

Discussion HIV treatment could result in large reductions in PWID HIV incidence, but is unlikely to result in local elimination except in low HIV prevalence settings.

S04.3 ANTIRETROVIRAL THERAPY AMONG FEMALE SEX WORKERS IN BURKINA FASO: CURRENT SITUATION

doi:10.1136/sextrans-2013-051184.0027

¹I Konate, ¹A Ouédraogo, ¹W W Bazié, ²T Traoré, ¹A Sanon, ¹C Huet, ^{1,2}N Méda, ³P Mayaud, ⁴P Van De Perre, ⁴N Nagot. ¹UR_SR/VIH/TB, Centre MURAZ, Bobo-Dioulasso, Burkina Faso; ²Centre de Recherche International en Santé/UO, Ouagadougou, Burkina Faso; ³LSHTM, London, UK; ⁴Université de Montpellier, INSERM U 1058, Montpellier, France

Background In Burkina Faso, female sex workers (FSW) are a core group in HIV transmission with 16.5% cases of HIV-infection, vs. 1.0% in the general population. Provision of ART to this core group may be key to controlling the HIV epidemic. We aimed at documenting the current situation regarding ART provision to FSW in Burkina Faso.

Methods We reviewed the interventions of the national HIV/AIDS control programme (NACP) regarding ART provision to FSW, using grey literature and interviews of stakeholders. In addition, we reviewed the findings of the local research on this topic.

Results Before 2010, programmes targeting FSW in Burkina Faso focused only on prevention, mainly through local NGOs. From 2010, in each of the 13 regions, the NACP designated one primary health centre to care for FSW. No statistics are available yet from these centres, but NACP/MOH stakeholders admitted that few FSW adhere to it, mainly because of a lack of strategy to reach and retain FSW, and lack of health care workers training. The ANRS 1222 ‘Yerelon’ research programme tested a FSW peer-led intervention integrating prevention and care activities within one dedicated centre in Bobo-Dioulasso. Beside care providers, peers and a psychologist much contributed to support ART adherence before and after initiation. Preliminary data among the 47 FSW treated with ART showed high adherence to ART (> 95%) at 12 months (92%) and 36 months (100%). At 36 months, plasma viral load was undetectable in 82% of FSW, with a mean CD4 increase of 230 cells/ μ L (IQR: 90–400).

Conclusion Despite a specific programme, ART provision to FSW remains limited in Burkina Faso. The ‘Yerelon’ model of prevention and care intervention can serve as a strategy for the NACP FSW programme, and the research sites be used as training centres to improve ART provision coverage and efficiency.

S04.4 ART AMONG FSW IN SOUTH AFRICA

doi:10.1136/sextrans-2013-051184.0028

F Venter. *Wits Reproductive Health and HIV Institute (WRHI), Johannesburg, South Africa*

Until recently South Africa has targeted its HIV interventions towards the general population, but the National Strategic Plan of 2012–2016 has expanded this focus to include key populations with high HIV prevalence and incidence. Foremost in this group are sex workers who are identified in the 2011 South African Know Your Epidemic (KYE) as contributing as much as 20% of the HIV transmission burden in the country. A model of health care delivery through an inner city sex worker programme has steadily expanded over a decade from special services delivered in a clinic, to clinical outreach services delivered within brothels and a mobile clinic, that now includes ART. The intervention includes behaviour change communication delivered through peer educators and outreach staff, programme-supported free clinical services to detect and treat STIs, HIV counselling and testing (HCT), TB detection, contraceptive provision, male and female condom provision. Referral linkages to relevant state services including HIV and TB treatment services and reproductive health services are in place with outreach workers tracing defaulting patients. We will discuss the outcomes of community based ART for sex workers, the challenges of provision, and present outcomes including virological suppression and loss to follow up data.

S04.5 ART AMONG FSW IN INDIA

doi:10.1136/sextrans-2013-051184.0029

S Mishra, *M Boily. Imperial College, London, UK*

In India, the provision of antiretroviral treatment (ART) began in 2004. Around the same time, a large-scale, condom-based HIV prevention programme targeted to FSWs was implemented in several districts. In this talk, we present findings from a comprehensive review of available data exploring trends in the ART ‘care-cascade’ among FSWs in India, with a focus on routine HIV testing, linkage to pre-ART care, ART uptake, and retention on ART. We then examine the association between local condom-based targeted HIV prevention programmes and HIV testing among FSWs, and identify key data gaps in the ‘care cascade’ among FSWs. We conclude by exploring the interaction between turn-over in sex work and ART eligibility among FSWs, and discuss the potential implications for local ART programmes.

S04.6 EXAMINING THE POPULATION-LEVEL IMPACT OF SCALING-UP ART FOR FSWs ACROSS EPIDEMIC CONTEXT

doi:10.1136/sextrans-2013-051184.0030

M Boily, *S Mishra. Imperial College, London, UK*

In this presentation, we will summarise the results of a recent systematic review of the published literature on the key parameters of the treatment cascade among female sex workers. Using different examples from transmission dynamics modelling studies of HIV transmission and ART use, we will discuss the potential impact of scaling-up ART for FSWs in contrasting epidemiological settings. We will present results of the impact of ART programmes in settings with different HIV prevalence levels in the general population (very low, low/medium, high) and different history of condom use interventions and level of ART scale-up already achieved. For example, we will compare the impact of different ART scale-up scenarios and eligibility criteria in districts of India (low prevalence settings) where there has been and there has not been a large-scale condom use intervention implemented in recent years. Other settings with