

Methods From January 2010 to November 2012, all females and MSM were routinely tested for anorectal, oropharyngeal and urogenital *Chlamydia trachomatis* (Ct) and *Neisseria gonorrhoeae* (Ng). Data were collected on demographics and sexual behaviour.

Results The study population comprised 4829 consults. Overall Ct positivity was 6.8% (131/1937) in females and 10.0% (289/2892) in MSM, for Ng this was 4.3% (n = 83) and 6.7% (n = 193) respectively. The proportion isolated anorectal Ct was 16.8% (n = 22) in females and 63.0% (n = 182) in MSM, for Ng this was 10.8% (n = 9) and 44.0% (n = 85) respectively. Combined urogenital and anorectal Ct was 45.0% (n = 59) in females and 13.5% (n = 39) in MSM, for Ng this was 19.3% (n = 16) and 11.4% (n = 22) respectively. Isolated oropharyngeal Ct was 9.9% (n = 13) in females and 5.9% (n = 17) in MSM, for Ng this was 53.0% (n = 44) and 31.6% (n = 61) respectively. Isolated urogenital Ct was 28.2% (n = 37) in females and 17.6% (n = 51) in MSM, for Ng this was 16.9% (n = 14) and 13.0% (n = 25) respectively.

Discussion Anorectal and oropharyngeal STD were often isolated and thereby not coincidentally treated with urogenital STD. Therefore urogenital testing only is insufficient. Furthermore, these isolated infections remain undiagnosed if routine anorectal/oropharyngeal testing is lacking. Combined anorectal/urogenital Ct/Ng were also highly prevalent, highlighting the importance of an effective treatment for simultaneous genital and extragenital Ct/Ng.

P3.155 ASSESSMENT OF KNOWLEDGE OF SEXUAL TRANSMITTED DISEASES (STDs) AND A SEROPREVALENCE STUDY IN GAY MAN IN KAYSERI, TURKEY

doi:10.1136/sextrans-2013-051184.0613

¹E Balci, ²M Doganay, ²H Demiraslan, ³S Gokahmetoglu. ¹Erciyes University Medical Faculty Department of Public Health, Kayseri, Turkey; ²Erciyes University Medical Faculty Department of Infectious Diseases, Kayseri, Turkey; ³Erciyes University Medical Faculty Department of Microbiology, Kayseri, Turkey

Background Homosexual behaviour is prone to many sexual transmitted diseases (STDs) and many social problems. The aim of this study is to assess the knowledge, source, the reliability and accuracy of information resources and to assess the seropositivity for some STDs.

Method A questionnaire was prepared for the STDs including these; to learn their demographic and economical characteristics, having any chronic disease or, experienced any disease, use of any illegal drug, their sense for acceptability in public, number of partner in one year, main transmission route, prevention and source of their information about this diseases in gay people. In additional these, Another questions were to learn their to join an education programme for prevention of STDs and to give blood samples for the serologic investigation against hepatitis B, hepatitis C, syphilis and HIV. Voluntary people defined himself as gay, was entered to the study.

Results The questionnaire was replied by 94 people. Mean age was 23.32 ± 4.6 year and 80.4% defined himself as gay. Of these, 72.6% was graduated from high school or university, 5.9% had a chronic disease and 3.9% had receiving any medicine. 29.4% had no any health insurance. Using one of these; cigarette, alcohol, heroin or cocaine were at percentage of 62.8, 46.8, 10.6 and 1.1 respectively. 23.4% had multiple partners and 76.7% single partners. Majority people knew the transmission of HIV/AIDS by sexual intercourse (88.3%) or blood/blood products (77.7%). 69.1% had a history of unprotected sexual relation. Major source of their knowledge about STDs was internet or other media. Blood samples could be taken from 39, one was positive for HIV. An education programme was used as a small group after questionnaire.

Conclusion This is a first basic study on homosexual group, summarising their problems in our region.

P3.156 CONTRIBUTION OF SEXUAL PRACTISES OTHER THAN ANAL SEX TO THE TRANSMISSION OF BACTERIAL SEXUALLY TRANSMITTED INFECTIONS IN MEN WHO HAVE SEX WITH MEN

doi:10.1136/sextrans-2013-051184.0614

^{1,2,3}J L Nash, ^{1,2}J S Hocking, ^{1,2}T R H Read, ^{1,2}M Y Chen, ^{1,2,3}C S Bradshaw, ^{1,2}D S Forcey, ^{1,2}C K Fairley. ¹Melbourne Sexual Health Centre, Melbourne, Australia; ²Sexual Health Unit, School of Population Health, University of Melbourne, Melbourne, Australia; ³Monash University, Melbourne, Australia

Background Syphilis, chlamydia and gonorrhoea are increasingly common amongst Australian men who have sex with men (MSM). Our objective was to quantify the proportion of cases and risk of primary syphilis (PS), urethral chlamydia (UCT) and urethral gonorrhoea (UGC) attributable to sexual practises other than anal sex.

Methods Electronic medical records for MSM who attended Melbourne Sexual Health Centre between July 2002 (for PS) or January 2006 (for UCT and UGC) and October 2012 inclusive were examined.

Results There were 37 533 eligible consultations; 2374 (6%) of these reported no anal sex. There were 204 PS diagnoses, 673 UCT diagnoses, and 618 UGC diagnoses; 12 (6%), 16 (2%) and 44 (7%) cases respectively occurred in consultations where no anal sex was reported in the previous three months (PS, UGC) or twelve months (UCT). Amongst MSM who reported no anal sex, PS was diagnosed in 0.5 cases/100 consultations, UCT was diagnosed in 1.5 cases/100 tests for UCT and UGC was diagnosed in 14 cases/100 tests for UGC. UCT was significantly more common in MSM who reported anal sex (OR 2.18, 95% CI 1.32 – 3.59, p = 0.002), but PS (p = 0.82) and UGC (p = 0.14) were not. For MSM who reported anal sex, condom use was protective for all three infections (all p ≤ 0.03).

Conclusions UCT rarely occurs in MSM due to sexual practises other than anal sex; however these practises appear to contribute significantly to the acquisition of PS and UGC. Our figures probably underestimate the risk involved, as men who have anal sex also engage in other sexual practises. The implication is that successful programmes to control sexually transmitted infections, particularly PS and UGC, will need to utilise additional strategies, such as frequent testing, as well as promoting condom use.

P3.157 INCREASING STI RATES IN YOUNG MSM IN THE NETHERLANDS, 2006–2011

doi:10.1136/sextrans-2013-051184.0615

¹F D H Koedijk, ¹B H van Benthem, ²E M D C Vrolings, ^{1,3}M A B van der Sande, ⁴On behalf of the Dutch STI clinics. ¹National Institute of Public Health and the Environment, Bilthoven, The Netherlands; ²STI AIDS The Netherlands, Amsterdam, The Netherlands; ³Julius Center for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht, The Netherlands; ⁴A van Daal (East), AP van Leeuwen (North-Holland Flevoland), F de Groot (North), CJA Hoebe (Limburg), M Langevoort (Utrecht), AM van Camerijk (South-Holland North), J van de Sande (Zeeland-Brabant), V Wieërs (South-Holland South), Bilthoven, The Netherlands

Objective Young men having sex with men (MSM) are a high-risk group involved in on-going transmission of sexually transmitted infections (STI). To improve our understanding of this epidemic, distribution of STI and testing behaviour in young MSM attending STI clinics in the Netherlands was analysed.

Methods National STI surveillance data on young MSM (15–24 years) for 2006–2011 were analysed. STI included were chlamydia, gonorrhoea, infectious syphilis and HIV. Chi-square test for trend was performed to analyse time trends in testing behaviour and STI positivity rates, multivariable logistic regression was used to identify factors associated with STI positivity.

Results Number of consultations in young MSM increased from 1161 in 2006 to 3167 in 2011 (+166%). The proportion not aware of