

Further studies are needed on utilisation and linkage to care and prevention services.

P17.03 CONVENIENT HIV TESTING SERVICE MODELS COMBINED WITH RAPID TESTING ARE ATTRACTING PREVIOUSLY UNTESTED GAY AND BISEXUAL MEN

^{1,2}V Knight*, ²H Wand, ³J Gray, ²P Keen, ^{1,4}A McNulty, ²R Guy. ¹Sydney Sexual Health Centre, South East Sydney Local Health District, Sydney, 2000, Australia; ²The Kirby Institute, University of New South Wales, Sydney, NSW, 2052, Australia; ³ACON Health, 414 Elizabeth Street Surry Hills, Sydney, NSW, 2010, Australia; ⁴School of Public Health and Community Medicine, University of NSW, Kensington, NSW, 2052, Australia

10.1136/sextrans-2015-052270.581

Introduction HIV testing is a cornerstone of the treatment as prevention (TasP) approach. In Australia, gay and bisexual men (GBM) account for the majority of HIV diagnoses each year, but less than a quarter are testing at the recommended frequency. In response, a range of new service delivery models were introduced in Sydney over the past few years. We assessed which HIV testing service delivery models were more likely to attract GBM who had never previously tested or were testing infrequently.

Methods We compared demographics, risk behaviour and HIV testing history among new GBM clients attending three different HIV testing service models (fast-track Xpress clinic, fixed-site community-based service and time-limited community-based shopfront) between August 2013 and May 2014. All services offered HIV rapid testing. We used multivariate regression to assess factors (including service model) associated with being untested or infrequent testers (not tested within the past 12 months).

Results Overall, 1704 new GBM attended the services; 19% were untested and 41% were infrequent testers. Across the services, there were significant differences in demographics, risk behaviour and past HIV testing history. The overall HIV seropositivity was 1.2% (95% CI: 0.8%–1.9%) and STI positivity was 12.4% (95% CI: 11.6–17.2) with no significant differences across services. Factors independently associated with being untested were attendance at the two community sites, younger age, being born in Asia, living in North Sydney, being bisexual and reporting fewer male sexual partners. Factors independently associated with infrequent testers were attending the fast track Xpress clinic, being older, being born in Asia and reporting fewer male partners.

Conclusion The findings show the two community sites reached more untested men and the fast-track clinic model more infrequent testers but the HIV/STI diagnosis rate was consistent across services, indicating that all three testing models are important to increase HIV testing among GBM.

Disclosure of interest statement Vickie Knight is supported by an Australian Postgraduate Association scholarship. No pharmaceutical grants were received in the development of this study.

P17.04 THE RELATIONSHIP BETWEEN HIV TESTING FREQUENCY AND HIV RISK PERCEPTION AMONG PERUVIAN MSM AND TRANSGENDER WOMEN

¹Sk Vargas*, ^{1,2}KA Konda, ¹SR Leon, ³G Calvo, ⁴HJ Salvatierra, ⁵B Brown, ²JD Klausner, ¹CF Caceres. ¹Unit of Health, Sexuality and Human Development, and Laboratory of Sexual Health, Universidad Peruana Cayetano Heredia, Lima, Peru; ²University of California, Los Angeles, CA, USA; ³Epicentro, Lima, Peru; ⁴Alberto Barton Health Center, Health Directorate of Callao, Lima, Peru; ⁵University of California, Irvine, CA, USA

10.1136/sextrans-2015-052270.582

Introduction HIV in Peru is concentrated among men who have sex with men (MSM) and transgender women (TW). Over 70% of HIV+ MSM/TW ignore their status, as HIV testing frequency is low, delaying diagnosis and treatment. We aimed to assess the relationship between HIV testing frequency and perceived HIV risk among MSM/TW.

Methods This analysis included baseline data from MSM/TW who reported a negative or unknown HIV status from a STI clinic cohort of MSM/TW at high risk in Lima. The behavioural survey assessed HIV risk perception (high, moderate, low) and prior testing frequency (testing was defined as frequent if it had occurred at least biannually). For HIV diagnosis, we used a 3rd generation rapid test, a 4th generation EIA, and Western Blot confirmation. Chi-square tests and multivariable regression were used to estimate adjusted prevalence ratios (aPRs).

Results Among eligible subjects (243 MSM and 67 TW), 122 (39%) reported frequent prior HIV testing, while HIV risk was perceived as high, moderate and low by 72 (23%), 148 (48%) and, 90 (29%) participants, respectively. Frequent HIV testing was more prevalent among participants reporting low/no HIV risk (aPR: 1.62; CI: 1.20–2.20) and among those reporting previous syphilis infection (aPR: 1.58; CI: 1.22–2.04); but was less prevalent among participants reporting recent condomless anal intercourse (aPR: 0.67; CI: 0.51–0.90). Baseline HIV status and HIV risk perception were not associated: HIV prevalence was 11%, 13% and 10% among those reporting high, moderate and low/no HIV risk (chi-square p-value >0.1).

Conclusion Frequent HIV testing was more prevalent among MSM/TW reporting low/no HIV risk. Risk perception did not correlate with HIV status among previously undiagnosed subjects. WHO-recommended biannual HIV testing should be promoted, and factors potentially affecting risk perception and testing frequency (e.g. access, stigma) should be identified and addressed by HIV programming for MSM/TW in Peru.

P17.05 HIV TEST REFUSALS AMONG BLACK AFRICANS ATTENDING SEXUALLY TRANSMITTED INFECTION CLINICS IN ENGLAND, 2013

H Mohammed*, M Furegato, A Nardone, G Hughes. *HIV & STI Department, Public Health UK*

10.1136/sextrans-2015-052270.583

Introduction Black Africans comprise the second largest group affected by HIV in England, but it is estimated that 34% are

undiagnosed. We identified factors associated with HIV test refusals among Black African sexually transmitted infection (STI) clinic attendees.

Methods Data on all STI clinic attendances in England in 2013 were obtained from the genitourinary medicine clinic activity dataset (GUMCADv2). Individuals diagnosed with HIV on or before 31st December 2012 were excluded from analysis. Ethnic variations in HIV test refusal at an attendance were determined. Further analyses were restricted to Black Africans, and bivariate and multivariate associations between demographic and clinical characteristics and HIV test refusal were assessed. All associations were determined using generalised estimating equations regression and adjusted odds ratios (aORs) with 95% confidence intervals (CIs) are reported.

Results Black Africans made 92,331 attendances at STI clinics in 2013 and refused an HIV test on 7,666 (8.3%) occasions. After adjusting for gender/sexual orientation, Black Africans were least likely to have refused an HIV test [aOR (95% CI): 0.503 (0.490–0.517)] (*vs.* White British persons). Among Black Africans, the odds of refusing an HIV test decreased with age [0.983 (0.980–0.986)] and were lower among those born outside the UK [0.676 (0.637–0.717)], those who were tested for HIV within the last year [0.802 (0.762–0.845)] and in men who have sex with men [0.307 (0.235–0.402)] (*vs.* heterosexual men and all women), while the odds were higher among those diagnosed with a new STI at the same attendance [1.272 (1.193–1.357)].

Conclusion Among Black Africans, targeted health promotion may be needed to improve HIV testing rates and decrease the proportion undiagnosed, especially for heterosexual men, women, younger, UK-born persons and those newly diagnosed with an STI.

Disclosure of interest statement There are no conflicts of interest.

P17.06 HIV PREVALENCE AND FACTORS ASSOCIATED WITH HIV TESTING AMONG YOUNG PEOPLE (15–24 YEARS) IN TANZANIA: A SECONDARY ANALYSIS OF THMIS DATA

^{1,2}DJ Damian*, ^{1,2}SE Msuya. ¹Kilimanjaro Christian Medical University College (KCMU Co), Moshi, Tanzania; ²Kilimanjaro Christian Medical Centre (KCMC), Moshi, Tanzania

10.1136/sextrans-2015-052270.584

Background HIV remained to be a major public health problem affecting the lives and livelihood of many in Sub-Saharan Africa (SSA). Young people (15–24 years) accounted for 42% of new HIV infections in people aged 15 years and older; and about 80% of them live in SSA. Knowing HIV status among young people can be one of key strategies in reducing transmission. Knowing HIV status is associated with increased safe sex practices, decreased number of partners and early/timely access to ART in HIV-positive. In Tanzania, only 54% of females and 34% of males aged 15–24 years have ever been tested HIV with limited information regarding factors associated with testing among young people.

Methods This was a secondary analysis of Tanzania HIV and Malaria Indicator Surveys data (THMIS) conducted in 2011–12. Descriptive statistics, bivariate and regression modelling of survey data were used to determine predictors of HIV testing among young people in Tanzania.

Results Prevalence of HIV among young people is 2.0% (young women 3% *vs.* young men 1%). Nearly half of all young people have ever tested for HIV (44.9%). HIV testing was associated

with older age (20–24 years) [AOR = 3.0, 95% CI: 2.4–3.6]; female gender [AOR = 2.9, 95% CI: 2.4–3.5]; currently married [AOR = 3.0, 95% CI: 2.3–3.8]; formerly married [AOR = 2.4, 95% CI: 1.6–3.6]; primary education [AOR = 1.8, 95% CI: 1.3–2.5]; secondary education [AOR = 3.5, 95% CI: 2.4–5.1]; living in urban areas [AOR = 1.4, 95% CI: 1.2–1.7] and having stigmatising attitudes towards HIV [AOR = 1.2, 95% CI: 1.0–1.4].

Conclusion HIV prevalence in young women is three times higher than in young men. In this high HI-prevalence setting almost half of young people have not tested for HIV. Provision of information and sensitisation among young people on HIV testing should focus on young people who are single, uneducated and living in rural areas.

P17.07 SEXUAL RISK BEHAVIOUR PREDICTS MORE FREQUENT USE OF HIV SELF-TESTING: EARLY FINDINGS FROM THE FORTH TRIAL

¹R Guy*, ¹MS Jamil, ²C Fairley, ¹K Smith, ¹A Grulich, ¹J Bradley, ¹J Kaldor, ²M Chen, ³A McNulty, ⁴M Holt, ⁵D Ryan, ⁶C Batrouney, ⁷D Russell, ¹B Bavinton, ¹P Keen, ¹DP Conway, ¹H Wand, ^{1,8}G Prestage, on behalf of FORTH Study Group. ¹The Kirby Institute, UNSW Australia, Sydney, Australia; ²Melbourne Sexual Health Centre, Melbourne, Australia; ³Sydney Sexual Health Centre, Sydney Hospital Sydney, Australia; ⁴Centre for Social Research in Health, UNSW Australia, Sydney, Australia; ⁵ACON, Sydney, Australia; ⁶Victorian AIDS Council, Melbourne, Australia; ⁷Cairns Sexual Health Service, Cairns, Australia; ⁸Australian Research Centre in Sex Health and Society, La Trobe University, Melbourne, Australia

10.1136/sextrans-2015-052270.585

Background Most HIV diagnoses in Australia are among gay and bisexual men (GBM), yet less than a quarter of higher-risk GBM are testing at the recommended frequency (3–6 monthly). In the context of a randomised trial of HIV self-testing (FORTH), we examined the predictors of more frequent use of HIV self-testing among higher-risk GBM.

Methods Participants in FORTH trial included higher-risk GBM (>5 sexual partners or condomless anal intercourse in the past 3 months). The trial is being conducted over 12 months, and men in the intervention arm receive 4 self-tests (OraSure's OraQuick home test) at baseline and additional self-tests on request. Using data from the baseline and 6 month surveys, we used logistic regression to examine predictors of using >2 self-tests over 6 months among participants in the intervention arm.

Results Of the GBM (n = 154) in the intervention arm, 59% reported in the baseline survey they had condomless anal intercourse with casual partners (CLAIC) in the past 6 months and 56% had a HIV test every six months. Men who reported CLAIC in the past 6 months were more likely to use >2 HIV self-tests in the first 6 months of the trial (odds ratio:2.8, 95% CI: 1.2–6.7). No other baseline survey factors were associated with >2 self-tests, including; demographics, testing frequency, likelihood to self-test in the future, and reported testing barriers (the process of getting tested is too much hassle, I don't like having to return for results, I don't want to go to a clinic/doctor to get tested).

Conclusion These findings indicate men who report sexual risk behaviour are more likely to increase their testing frequency through self-tests, which is a key HIV prevention goal. However there is also a need to ensure the longer window period of the OraQuick self-test is understood to avoid infections being missed.