



CrossMark

Highlights from this issue

doi:10.1136/sextrans-2016-052567

Jackie A Cassell, *Editor in Chief*

Syphilis, has always been known the "protean" disease, capable of manifesting in many forms and deceiving physicians as to its true nature. It confused Hunter, whose inoculation experiments¹ were the forerunner of the Tuskegee and Guatemala experiments²—it is disputed whether he experimented on himself, but not that he inoculated others in support of a unitary theory of gonorrhoea and syphilis. The huge impact of syphilis on pre-penicillin public health together with its place at the centre of moral debate, and the difficulty of diagnosis made it central to paradigm shifts in the philosophy of science as described in Fleck's seminal work "The Genesis and Development of a Scientific Fact",³ first published in German in 1935. Many developed countries have seen a huge resurgence of syphilis, particularly among men who have sex with men (MSM), though it is now generally a benign condition since we have effective treatment. This month's case report by Muldoon *et al*⁴ is a timely reminder that syphilis can be dangerous even in its earliest stages. The authors report a case of secondary syphilis presenting with coronary ostial occlusion and aortitis. It's important too that we regularly reflect on the clinical characteristics of long established conditions, which can evolve with selection pressures on their causative organisms, and depend on the curiosity and drive of clinicians to document what they are seeing. On a related note, Towns *et al*⁵ report a high frequency of multiple and painful anogenital syphilis lesions, in the absence of herpes virus infection.

A striking feature of global epidemiology is the resistance of sexual health inequalities in vulnerable ethnic groups to health improvements in the wider societies surrounding them. Though minority ethnic groups experience many dimensions of poor health, the rate ratios for poor sexual health outcomes continue to be staggering in many developed world settings. This has been shown time and time again. Recent years have seen an emphasis on structural determinants of sexual health. This month we publish a call to arms for Indigenous sexual health in Australia, by the redoubtable Kerry Arabena. Her Editorial relates primarily to Australia—where,

astonishingly, trachoma remains endemic among Indigenous peoples and generates diagnostic debates about the source genital chlamydia in children.⁶ Health inequalities on this scale affecting First Peoples present all of us health professionals and politicians with a radical challenge, that we must all embrace. We all know the special character of this challenge in our own setting, and have a daily duty to put it at the heart of our service planning and advocacy.

How much does oral sex matter in STI transmission? While it's clearly a minor player in HIV transmission, there is growing interest in the role of oral sex in the transmission of bacterial STI,⁷ and in the significance of oral human papilloma virus (HPV).⁸

Population based studies of adolescents are difficult to do, so we are pleased to publish sister Norwegian studies by Gravningen *et al*^{9, 10} which report self-perceived risk and online partner recruitment in relation to chlamydia infection. Continuing the chlamydial theme, Kampman *et al*¹¹ report on the effectiveness of recalling young people for re-testing following a chlamydia diagnosis.

Other studies explore the value of integrating HIV and sexual health services in an African setting,¹² bacterial vaginosis in women at risk of HIV in Uganda,¹³ mother to child HIV transmission in China,¹⁴ methods of training for STI treatment,¹⁵ the HIV care cascade in Russia¹⁶ and provider offers of HPV vaccine.¹⁷

Finally, we welcome reader letters, and are pleased this month to publish on treatment for rectal chlamydia¹⁸ and Raltegravir for post-exposure HIV prophylaxis.¹⁹

Competing interests None.

Provenance and peer review Commissioned; Not peer reviewed.

REFERENCES

- 1 Wright DJM. Annals of the Royal College of Surgeons of England. *John Hunter and Venereal Disease* 1981;63:198–201.
- 2 Zenilman J. Ethics gone awry: the US Public Health Services studies in Guatemala. *Sexually Transmitted Infections* 2013;89:295–300.
- 3 Fleck L. *Genesis and Development of a Scientific Fact*. 1981. University of Chicago Press, Chicago.
- 4 Chadwick JA, MacNab A, Sarma J, *et al*. Secondary syphilis presenting with aortitis and coronary ostial occlusion. *Sex Transm Infect* 2016;92:108–9.
- 5 Towns JM, Leslie DE, Denham I, *et al*. Painful and multiple anogenital lesions are common in men with

Treponema pallidum PCR-positive primary syphilis without herpes simplex virus coinfection: a cross-sectional clinic-based study. *Sex Transm Infect* 2016;92:110–15.

- 6 Giffard PM, Brenner NC, Tabrizi SN, *et al*. Chlamydia trachomatis genotypes in a cross-sectional study of urogenital samples from remote Northern Central Australia. *BMJ Open* 2016;6:e009624.
- 7 Barbee LA, Khosropour CM, Dombrowski JC, *et al*. An estimate of the proportion of symptomatic gonococcal, chlamydial and non-gonococcal nonchlamydial urethritis attributable to oral sex among men who have sex with men: a case-control study. *Sex Transm Infect* 2016;92:155–60.
- 8 Rosen BJ, Walter L, Gilman RH, *et al*. Prevalence and correlates of oral human papillomavirus infection among healthy males and females in Lima, Peru. *Sex Transm Infect* 2016;92:149–54.
- 9 Gravningen K, Braaten T, Schirmer H. Self-perceived risk and prevalent chlamydia infection among adolescents in Norway: a population-based cross-sectional study. *Sex Transm Infect* 2016;92:91–6.
- 10 Gravningen K, Aicken CRH, Schirmer H, *et al*. Meeting sexual partners online: associated sexual behaviour and prevalent chlamydia infection among adolescents in Norway: a cross-sectional study. *Sex Transm Infect* 2016;92:97–103.
- 11 Kampman CJG, Koedijk FDH, Driessen-Hulshof HCM, *et al*. Retesting young STI clinic visitors with urogenital *Chlamydia trachomatis* infection in the Netherlands; response to a text message reminder and reinfection rates: a prospective study with historical controls. *Sex Transm Infect* 2016;92:124–9.
- 12 Obure CD, Guinness L, Sweeney S, *et al*. Does integration of HIV and SRH services achieve economies of scale and scope in practice? A cost function analysis of the Integra Initiative. *Sex Transm Infect* 2016;92:130–4.
- 13 Francis SC, Looker C, Vandepitte J, *et al*. Bacterial vaginosis among women at high risk for HIV in Uganda: high rate of recurrent diagnosis despite treatment. *Sex Transm Infect* 2016;92:142–8.
- 14 Zeng H, Chow EPF, Zhao Y, *et al*. Prevention of mother-to-child HIV transmission cascade in China: a systematic review and meta-analysis. *Sex Transm Infect* 2016;92:116–23.
- 15 Weaver MR, Pillay E, Jed SL, *et al*. Three methods of delivering clinic-based training on syndromic management of sexually transmitted diseases in South Africa: a pilot study. *Sex Transm Infect* 2016;92:135–41.
- 16 Wirtz AL, Zelaya CE, Latkin C, *et al*. The HIV care continuum among men who have sex with men in Moscow, Russia: a cross-sectional study of infection awareness and engagement in care. *Sex Transm Infect* 2016;92:161–7.
- 17 Gerend MA, Shepherd MA, Lustria MLA, *et al*. Predictors of provider recommendation for HPV vaccine among young adult men and women: findings from a cross-sectional survey. *Sex Transm Infect* 2016;92:104–7.
- 18 Hathorn E, Ward D, Goold P. What is the most appropriate treatment for rectal *Chlamydia trachomatis* infection? *Sex Transm Infect* 2016;92:115.
- 19 Mulka L, Annandale D, Richardson C, *et al*. Raltegravir-based HIV postexposure prophylaxis (PEP) in a real-life clinical setting: fewer drug–drug interactions (DDIs) with improved adherence and tolerability. *Sex Transm Infect* 2016;92:107.