



OPEN ACCESS

ORIGINAL ARTICLE

Contaminated fingers: a potential cause of *Chlamydia trachomatis*-positive urine specimens

Philip M Giffard,^{1,2} Rachael A Lilliebridge,¹ Judith Wilson,¹ Gerald Murray,^{3,4} Samuel Phillips,^{3,4} Sepehr N Tabrizi,^{3,4,5,6} Suzanne M Garland,^{3,4,5,6} Louise Martin,⁷ Gurmeet Singh,^{7,8,9,10} Steven Y C Tong,^{1,11} Deborah C Holt,^{1,2} Patiyann Andersson¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/sextrans-2016-053081>).

For numbered affiliations see end of article.

Correspondence to

Dr Philip M Giffard, Global and Tropical Health, Menzies School of Health Research, Royal Darwin Hospital Campus, PO Box 41096, Casuarina, Darwin, Northern Territory, Australia; phil.giffard@menzies.edu.au

Received 2 January 2017

Revised 21 April 2017

Accepted 29 April 2017

ABSTRACT

Objectives The detection of an STI agent in a urogenital tract (UGT) specimen from a young child is regarded as being indicative of sexual abuse. However, the probabilities of contamination events that could conceivably lead to STI positive specimens in the absence of sexual contact are unclear. The objective was to estimate the potential for fingers that have come in contact with *Chlamydia trachomatis*-positive urine to detectably contaminate *C. trachomatis*-negative urine.

Methods The study design was based on self-experimentation. Dilutions of *C. trachomatis* elementary bodies (EBs) were prepared. A participant contacted an EB dilution then a urine surrogate specimen. The experiment was performed by three participants using three *C. trachomatis* isolates, of genotype E, F and B. Two surrogate urine contact methods were used to mimic contamination of a carer assisting with a child's urine collection. All EB dilutions and urine surrogate specimens were subjected to *C. trachomatis* assay and quantification in a real-time PCR-based diagnostic system.

Results The amplicon crossing point (Cq) for EB dilutions was 10.0±1.6 less than for corresponding finger contacted urine specimens, which corresponds to ~10 µL of EB suspension transferred. This was largely independent of participant identity, *C. trachomatis* strain or EB dilution. Hand decontamination led to large reductions in EBs transferred, but transfer remained consistently detectable. Recent Cq data from *C. trachomatis*-positive clinical urine specimens were collated, and 20% clearly contained sufficient *C. trachomatis* to detectably contaminate another specimen by finger-mediated transfer, as in this experiment.

Conclusions This study directly demonstrated the potential for urine contaminated fingers to convert a *C. trachomatis*-negative urine specimen to *C. trachomatis* positive as a result of contact. Accordingly, procedures for urine specimen collection, particularly from children, need to be designed to prevent contamination.

INTRODUCTION

The exquisite sensitivity of nucleic acid amplification tests for STIs can lead to the perception that positive results can arise from specimen contamination events. This is particularly significant in the Australia's Northern Territory (NT). Approximately 30% of the NT population are Indigenous Australians, with about half living in remote communities.

Key messages

- Human fingers contaminated with *Chlamydia trachomatis* can detectably contaminate urine surrogate specimens by contact.
- The relationship between level of finger contamination and urine surrogate contamination is highly consistent.
- Three contaminated fingers, dried on paper towel, transfer the equivalent of approximately 10 µL *C. trachomatis* suspension to a urine surrogate specimen.
- It is critical that urine specimen collection procedures, in particular from children, should be in accordance with procedures designed to prevent contamination from any source.

The notification rates for STIs are high in the NT, especially in the Indigenous population. For example, the rate of *Chlamydia trachomatis* notifications in Indigenous people in the NT is approximately five times higher than the notification rate in the Australian population as a whole.¹

The 'Little Children Are Sacred' report from 2007² detailed widespread concern of child sexual abuse in NT remote Indigenous communities. This led to large-scale and controversial interventions by the Australian Commonwealth Government, which are ongoing in modified form. Long-standing tensions between the imperatives of child protection and avoidance of disruption and stereotyping of communities and families remain.³ This may manifest in difficulties formulating service provider responses to instances of detection of STI agents in the urine of young children, when there is no other evidence of sexual abuse. While recognising that an STI in a child is a strong indicator of sexual abuse, an absence of other indicators can lead some service providers to raise the possibility of a urine specimen or anatomical site contamination. Studies outside Australia have lent credence to this concept.⁴⁻⁶

We are investigating conceivable mechanisms that diagnostic specimens may become contaminated. We have previously reported the potential of contaminated toilet-bathroom facilities to give rise to contaminated specimens,⁷ and also addressed the question as to whether *C. trachomatis* derived from ocular infections rather than sexual contact



CrossMark

To cite: Giffard PM, Lilliebridge RA, Wilson J, et al. *Sex Transm Infect* Published Online First: [please include Day Month Year]. doi:10.1136/sextrans-2016-053081

can be reliably identified by genotyping.^{8,9} Here, we test another conceivable mechanism; that a carer with *C. trachomatis* contaminated fingers who assists a child to provide a urine specimen may convert a specimen from *C. trachomatis* negative to *C. trachomatis* positive, as a result of contact between their fingers and the specimen.

METHODS

The study design basis was simulation by self-experimentation performed by three members of the investigator team (termed participants). Three *C. trachomatis* isolates were used. They were all derived from the 'Mother-Child' study, a survey of *C. trachomatis* in mothers and their babies, performed in the north of the NT between 1986 and 1992.^{10–12} Isolate E_Aus56 is serovar E, isolate F_Aus51 is serovar F and isolate B_Aus45 is serovar B. All three are urogenital tract isolates and B_Aus45 has been genome sequenced.¹²

Each participant contacted suspensions of live *C. trachomatis* with their bare fingers, and then immediately after contacted a 10 mL aliquot of a urine surrogate solution, made as described by Martino and coworkers.¹³ The relative quantities of *C. trachomatis* DNA in the suspensions and urine surrogate solutions were then determined. The experimentation was performed on three different days, with 2 weeks between each occurrence. On each day of experimentation, a single *C. trachomatis* isolate was tested with all three participants. In advance of experimentation, the isolate to be used was grown to high infection levels in McCoy cells (90%–95% of host cells infected) as previously described.¹² Elementary bodies (EBs) were released from host cells and concentrated, as previously described,¹² and a 100-fold dilution series in urine surrogate prepared yielding dilutions of 10^0 – 10^{-12} for E-Aus56 and F_Aus51 and 10^0 – 10^{-14} for B_Aus45. Between 30 and 60 min in advance of the experiment, the experimenters washed their hands in a general purpose hand and body wash (Avagard 9230-D (3M)), rinsed with tap water, then rinsed their hands with 0.5% chlorhexidine gluconate in 70% ethanol (Avagard 9250 P (3M)) and allowed this to dry.

Two different methods of contact between the experimenter's *C. trachomatis* contaminated fingers and urine surrogate were used: (1) dipping the contaminated fingers into 10 mL urine surrogate in a urine pot for 10 s (method A—dipping method (performed first, using the left hand)) and (2) running 10 mL of urine surrogate over the contaminated fingers into a container (method B—pouring method (performed second, using the right hand)). These methods were first used to generate negative control (N1) specimens, where there was no prior contact between fingers and an EB suspension on that day. Then, commencing with the most dilute EB suspension, the participant contacted each EB dilution with the first, second and third fingers, dried the fingers briefly on a paper towel, then contacted a 10 mL urine surrogate specimen with the same fingers. On completion of both contact methods with all dilutions of EB suspension, the participant decontaminated their hands using the same procedure as the pre-experimentation hand decontamination. After this, on the same day, the participant generated 'postexperiment, postdecontamination' (N2) specimens by contacting fresh 10 mL urine surrogate with the same fingers of each hand as had been used previously. In this instance, the dipping method was used with both hands.

In the interests of practicality and safety, each experiment was performed using a class II biosafety cabinet by participant and two assistants. This ensured the participant touched nothing with their fingers in the course of the experiment apart from the

EB dilutions, the urine surrogate samples and the paper towel used for drying.

The EB dilutions and the urine surrogate specimens were all subjected to analysis for *C. trachomatis* DNA, using the Roche Molecular Diagnostics Cobas 4800 CT/NG commercial diagnostic system (Roche Diagnostics Australia). This provides quantitation information in the form of the amplimer concentration 'crossing point' (Cq). Provided there is doubling of the amplimer concentration with each PCR cycle, the Cq is inversely proportional to the \log_2 (amplimer), with a gradient of -1 . In general, Cq results ≥ 40 were not included in subsequent analyses. We reasoned that results right at the limit of detection of the Roche instrument are potentially due to contamination from, for example, the air in the biohazard hood, and may confound rather than enhance our efforts to reliably determine the relationship between *C. trachomatis* loads in EB suspensions and corresponding finger-contacted urine surrogate specimens. Similarly, Cq values from positive reactions where a less diluted EB suspension yielded a negative reaction were excluded from further analysis. All results, including those from specimens that yielded positive results with Cq values ≥ 40 , are presented in the supplementary data.

The difference between the Cq for an EB suspension and the Cq for a corresponding finger contacted urine surrogate specimen was defined as the ΔCq . Assuming amplimer doubling with every PCR cycle, the volume of EB suspension transferred to the urine surrogate (in μL) = $10\,000/2^{\Delta Cq}$, where 10 000 is the volume of each urine surrogate specimen in μL and $2^{\Delta Cq}$ is the ratio of target concentration between EB suspension and derived urine surrogate specimen. A similar approach was used to calculate the reduction in *C. trachomatis* load on the fingers concomitant with hand decontamination after the completion of the transfer experiment. In this case, the parameter $\Delta Cq(N2)$ is the difference between the Cq value for a urine surrogate specimen arising from transfer from the undiluted EB suspension (ie, the final data point in the transfer experiment) and the Cq for the urine surrogate specimen contacted by the postdecontamination fingers. The numerical reduction in load is thus $2^{\Delta Cq(N2)}$.

The statistical methods used were primarily one-way analysis of variance (ANOVA) and Tukey 'Honest Significant Difference' (HSD) for comparisons of more than two categories and the t-test for comparison of two categories. In general, ΔCq values were used rather than calculated values for volumes of EB dilution transferred. This is because the statistical methods used are reliant on normal distribution of data. It was determined that the distribution of ΔCq values was consistent with normality, according to the Shapiro-Wilks test ($p=0.92$). In contrast, the distribution of deduced transferred volumes was inconsistent with normality, with a long tail towards high values.

RESULTS

Initial inspection of the data revealed a consistent relationship between the Cq values derived from EB suspensions, and Cq values derived from the corresponding finger contacted urine surrogate specimens. Results from a typical experiment are shown in [figure 1](#), and all results are shown in supplementary information.

We tested the assumption that there was amplimer doubling in every PCR cycle. A linear regression of \log EB dilution versus Cq for all Cq values < 40 was performed. This yielded $Cq=3.3 \times \log \text{EB dilution} + 16.0$ ($R^2=0.98$). Because $3.3 = \log_2 10$, it was concluded that the assumption of target doubling in each




Figure 1 Typical result from an EB transfer experiment for one strain, with the solid line representing Cq values from dilutions of an EB suspension and the dotted line representing the mean Cq from the corresponding finger contacted urine surrogate specimen, for all three participants. EB, elementary body.

PCR cycle is justified, so allowing the volumes of EB suspensions transferred to be calculated as described in the Methods.

Collated results from all specimens apart from the N1 pre-experimentation specimens and the N2 posthand decontamination specimens are summarised in [table 1](#). In accordance with the Methods, six Cq values from finger contacted urine surrogate specimens were excluded from analysis as they had Cq values ≥ 40 (three specimens), yielded positive reactions where a less diluted EB specimen yielded a negative reaction (one specimen) or fulfilled both criteria (two specimens) (see online supplementary data). The mean ΔCq value across the entire experiment (not including the N1 and N2 specimens) was 10.0 ± 1.6 , which equates to $\sim 10 \mu\text{L}$ of EB dilution being transferred to the urine surrogate specimens, with an inferred range of volumes transferred being $\sim 1\text{--}100 \mu\text{L}$. The ΔCq values from participant 1 were significantly lower than the ΔCq values from the other two experimenters, indicating that participant 1 transferred significantly more EBs than the other two. Similarly, the ΔCq for the pouring method was slightly but significantly less than for the dipping method. In contrast, there was no significant difference in the ΔCq values between the three different *C. trachomatis* strains. The 10^{-2} dilutions gave rise to significantly

higher ΔCq values than either the undiluted suspension or 10^{-4} diluted suspension.

The *C. trachomatis* diagnostic test yielded a positive reaction for five of the 18 N1 specimens, with Cq values ranging from 39.6 to 40.8 (see online supplementary data for complete results). All the positive reactions were from specimens obtained immediately before the pouring experiments, after the dipping experiment had been completed.

All of the N2 postdecontamination specimens yielded positive PCR reactions, with Cq values ranging from 32.5 to 40.0. After exclusion of the single Cq value of 40.0, the reductions in *C. trachomatis* DNA resulting from hand decontamination were quantified. The mean of the $\Delta Cq(N2)$ values was 10.3 ± 2.1 , with an absolute range of 6.4–14.8. This equates to a mean reduction of 1271-fold and an absolute range of 85.0-fold to 28526-fold.

To assess the clinical implications of this study, we compared our results to 30 successive Cq values from analysis of specimens analysed in the course of clinical service provision. The Cq values were derived from the same instrument as used to analyse the urine surrogate specimens in this study. We conservatively classed any specimen with a $Cq < 40 - 10 = 30$ as clearly having potential to contaminate a finger sufficiently to potentially give rise to a false positive. Six of the 30 (20%) of the Cq values were ≤ 30 , indicating that such potential exists.

DISCUSSION

The principal finding was remarkably consistent ΔCq values of ~ 10 cycles, with respect to the EB suspensions and corresponding finger-contacted urine surrogate specimens. This was largely independent of suspension density, *C. trachomatis* isolate, participant identity or EB transfer procedure. Comparison with Cq values from a sample of *C. trachomatis*-positive urogenital tract (UGT) clinical specimens indicated that 20% of such specimens can contaminate a human finger sufficiently to be able to convert a *C. trachomatis*-negative specimen to positive as a result of contact. While ΔCq values may provide an indication of the likelihood of detectable contamination of clinical specimens, to make the results more generalisable, we inferred the volumes of EB suspensions transferred from human skin to the 10 mL urine surrogate specimens. The

Table 1 Ranges of ΔCq values and inferred volumes of EB dilutions transferred

Category (n)	Mean ΔCq (SD)	ΔCq range	Mean μL transferred (inferred from $\pm 2 \times \Delta Cq$ SD range)(inferred from observed ΔCq range)	
All (49)	10.0 (1.6)	6.7–13.3	9.8 (1.1–89.7)(1.0–96.2)	
Participant 1 (17)	8.8 (0.9)	6.7–9.8	22.4 (6.4–78.1)(11.2–96.2)	$p < 0.0001^*$
Participant 2 (16)	10.5 (1.2)	8.6–13.0	7.1 (1.4–32.6)(1.2–25.8)	Participant 1 vs participant 2: $p < 0.01 \dagger$
Participant 3 (16)	10.9 (1.6)	7.7–13.3	5.2 (0.6–48.1)(1.0–48.1)	Participant 1 vs participant 3: $p < 0.01 \dagger$ Participant 2 vs participant 3: not significant \ddagger
Dipping (23)	10.6 (1.7)	7.4–13.3	6.4 (0.6–68.0)(1.0–59.2)	$p = 0.015 \ddagger$
Pouring (26)	9.5 (1.3)	6.7–12.4	13.8 (2.3–83.7)(1.0–96.2)	
E_Aus56 (17)	9.7 (1.6)	6.7–12.2	12.0 (1.3–110.5)(2.1–96.2)	Not significant; $p = 0.41^*$
F_Aus51 (15)	10.0 (1.6)	7.4–13.3	9.8 (1.1–89.8)(1.0–59.2)	
B_Aus45 (17)	10.4 (1.5)	8.4–13.0	7.4 (0.9–59.2)(1.2–29.6)	
Undiluted (18)	9.4 (1.7)	6.7–12	14.8 (1.4–156.3)(2.4–96.2)	$p = 0.004^*$
10^{-2} diluted (18)	11.0 (1.5)	9.0–13.3	4.8 (0.6–39.1)(1.0–19.5)	10^0 vs 10^{-2} : $p < 0.01 \dagger$
10^{-4} diluted (13)	9.5 (1.0)	7.7–11.1	13.8 (3.5–55.2)(4.6–48.1)	10^0 vs 10^{-4} : not significant \ddagger 10^2 vs 10^{-4} : $p < 0.05 \dagger$

The 49 data points were collated according to different criteria to reveal possible correlates. This encompasses data only from specimens that yielded a positive *C. trachomatis* test with a Cq < 40 .

*One way analysis of variance.

\dagger Tukey HSD.

\ddagger t-test (two tailed).

self-experimentation procedure transferred into each surrogate urine specimen *C. trachomatis* or DNA equivalent corresponding to a mean of 10 μ L and a range of ~1–100 μ L of EB suspension. It was shown that hand decontamination reduced the amount of *C. trachomatis* DNA transferred from contaminated fingers by ~1000-fold, thus reducing potential for specimen contamination. The principal strength of the study is that it was a very direct test of the potential for *C. trachomatis*-contaminated fingers to contaminate diagnostic specimens, with the remarkable consistency of Δ Cq values, indicating integrity in the experimental design.

We considered whether the positive diagnostic tests for N1 specimens (5/18) indicated that the experiment had been confounded, such as by pre-existing finger contamination. All five positive reactions were from specimens taken after the dipping method experiments and prior to pouring method experiments. This, in combination with extremely high Cq values, suggests the likely basis for the positive N1 reactions was aerosolised *C. trachomatis* inside the biosafety hood arising from the dipping method experiments. While we cannot rule out that such contamination has impacted the experimental results, this can only be the case for specimens that gave Cq values close to 40 and will not have had a significant effect on the study outcomes.

The experimental design did not encompass a systematic determination of the persistence of *C. trachomatis* on the skin. This was primarily because it is unreasonable to expect a participant to contaminate their fingers with live *C. trachomatis* and then refrain from hand decontamination for a long period of time. However, the results from the N1 specimens from the experiments carried out on the second and third days of experimentation provide some insight. The three experimentation days were 2 weeks apart. No *C. trachomatis* DNA was detected in the N1 specimens that were obtained at the start of the second and third days of experimentation, indicating that no detectable finger contamination had persisted from the experimentation 2 weeks, or 2 weeks and 4 weeks prior. This is despite the N2 specimens obtained at the end of the experimentation in experimentation days 1 and 2 being universally positive. Thus, between experiments on different days, the amount of *C. trachomatis* DNA on the fingers of the participants dropped below the level in which they could contaminate specimens. Given that hand decontamination reduced the amount of *C. trachomatis*/*C. trachomatis* DNA transferred by an average of ~1000, this is unsurprising.

The assumption implied by the data analysis method was that the quantitative PCR device accurately measured *C. trachomatis* DNA. However, this may not be justified. It was unexpected that the distribution of Δ Cq values conforms to normality more closely than the distribution of inferred volumes transferred to the urine surrogate. The volume transferred is the physical phenomenon expected to be normally distributed, while the Δ Cq is the product of a log transformation. This suggests that variation in the Δ Cq values may encompass instrumental artefacts as well as variation in the volumes of amounts of EB suspensions transferred.

We observed that participant identity and EB transfer method are correlated with EBs transferred. We do not regard speculation regarding mechanisms as useful, as the effects are small and do not change translational implications. Also, we regard the higher Δ Cq observed with the 10^{-2} dilutions of EB as not mechanistically meaningful. Phenomena such as a limited number of high affinity EB binding sites on the skin could reduce EB transfer below what would be expected when the EB density is low. However, there was no significant difference between the Δ Cq values arising from undiluted suspension and most dilute (10^{-4} diluted) suspensions, so no evidence for loss of EBs at low EB densities. The similar

results from three *C. trachomatis* isolates suggest that our results are generalisable to the species as a whole.

This study has directly shown that contact between contaminated fingers and urine specimens has potential to contaminate the specimen and underpin a positive *C. trachomatis* test. This provides justification for standardised procedures for collection of urine specimens from children, encompassing controlled environments, with clear guidance, and if possible, close supervision by clinical staff. In the absence of other measures such as wearing gloves, hand decontamination has been shown to greatly reduce the potential for specimen contamination. There must be clear instruction on hand washing and avoidance of contact with the urine specimen and interior of the container to all involved in collecting a child's urine sample for STI testing. Regardless, a positive STI test in a child remains a strong indicator of child sexual assault necessitating further investigation by police, statutory child protection agencies and relevant forensic medical teams.

More broadly, this study adds to accumulating knowledge regarding mechanisms that could contaminate specimens, and so give rise to spurious STI positive results. Previous studies have focused on the potential for STI nucleic acid on clinic or laboratory surfaces to find its way into specimens.^{4–7} The emerging picture is that such surface contamination is commonly found but rarely leads to STI positive specimens. The difficulty of proving a negative, plus the demonstration by Andersson and coworkers⁷ of a non-zero frequency of specimen contamination, justifies measures to control the environment and procedure for STI specimen collection and emphasises the benefits of specimen collection under controlled conditions, particularly from young children. This current study adds to the imperative for this, by showing that potential for contact between specimen and uncovered skin should be avoided. In general, guidelines for urine specimens for STI testing simply specify 'first catch urine' and do not specify specific precautions for avoidance of urine contamination. In the NT, poster material available to medical practitioners specifies that patients 'wash your hands with soap'.¹⁴ This is not children specific, and the extent that this procedure is followed is unknown. The absence of such precautions against contamination of specimens from young children could potentially be significant in the judicial system. Interestingly, guidelines for clean catch urine sampling, which is not used in STI testing, do specify avoidance of contact between specimens and fingers for example.¹⁵ Determining the most effective and culturally appropriate means of obtaining UGT specimens from young children while avoiding contamination risks could be a useful area for future research.

Author affiliations

¹Global and Tropical Health Division, Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory, Australia

²School of Psychological and Clinical Sciences, Charles Darwin University, Darwin, Northern Territory, Australia

³Murdoch Childrens Research Institute, Melbourne, Victoria, Australia

⁴Department of Microbiology and Infectious Diseases, The Royal Women's Hospital, Melbourne, Victoria, Australia

⁵Department of Obstetrics and Gynaecology, University of Melbourne, Melbourne, Victoria, Australia

⁶Department of Microbiology, Royal Children's Hospital, Melbourne, Victoria, Australia

⁷Royal Darwin Hospital, Top End Health Service, Northern Territory Government, Darwin, Northern Territory, Australia

⁸Sexual Assault Referral Centre, Northern Territory Government, Northern Territory, Australia

⁹Northern Territory Medical Program, Flinders University, Darwin, Northern Territory, Australia

¹⁰Child Health Division, Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory, Australia

¹Victorian Infectious Disease Service, The Royal Melbourne Hospital, and The University of Melbourne, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia

Contributors PMG: Contributed to study conception, contributed to study design and formulation of detailed experimental procedure, carried out practical work, analysed data, wrote manuscript. RAL: Contributed to study design, contributed to study design and formulation of detailed experimental procedure, carried out practical work, analysed data. JW: Contributed to formulating detailed experimental procedure, carried out practical work, analysed data, commented on manuscript. GM: Carried out practical work, analysed data, commented on manuscript. SP: Carried out practical work, analysed data, commented on manuscript. SNT: Contributed to study conception, contributed to study design, provided clinical data. SMG: Contributed to study conception, contributed to study design, commented on manuscript. LM: Identified problem that this study addressed, contributed to study design, commented on manuscript. GS: Contributed to study design, commented on manuscript. SYCT: Contributed to study conception, contributed to study design, commented on manuscript. DCH: Contributed to study design and formulation of detailed experimental procedure, carried out practical work, analysed data, commented on manuscript. PA: Contributed to study conception, contributed to study design and formulation of detailed experimental procedure, carried out practical work, analysed data, commented on manuscript.

Funding This study was funded by Australian National Health and Medical Research Council project grant 1060768. SYCT was supported by an Australian National Health and Medical Research Council Career Development Fellowship 1065736.

Competing interests None declared.

Patient consent Obtained.

Ethics approval This study was approved by the Human Research Ethics Committee of the Northern Territory Department of Health and the Menzies School of Health Research, under ethical clearance number 2011-1673.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data are included in the publication itself.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 The Kirby Institute. *HIV, viral hepatitis and sexually transmissible infections in Australia. Annual Surveillance Report 2015*. Australia, Sydney NSW: The Kirby Institute UNSW, 2015.
- 2 Wild R, Anderson P, Ampe Akelyernemane Meke Mekarle, 'Little Children Are Sacred' Report of the board of enquiry into the protection of Aboriginal children from sexual abuse. 2007 http://www.inquirysaac.nt.gov.au/pdf/bipacsa_final_report.pdf
- 3 Brown A, Brown NJ. The Northern Territory intervention: voices from the centre of the fringe. *Med J Aust* 2007;187:621–3.
- 4 Chan SY, Jose S, King R, et al. How likely is environmental or patient cross-contamination of *Chlamydia trachomatis* DNA to lead to false positive results in patients attending our clinic? *Sex Transm Infect* 2013;89:105–7.
- 5 Lewis N, Dube G, Carter C, et al. Chlamydia and gonorrhoea contamination of clinic surfaces. *Sex Transm Infect* 2012;88:418–21.
- 6 Meader E, Waters J, Sillis M. Chlamydia trachomatis RNA in the environment: is there potential for false-positive nucleic acid amplification test results? *Sex Transm Infect* 2008;84:107–10.
- 7 Andersson P, Tong SY, Lilliebridge RA, et al. Multisite direct determination of the potential for environmental contamination of urine samples used for diagnosis of sexually transmitted infections. *J Pediatric Infect Dis Soc* 2014;3:189–96.
- 8 Giffard PM, Brenner NC, Tabrizi SN, et al. Chlamydia trachomatis genotypes in a cross-sectional study of urogenital samples from remote Northern and Central Australia. *BMJ Open* 2016;6:e009624.
- 9 Giffard PM, Singh G, Garland SM. What does Chlamydia trachomatis detection in a urogenital specimen from a young child mean? *Sex Transm Infect* 2016;sextrans-2015-052473.
- 10 Bowie WR, Caldwell HD, Jones RP, et al. eds. Proceedings of the Seventh International Symposium on Human Chlamydial Infections. International Symposium on Human Chlamydial Infections; 1990; Harrison Hot Springs. Press Syndicate of the University of Cambridge.
- 11 Oriel D, Ridgway GL, Schachter J, et al. eds. Proceedings of the Sixth International Symposium on Human Chlamydial Infections. International Symposium on Human Chlamydial Infections; 1986; Sanderstead, Surrey. Press Syndicate of the University of Cambridge.
- 12 Andersson P, Harris SR, Seth Smith HM, et al. Chlamydia trachomatis from Australian aboriginal people with Trachoma are polyphyletic composed of multiple distinctive lineages. *Nat Commun* 2016;7:10688.
- 13 Martino PD, Fursy R, Bret L, et al. Indole can act as an extracellular signal to regulate biofilm formation of *Escherichia coli* and other indole-producing bacteria. *Can J Microbiol* 2003;49:443–9.
- 14 Anonymous. Specimen collection posters. 2011 <http://digitalibrary.health.nt.gov.au/prodjspsui/handle/10137/531>
- 15 Kirkwood J. Clean catch urine sample and culture. 2016 <http://www.healthline.com/health/urine-culture-clean-catch#overview1>

STI

Contaminated fingers: a potential cause of *Chlamydia trachomatis*-positive urine specimens

Philip M Giffard, Rachael A Lilliebridge, Judith Wilson, Gerald Murray, Samuel Phillips, Sepehr N Tabrizi, Suzanne M Garland, Louise Martin, Gurmeet Singh, Steven Y C Tong, Deborah C Holt and Patiyan Andersson

Sex Transm Infect published online June 9, 2017

Updated information and services can be found at:

<http://sti.bmj.com/content/early/2017/06/08/sextrans-2016-053081>

These include:

References

This article cites 8 articles, 4 of which you can access for free at:

<http://sti.bmj.com/content/early/2017/06/08/sextrans-2016-053081#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (249)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>