

Supplemental Table 1. Cytology and histology results among men who have sex with men living with HIV.

Cytology	HRA					
	No biopsy (n=95) n (%)	Negative (n=45) n (%)	LSIL-AIN1 (n=14) n (%)	HSIL-AIN2 (n=14) n (%)	HSIL-AIN3 (n=6) n (%)	Unsatisfactory (n=2) n (%)
NILM	57 (60)	21 (46.7)	6 (42.9)	1 (7.1)	0 (0.0)	2 (100)
ASC-US	13 (13.7)	10 (22.2)	2 (14.3)	2 (14.3)	0 (0.0)	0 (0.0)
LSIL	18 (18.9)	8 (17.8)	6 (42.9)	5 (35.7)	1 (16.7)	0 (0.0)
ASC-H	2 (2.1)	0 (0.0)	0 (0.0)	1 (7.1)	1 (16.7)	0 (0.0)
HSIL	1 (1.1)	5 (11.1)	0 (0.0)	5 (35.7)	4 (66.7)	0 (0.0)
Unsatisfactory	4 (4.2)	1 (2.2)	0 (0.0)	0 (0.0)	0 (0)	0 (0.0)

NILM: negative for intraepithelial lesion and malignancy; ASC-US: atypical squamous cells of undetermined significance, ASC-H: atypical squamous cells cannot exclude HSIL; LSIL: low grade squamous intraepithelial lesion; HSIL: high grade squamous intraepithelial lesion. HRA: High resolution anoscopy. AIN: Anal intraepithelial neoplasia.

Table created by the authors