caesarian section must have been contaminated by an ascending infection either before or after the rupture of the placental membranes, or alternatively, by contamination after birth. In a separate study (Goeman J, *et al*, unpublished observation) of 205 infants with ophthalmia neonatorum, only one out of 38 with chlamydial ophthalmia neonatorum was delivered by caesarian section whereas four of 19 *N gonorrhoeae* positive babies with ophthalmia neonatorum and 22 of 148 with non-chlamydial, non-gonococcal ophthalmia neonatorum were delivered by caesarian section. Thus chlamydial ophthalmia neonatorum was observed in 4% of babies with ophthalmia neonatorum delivered by caesarian section and 26% of those delivered vaginally.

Although our study size was too small to rule out the possibility of prepartal infection of the placenta by chlamydiae, it must be less common than gonococcal infection of the placenta.

Yours faithfully,
E Frost
A Leclerc
G Gioanni
J Goeman
M Peeters
M Collet

Centre International de Recherches
Médicales de Franceville,
BP 769,
Franceville, Gabon

References

- Yvert F, Frost E, Walter P, Gass R, Ivanoff B. Prepartal infection of the placenta with *Neisseria gonorrhoeae*. *Genitourin Med* 1985; **61**:103-5.
- Lacey CJN, Milne JD. Preterm labour in association with *Neisseria gonorrhoeae*. *British Journal of Venereal Diseases* 1984; **60**:123-4.
- Givner LB, Rennels MB, Woodward CL, Huang SW. *Chlamydia trachomatis* infection in infant delivered by caesarian section. *Pediatrics* 1981; **68**:420-1.
- Hammerschlag MR, Anderka M, Semine DZ, McComb D, McCormack WM. Prospective study of maternal and infantile infection with *Chlamydia trachomatis*. *Pediatrics* 1979; **64**:142-8.
- Rees E, Tait AI, Hobson D, Johnston FWA. Perinatal chlamydial infection. In: Hobson D, Holmes KK, eds. *Nongonococcal urethritis and related infections*. Washington DC: American Society for Microbiology, 1977: 140-7.
- Hansman D. Inclusion conjunctivitis. *Med J Aust* 1969; **1**:151-3.
- Barry WC, Teare EL, Uttley AHC, *et al*. *Chlamydia trachomatis* as a cause of neonatal conjunctivitis. *Arch Dis Child* 1986; **61**:797-9.
- Wilt JC, Wilt PC, Kordova N, Martin C. The human placenta as a possible reservoir of

chlamydial infection in northern Canada. *Can J Public Health* 1976; **67**:114-6.

- Pankuch GA, Appelbaum PC, Lorenz RP, Botti JJ, Schachter J, Naeve RL. Placental microbiology and histology and the pathogenesis of chorioamnionitis. *Obstet Gynecol* 1984; **64**:802-6.
- Mårdh P-A, Johansson PJ, Svenningsen N. Intrauterine lung infection with *Chlamydia trachomatis* in a premature infant. *Acta Paediatr Scand* 1984; **73**:569-72.
- Schachter J. Chlamydial infections. *N Engl J Med* 1978; **298**:540-9.
- Armstrong JH, Zacarias F, Rein MF. Ophthalmia neonatorum: a chart review. *Pediatrics* 1976; **57**:884-92.
- Frost E, Yvert F, Zué Ndong J, Ivanoff B. Ophthalmia neonatorum in a semi-rural African community. *Trans R Soc Trop Med Hyg* 1987; **81**:378-80.
- Leclerc A, Frost E, Collet M, Goeman J, Bedjabaga L. Urogenital *Chlamydia trachomatis* in Gabon: an unrecognised epidemic. *Genitourin Med* 1988; **64**:308-11.

TO THE EDITOR, *Genitourinary Medicine*

Genital herpes

Sir,
The number of cases of genital herpes nationally has increased. Returns to the DHSS showed 7327 cases in 1977¹ and 11 147 in 1981,² and other sexually transmitted viral diseases have also increased during the past decade.

Herpes genitalis is a disease characterised by recurrences, so to assess the relative contribution made to this increasing total by first attacks of herpes and by recurrences, we reviewed the notes of patients attending the department of genitourinary medicine in Leicester with herpes during the same six month periods in 1982 and 1987 (table). As expected, the total cases of herpes increased during those years (90 cases in 1982, 134 cases in 1987). Separating the episodes into first attacks and recurrences shows that there was a modest increase in first attacks (63 increasing to 72), whereas recurrences doubled; the largest increase being in men (12 increasing to 41).

These figures show the expected increase in recurrences as the number of infected people increases with time. With the increasing pool of infection, the number of uninfected people at risk will also increase (as long as the size of the pool is small compared with the population at risk, which it is). Assuming continued sexual activity, therefore, the number of first attacks will increase.

It is interesting that the number of first attacks in men actually decreased, but recurrences showed a threefold increase,

Table Incidence of genital herpes in Leicester in two six month periods

	1982	1987
Men:		
First attack	34	29
Recurrences	12	41
Total	46	70
Women:		
First attack	29	43
Recurrences	15	21
Total	44	64
Both:		
First attack	63	72
Recurrences	27	62
Total	90	134

whereas recurrences in women showed little increase during the study period.

Yours faithfully,
P G Fisk
D T P Evans

Department of Genitourinary Medicine,
Royal Infirmary, Leicester LE1 5WW

References

- Anonymous. Sexually transmitted diseases. Extract from the annual report of the CMO of the DHSS for 1977. *British Journal of Venereal Diseases* 1979; **55**:225-9.
- Anonymous. Sexually transmitted diseases. Extract from the annual report of the CMO of the DHSS for 1981. *British Journal of Venereal Diseases* 1984; **60**:199-203.