- 6 Morris DJ. Is human immunodeficiency virus (HIV) rather than cytomegalovirus the cause of retinitis and colitis in HIV-infected patients? *Rev Infect Dis* 1990;12:557-9.
- 7 Morris DJ. Antiviral chemotherapy for cytomegalovirus disease. *J Antimicrob Chemother* 1988;21:519-22.

Accepted for publication 13 September 1990.

Gastrointestinal obstruction associated with *Chlamydia trachomatis*

I read with great interest the recent report of Pegg and Owen regarding small bowel obstruction associated with *Chlamydia trachomatis*.¹ I would like to point out, however, that this phenomenon was first suggested in a similar case report in 1987,² and recently included in a review of abdominal pain syndromes caused by chlamydial infections.³

While both cases^{1,2} presented clinical, historical and serologic evidence of a chlamydial aetiology for the small bowel obstruction, the report of Pegg and Owen additionally demonstrated *C trachomatis* in the genital tract by ELISA testing.

The women in both reports had no genital tract complaints at the time of presentation, and their fallopian tubes appeared normal. Both of these cases strongly suggest that infection with *C trachomatis* may result in small bowel obstruction, and that pelvic symptoms may not be temporally associated with the abdominal disease.

DENNIS J BAUMGARDNER
University of Illinois
College of Medicine-Urbana/Champaign,
The Carle Foundation,
611 West Park St,
Urbana, IL 61801, USA

- 1 Pegg DJ, Owen AWM. Gastrointestinal obstruction associated with *Chlamydia trachomatis*. *Gentourin Med* 1990;66:26-7.
- 2 Baumgardner DJ, McCann DE. Peritonitis and small bowel obstruction from occult pelvic inflammatory disease. *Illinois Medical Journal* 1987;171:75-8.
- 3 Baumgardner DJ. Abdominal pain: Chlamydia as culprit. *Postgrad Med* 1989;85:281-8.

Value of performing biopsies in genitourinary clinics

We read with interest the recent letter in your journal concerning the value of

performing biopsies in genitourinary clinics.¹

Men with abnormalities of the skin of the penis frequently present to genitourinary physicians, who must therefore also be skilled in dermatology. This is particularly so in the light of the probable re-definition of our specialty as "dermato-venereology" in the unified post-1992 European Community.

In order to assess the specific dermatological problems encountered by a busy genitourinary medicine clinic, we set up an internal clinic for penile dermatoses. The criteria for referral were a penile dermatosis of uncertain diagnosis for which the clinician thought that a biopsy might prove helpful. The technique used was that of local anaesthesia followed by skin snip biopsy. All patients were thoroughly counselled before the procedure, which was generally regarded by the patient as not being as traumatic as initially envisaged.

Over a period of two months a total of 18 biopsies were performed with the following histopathological results: 5 nonspecific dermatitis, 3 viral warts, 3 lichen sclerosis et atrophicus, 2 lichen planus, 2 symptomatic subclinical papillomavirus infection, 1 granulomatous disease (currently undergoing investigation) and 1 trauma (probable dermatitis artefacta). Although only a small sample was taken, typically 0.5 mm in diameter, in only one case did the biopsy prove non-diagnostic.

It can be seen that in 11 out of 18 of cases, a specific diagnosis was able to be made for which a management plan could be devised. We would draw particular attention to the diagnoses of lichen sclerosis et atrophicus (LSA). Not only are there multiple names for the same histopathological condition (LSA, balanitis xerotica obliterans and kraurosis vulvae), but the malignant potential of this common² condition remains undefined and a standard text³ suggests six to twelve monthly follow-up for life.

Despite the longstanding combination of the specialties of dermatology and venereology on the continent, the literature on genital dermatology is scant. The three specialties of genitourinary medicine, dermatology and urology have overlapping interests in penile cutaneous disorders, but rarely have in-depth knowledge. Unlike the vulval cutaneous disorders,⁴ for exam-

ple, there is no standard textbook in the English language on penile dermatoses. Furthermore, confusions still exist over relatively common disorders, as described above. It is thus important to develop and improve lines of communication between genitourinary physicians, dermatologists and histopathologists at regular audit meetings.

We would therefore wholeheartedly agree with the conclusion of Drs Arumainayagam and Sumathipala that penile biopsy is a very useful diagnostic procedure in the setting of a genitourinary clinic. The more widespread use of this simple and minimally invasive procedure would allow us to gain greater insight into the ill-understood incidence and nature of genital dermatoses.

R J HILLMAN
J R W HARRIS
Department of Genitourinary Medicine
M M WALKER
Department of Histopathology,
St Mary's Hospital, Praed St,
London W2 1NY, UK

- 1 Arumainayagam JT, Sumathipala AHT. Value of performing biopsies in genitourinary clinics. *Gentourin Med* 1990;66:407.
- 2 Chalmers RJG, Burton PA, Bennett R, Goring CC. Lichen sclerosis et atrophicus—a distinctive and common cause of phimosis in boys. *Br J Dermatol* 1982;107 (suppl):29-30.
- 3 Rowell NR. Lichen sclerosis. In: Rook A, Wilkinson DS, Ebling FJG, Champion RH, Burton JL eds. *Textbook of Dermatology*. Oxford: Blackwell Scientific Publication, 1986: 1368-74.
- 4 Wilkinson EJ ed. *Pathology of Vulva and Vagina*. New York: Churchill Livingstone, 1987.

Sexual assault of men: a series

The sexual assault of men has attracted little attention. The report of Hillman and colleagues describes five cases of male sexual assault from two large genitourinary medicine (GUM) departments during an unspecified time period.¹ We suspect their report is not representative of men attending GUM departments after sexual assault and write to report our experience.

During 1989, 10 male patients attended this department reporting penetrative sexual assault by men. They presented four days to one year after the assault and patient details are