ORIGINAL ARTICLE

An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester

M Ashton, W Sopwith, P Clark, D McKelvey, L Lighton, D Mandal

Sex Transm Infect 2003;**79**:291–293

See end of article for
authors' affiliations

Correspondence to:
Matthew Ashton,
Communicable Disease
Surveillance Centre (North
West), 57a Upper
Northgate Street, Chester
CH1 4EF, UK;
matthew.ashton@hpa.org.uk

Accepted for publication
14 April 2002

Background: In the past few years, increased diagnoses of syphilis have been reported in cities around Britain and Europe. Enhanced surveillance of cases began in 1999 to identify the epidemiology of this increase in Greater Manchester.

Methods: Information was collected on all cases of syphilis newly diagnosed in genitourinary medicine (GUM) clinics in Greater Manchester between January 1999 and November 2002. The data collected included demographic information and information about other sexually transmitted infections, sexual behaviour, perception of risk of infection, and awareness of syphilis transmission.

Results: The majority of cases identified were white homosexual men resident in Greater Manchester. Of the 414 cases diagnosed, 74% had either a primary or secondary stage of syphilis infection and 37% of cases were HIV positive. High numbers of individuals practised unprotected oral sex despite good awareness of the risk of infection with syphilis. There is evidence that the way people are meeting sexual contacts is changing, with increasing numbers meeting most of their partners through the internet.

Conclusions: These findings have implications for targeting interventions. The provision of rapid diagnostic and treatment services is likely to be key for the control of syphilis and potentially of subsequent increases in HIV in the region.

The incidence of infectious syphilis diagnosed from genitourinary medicine (GUM) clinics in the United Kingdom increased almost sixfold between 1996 and 2001.¹ Through this period, the greatest proportional increase in cases has been among men who have sex with men (MSM).² Several local outbreaks of syphilis have contributed to this increase, including one identified in Greater Manchester in 1999.³ Initial investigations in Manchester revealed a high proportion of cases among MSM⁴ and identified oral sex as a key transmission route for syphilis in Manchester.^{4,5}

We initiated a system of "enhanced surveillance" in February 2000, collecting epidemiological information to establish the extent of infection in Greater Manchester and illuminate the patterns of transmission.

METHODS

We devised a form for the collection of epidemiological data, including demographic data, diagnosis, and risk factors and distributed it to each of the 11 GUM clinics in Greater Manchester. Health workers completed one form for each case of syphilis by means of an interview or from case notes where this was not possible. The case definition used was all cases of laboratory confirmed infectious syphilis diagnosed in Greater Manchester from 1999 onwards. For cases diagnosed in 1999 and early 2000, as much data as possible were collected retrospectively from case notes. Completed forms were returned to the North West Office of the Communicable Disease Surveillance Centre (CDSC NW) for collation, data entry, and data analysis.

RESULTS

Between January 1999 and November 2002, there were 414 cases of syphilis reported to the enhanced surveillance database by GUM clinics in Greater Manchester. Of these cases, 93% (377/405) were male and 81% (330/405) were homosexual. Of those for whom data were available, 93% of cases (354/382) were born in the United Kingdom, and 82%

(310/380) were residents of Greater Manchester. There was no particular residential clustering of cases. The majority of cases (90%; 342/381) were white. The next most significant defined ethnic groups of cases were "Black Caribbean" and "Indian" (each 2.1%; 8/381).

The epidemic curve demonstrates some seasonal variation in reporting of infection, with the New Year and summer periods appearing to be key times for identifying new cases (fig 1). Of those for whom a diagnosis was available, 79% of cases (289/364) were either at primary or secondary stages of infection and the ratio of primary to secondary to early latent cases did not vary significantly during the period reported. Since 1999, two thirds of cases were under 35 years old and the peak incidence was among 31–35 year olds.

Most cases (71%; 226/317) were self referrals to GUM clinics and only 10% of cases were referred by general practitioners. Cases often reported other infections in addition to syphilis; 30% (85/288) of the two thirds for whom data were available reported a concurrent infection and 35% of cases were HIV positive (107/308). The HIV status of 106 cases (26% of the total) was not known. In addition to concurrent infections, 52% of cases (201/386) reported having had a previous STI.

Oral sex is a key transmission route for syphilis in Greater Manchester. Very few cases (7%, 12/169) reported using a condom at any time for oral sex (that is, using one sometimes, mostly, or always) and this was in stark contrast to practices for anal sex, for which 95% (148/156) of cases reported using a condom at some time (table 1). Cross tabulation of condom use by perceived risk of catching syphilis from an infected person through unprotected oral sex shows that 61% of homosexual men (87/143) believed the chance of catching syphilis was high, but reported never using a condom. Of these, 70% (61/87) were also aware of the outbreak of syphilis in Greater Manchester before attending the clinic.

There is evidence that the means by which cases meet sexual contacts are changing. Between 2001 and 2002, the proportion of cases locating contacts through the internet

Figure 1 Reported cases of syphilis in Greater Manchester between January 1999 and November 2002, by month and showing 3 month rolling average.

Table 1 Frequency of condom use reported by cases of syphilis for oral and anal sex

Condom use	Oral sex		Anal sex	
	Cases	%	Cases	%
Never	157	93	8	5
Sometimes	10	6	33	21
Mostly	1	1	51	33
Always	1	1	64	41
Total	169	100	156	100
(Unknown)	(245)		(258)	

markedly increased, compared with the more traditional venues of pubs and clubs. In 2002, almost 30% of “high risk” cases (those reporting 10 or more partners in the past year) used the internet to locate most of their contacts.

DISCUSSION

The sustained increase in the number of cases of syphilis in Greater Manchester demanded ongoing surveillance following the initial outbreak investigation. This reveals a high proportion of those practising oral sex reporting never using a condom. Almost two thirds of these cases believed that the chance of catching syphilis from oral sex was high. Of those who believed oral sex to be risky, 70% were also aware of the syphilis outbreak before attending the clinic. These findings suggest not only an increase in high risk sexual behaviour but also an increase despite widespread awareness of the risks.

There is evidence that safer sex messages targeted at the homosexual community in Manchester since 2000 have had some impact. In August 2000, only 37% of the 27 cases interviewed considered unprotected oral sex to be a high risk activity for catching syphilis,⁴ compared with 65% shown here. Despite the increased awareness, transmission of syphilis has continued in Manchester and remains focused within the gay community.

Long waiting lists for GUM services are identified as possible factors contributing to missed cases of syphilis.⁴ Prolonged waiting times for appointments at GUM services are reported nationwide.⁶ The majority of syphilis cases reported here were identified through self referral at GUM clinics and other health services have not generally been considered by patients in the past as reliable sources of advice or care.⁴ Improvements in GUM services are urgently required, therefore, to enable effective control of syphilis in Manchester, and strengthening

the role of other health services such as general practitioners should also be considered.⁷

There is evidence that the routes by which people locate sexual contacts in Manchester are changing. Between 2001 and 2002, the greatest percentage increase was the use of the internet to locate contacts. This change in behaviour has implications for the targeting of intervention strategies.

As with other studies, a significant proportion of cases of syphilis were coinfecting with HIV. Recent studies in Amsterdam and Switzerland report increasing levels of high risk sexual behaviour among individuals with HIV^{8,9} and there is some indication that the introduction of antiretroviral therapies has contributed to this.⁸ A twofold to fivefold increased risk of HIV infection has been reported for patients with existing sexually transmitted infections, including syphilis.¹⁰ It is not clear in the Manchester cohort whether the levels of concurrent infection are linked with unsafe sexual behaviour or a biological effect of concurrent infection. There is a very real risk of a significant increase in HIV infection among those infected with syphilis and this potential association should be made clear in intervention strategies. The rapid detection and treatment of syphilis may be a key factor in the successful attainment of the government's target to reduce new cases of HIV by 25% by 2007.¹¹

ACKNOWLEDGEMENTS

We thank all the members of the Outbreak Investigation Team, including the original leads Dr Ann Hoskins and Professor Qutub Syed, for the hard work done since the outbreak was identified. We are also indebted to the GUM services in Greater Manchester for their support of enhanced surveillance in the region.

CONTRIBUTORS

MA and WS analysis and writing; PC, LL, DMcK, and DM, members of Outbreak Investigation Team.

Authors' affiliations

M Ashton, W Sopwith, Communicable Disease Surveillance Centre (North West), 57a Upper Northgate Street, Chester CH1 4EF, UK

P Clark, Public Health Sector, Liverpool John Moores University, 70 Great Crosshall Street, Liverpool L3 2AB, UK

D McKelvey, Trafford South PCT, Oaklands House, Washway Road, Sale, Manchester M33 6FS, UK

L Lighton, Greater Manchester Health Protection Unit, Floor 7b, Peel House, Albert Street, Eccles, Manchester M30 0NJ, UK

D Mandal, Manchester Royal Infirmary, GUM Department, Oxford Road, Manchester M13 9WL, UK

REFERENCES

- 1 **Communicable Disease Surveillance Centre.** Syphilis: epidemiological data (web page). 2002; Available at www.phls.org.uk/topics_az/hiv_and_sti/sti-syphilis/epidemiology/epidemiology.htm. (Accessed November 2002).
- 2 **Nicoll A, Hamers FF.** Are trends in HIV, gonorrhoea, and syphilis worsening in western Europe? *BMJ* 2002;**324**:1324–7.
- 3 **Communicable Disease Surveillance Centre.** Increased transmission of syphilis in Manchester. *Commun Dis Rep CDR Weekly [Serial Online]* 2000;**10**:89.
- 4 **Clark P, Cook PA, Syed Q, et al.** Re-emerging syphilis in the North West. Liverpool: Public Health Sector, Liverpool John Moores University, 2001.
- 5 **Cook PA, Clark P, Bellis MA, et al.** Re-emerging syphilis in the UK: a behavioural analysis of infected individuals. *Commun Dis Public Health* 2001;**4**:253–8.
- 6 **Djuretic T, Catchpole M, Bingham JS, et al.** Genitourinary medicine services in the United Kingdom are failing to meet current demand. *Int J STD AIDS* 2001;**12**:571–2.
- 7 **Spence D.** Trends in HIV, gonorrhoea, and syphilis. Sexual health services in general practice can be improved. *BMJ* 2002;**325**:494.
- 8 **Stolte IG, Dukers NHTM, de Wit JBF, et al.** A summary report from Amsterdam: increase in sexually transmitted diseases and risky sexual behaviour among homosexual men in relation to the introduction of new anti-HIV drugs. *Euro Surveillance: Bulletin Europeen Sur Les Maladies Transmissibles European Communicable Disease Bulletin* 2002;**7**:19–22.
- 9 **Dubois-Arber F, Moreau-Gruet F, Jeannin A.** Men having sex with men and HIV/AIDS prevention in Switzerland: 1987–2000. *Euro Surveillance: Bulletin Europeen Sur Les Maladies Transmissibles European Communicable Disease Bulletin* 2002;**7**:16–8.
- 10 **Centers for Disease Control and Prevention.** HIV prevention through early detection and treatment of other sexually transmitted diseases—United States. Recommendations of the Advisory Committee for HIV and STD prevention. *MMWR Recommendations & Reports* 1998;**47**:1–24.
- 11 **UK Department of Health.** The national strategy for sexual health and HIV: implementation action plan. London: Department of Health Publications, 2002.

Clinical Evidence—Call for contributors

Clinical Evidence is a regularly updated evidence based journal available worldwide both as a paper version and on the internet. *Clinical Evidence* needs to recruit a number of new contributors. Contributors are health care professionals or epidemiologists with experience in evidence based medicine and the ability to write in a concise and structured way.

Currently, we are interested in finding contributors with an interest in the following clinical areas:

Altitude sickness; Autism; Basal cell carcinoma; Breast feeding; Carbon monoxide poisoning; Cervical cancer; Cystic fibrosis; Ectopic pregnancy; Grief/bereavement; Halitosis; Hodgkins disease; Infectious mononucleosis (glandular fever); Kidney stones; Malignant melanoma (metastatic); Mesothelioma; Myeloma; Ovarian cyst; Pancreatitis (acute); Pancreatitis (chronic); Polymyalgia rheumatica; Post-partum haemorrhage; Pulmonary embolism; Recurrent miscarriage; Repetitive strain injury; Scoliosis; Seasonal affective disorder; Squint; Systemic lupus erythematosus; Testicular cancer; Varicocele; Viral meningitis; Vitiligo

However, we are always looking for others, so do not let this list discourage you.

Being a contributor involves:

- Appraising the results of literature searches (performed by our Information Specialists) to identify high quality evidence for inclusion in the journal.
- Writing to a highly structured template (about 2000–3000 words), using evidence from selected studies, within 6–8 weeks of receiving the literature search results.
- Working with *Clinical Evidence* Editors to ensure that the text meets rigorous epidemiological and style standards.
- Updating the text every eight months to incorporate new evidence.
- Expanding the topic to include new questions once every 12–18 months.

If you would like to become a contributor for *Clinical Evidence* or require more information about what this involves please send your contact details and a copy of your CV, clearly stating the clinical area you are interested in, to Claire Folkes (cfolkes@bmjgroup.com).

Call for peer reviewers

Clinical Evidence also needs to recruit a number of new peer reviewers specifically with an interest in the clinical areas stated above, and also others related to general practice. Peer reviewers are health care professionals or epidemiologists with experience in evidence based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity, and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and health care professionals, possibly with limited statistical knowledge). Topics are usually 2000–3000 words in length and we would ask you to review between 2–5 topics per year. The peer review process takes place throughout the year, and our turnaround time for each review is ideally 10–14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com or contact Claire Folkes (cfolkes@bmjgroup.com).