

# Multiple risk factors for persistent HBV viraemia in an adult receiving nucleos/tide analogue therapy

Sheila Lumley <sup>1</sup>, Maeve Barlow,<sup>2</sup> Khadija Said <sup>3,4</sup>, Emily Martyn,<sup>3,4</sup> Elizabeth Waddilove,<sup>3</sup> Marion Delphin,<sup>3</sup> Daisy Jennings,<sup>1</sup> Haiting Chai,<sup>1</sup> Agnes Kemper,<sup>2</sup> Joy Ko,<sup>2</sup> Azim Ansari,<sup>1</sup> Douglas Macdonald,<sup>5</sup> Indrajit Ghosh,<sup>2</sup> Samreen Ijaz,<sup>6</sup> Stuart Flanagan <sup>2</sup>, Philippa Clare Matthews <sup>2,3</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/sextrans-2024-056168>).

<sup>1</sup>Nuffield Department of Medicine, University of Oxford, Oxford, UK

<sup>2</sup>Central North West London NHS Foundation Trust, London, UK

<sup>3</sup>Francis Crick Institute, London, UK

<sup>4</sup>University College London, London, UK

<sup>5</sup>Royal Free Hospital, London, UK

<sup>6</sup>Virus Reference Department, National Infection Service, UK Health Security Agency-Colindale, London, UK

## Correspondence to

Dr Sheila Lumley, University of Oxford, Oxford, UK; [sheila.lumley@trinity.ox.ac.uk](mailto:sheila.lumley@trinity.ox.ac.uk)

Received 20 March 2024

Accepted 11 May 2024

Published Online First

24 June 2024

## ABSTRACT

Diagnosing and treating chronic hepatitis B virus (HBV) infection are key interventions to support progress towards elimination of viral hepatitis by 2030. Although nucleos/tide analogue (NA) therapy is typically highly effective, challenges remain for viral load (VL) suppression, including medication access, incomplete adherence and drug resistance. We present a case of a long-term HBV and HIV coinfecting adult prescribed with sequential NA therapy regimens, with episodes of breakthrough viraemia. Multiple factors contribute to virological breakthrough, including exposure to old NA agents, initial high HBV VL, therapy interruptions, intercurrent illnesses and potential contribution from resistance mutations. The case underscores the importance of individualised treatment approaches and adherence support in achieving HBV suppression. Furthermore, it emphasises the need for improved clinical pathways addressing education, support and access to care, particularly for marginalised populations. Comprehensive data collection inclusive of under-represented individuals is crucial for maintaining retention in the care cascade and informing effective interventions.

## BACKGROUND

In line with international goals to eliminate viral hepatitis as a public health threat by 2030,<sup>1</sup> there is a global drive to diagnose, treat and prevent hepatitis B virus (HBV) infection. Nucleos/tide analogue (NA) agents suppress HBV DNA to below quantifiable thresholds in the majority of people receiving treatment. However, viraemia persists in a proportion of those offered treatment (up to 20% after 1 year in a recent population analysis<sup>2</sup>), attributed to a range of influences which may include drug resistance due to the selection of polymorphisms (resistance-associated mutations (RAMs)) in the viral reverse transcriptase (RT). Resistance is predictably selected by exposure to lamivudine (3TC), which also influences susceptibility to entecavir (ETV) and adefovir (ADV).<sup>3</sup> In contrast, resistance to tenofovir (TFV) is uncommon due to a high genetic barrier.<sup>4,5</sup>

We here describe the case of an adult in whom HBV viraemia has not been consistently suppressed on treatment, to highlight a vulnerable population with risk factors for virological breakthrough.

## KEY MESSAGES

- ⇒ Hepatitis B virus (HBV) viraemia may persist or rebound in adults receiving nucleos/tide analogue (NA) treatment; a number of factors may collectively reduce the success of treatment in mediating long-term HBV suppression.
- ⇒ We illustrate a complex case where multiple factors contribute to non-suppressed HBV viral load (VL) on NA treatment.
- ⇒ We raise awareness of diverse factors which may be contributing to non-suppressed HBV VL, and advocate for improved clinical pathways and comprehensive data collection to improve care and address knowledge gaps surrounding VL non-suppression on treatment.

## HBV CASE REPORT: PRESENTATION, INVESTIGATIONS AND TREATMENT

An adult man received treatment for HBV and HIV coinfection over a period of 26 years in a central London clinic ([figure 1](#)). We retrospectively reviewed data from his routine clinical records.

At initial diagnosis, HBV viral load (VL) was 8.5 log<sub>10</sub> IU/mL and hepatitis B 'e' antigen (HBeAg) was positive. He screened negative for hepatitis D virus infection. Antiretroviral therapy (ART) was commenced, with HIV suppression from 3.7 log<sub>10</sub> RNA copies/mL to undetectable. Subsequently, HBV was diagnosed and HBV VL progressively suppressed on HBV-active regimens containing 3TC (from baseline), switched to ADV (year 8) and then TFV (year 12), prescribed in line with changing guidelines. HBeAg was undetectable by year 13 (although he did not develop anti-HBe). He received successful intercurrent treatment for hepatitis C virus infection.

Due to illness and emergency hospital admissions unrelated to bloodborne virus infection, he was unable to sustain NA treatment, with documented gaps at intervals between years 22 and 25. HIV and HBV rebounded, with associated liver inflammation (peak alanine aminotransferase (ALT) 902 IU/L), and HBeAg status reverted to positive. On reinstating HBV-active ART (including TFV and emtricitabine), HIV VL suppressed within 1 month, but HBV VL remained persistently elevated between 3.3 and 9.9 log<sub>10</sub> IU/mL ([figure 1](#)).



© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.


**To cite:** Lumley S, Barlow M, Said K, et al. *Sex Transm Infect* 2024;**100**:329–331.

## Case report


## A. Antiviral treatment history



## B. Viral markers



## C. Liver biochemistry and elastography



## D. HBV sequencing

		Resistance associated mutations								
		I169T	V173L	L180M	A181T/V	T184G	S202I/G	M204V/I	N236T	M250V
RAM Identified by sequencing	Reference lab (Sanger)	No	No	No	Yes	No	No	No	No	No
	Research lab (Illumina)	No	No	No	97% A	No	No	No	No	No
		% sequences with RAM								
		3% T								

**Figure 1** Summary details of non-suppression of HBV VL in an adult prescribed with antiviral therapy. (A) Timeline showing antiviral treatment for HBV, HIV and HCV, showing years since diagnosis on x-axis. (B) Timeline showing trends in viral loads for HBV, HIV and HCV, in addition to CD4+ count, quantitative HBsAg (qHBsAg) and HBeAg status (NB: x-axis not to scale). (C) Timeline showing ALT and liver elastography scores (NB: x-axis not to scale). (D) Resistance-associated mutations (RAMs) listed in EASL guidance<sup>8</sup> mapped to the HBV polymorphisms identified in HBV sequenced from this individual through Sanger and Illumina sequencing. 3TC, lamivudine; ADV, adefovir; ALT, alanine aminotransferase; ART, antiretroviral therapy; EASL, European Association for the Study of the Liver; ETV, entecavir; FTC, emtricitabine; FS, fibrosan (liver elastography), HBeAg, hepatitis B 'e' antigen; HBV, hepatitis B virus; HCV, hepatitis C virus; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate; VL, viral load.

A clinical diagnostic laboratory identified the presence of a resistance mutation at position 181 in the HBV RT sequence (A181T) and reported potential drug resistance to 3TC, ADV and telbivudine. HBV Illumina sequence analysis<sup>6</sup> demonstrated dual infection with HBV genotypes A and G (online supplemental methods), and confirmed the A181T polymorphism, although only in a minority of quasi-species (figure 1B). Alongside support to optimise adherence, ETV was added (in keeping with clinical guidelines<sup>7,8</sup>), followed by an HBV virological response to 2.4 log<sub>10</sub> IU/mL after 1 month and 3.2 log<sub>10</sub> IU/mL at 4 months.

## DISCUSSION

A number of factors can collectively reduce the success of NA treatment for HBV, including high baseline VL, HIV coinfection, exposure to historical regimens with low genetic barrier to resistance, therapy interruptions, physical and mental health comorbidities and complex barriers to continuity of care. These are inter-related and are likely to have a cumulative influence.

Individuals with these characteristics may be in vulnerable or marginalised groups, are unlikely to be eligible for clinical

studies, and may experience social stigma and discrimination. These influences mitigate against their inclusion in laboratory data, clinical cohorts and trials.<sup>4</sup> There is a need for focus on equitable representation of the real-world challenges of life-long therapy to fill this current 'blind spot' (further discussed in our Editorial<sup>9</sup>).

The contribution of drug resistance is doubtful in our patient, as A181T alone is not a recognised cause of tenofovir disoproxil fumarate resistance<sup>10</sup>; a resistant phenotype would typically require multiple associated RAMs. More clinical and in vitro data are still needed to ascertain the relative contributions of different RAMs, alone and in combination. Adding HBV active agents, and providing adherence support, resulted in progressive reduction in viraemia over time to <2000 IU/mL, but not to undetectable.

Persistent HBV viraemia poses risks of transmission and long-term inflammatory/fibrotic liver disease (indicated by elevated ALT in this case), highlighting the need for intervention. Service improvements should focus on flexible, patient-centric access to information, consistent supplies of medication, avoiding

out-of-pocket costs and providing access to peer support, particularly for those coping with other health and/or social challenges and for whom there are barriers to care access. As HBV treatment eligibility expands and we work towards elimination targets, research is needed to better determine the factors that contribute to the presence and impact of persistent viraemia, and to optimise surveillance and clinical intervention.

**Handling editor** Anna Maria Geretti

**X** Philippa Clare Matthews @pipppa\_matt

**Contributors** SL and PCM are the guarantors.

**Funding** SL is funded by a Wellcome clinical PhD fellowship (102176/B/13/Z). PCM has funding from Wellcome (reference: 110110/Z/15/Z), the Francis Crick Institute and UCL NIHR Biomedical Research Centre.

**Competing interests** None declared.

**Patient consent for publication** Consent obtained directly from patient(s).

**Ethics approval** This study involves human participants and was approved by the South East Coast–Brighton & Sussex Research Ethics Committee (study title: Characterising and modifying immune responses in chronic viral hepatitis; REC reference: 11/LO/0421; IRAS project ID: 43993). Written informed consent was obtained.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

#### ORCID iDs

Sheila Lumley <http://orcid.org/0000-0001-6825-9324>

Khadija Said <http://orcid.org/0000-0001-7467-7290>

Stuart Flanagan <http://orcid.org/0000-0003-1545-8615>

Philippa Clare Matthews <http://orcid.org/0000-0002-4036-4269>

#### REFERENCES

- 1 World Health Organization: Elimination of Hepatitis by 2030, Available: <https://www.who.int/health-topics/hepatitis/elimination-of-hepatitis-by-2030> [Accessed 11 Dec 2023].
- 2 Wang T, Campbell C, Stockdale AJ, *et al.* Classification of virologic trajectories during nucleos(t)ide analogue treatment of hepatitis B virus (HBV) infection. *Infectious Diseases (except HIV/AIDS)* 2023.
- 3 Rajoriya N, Combet C, Zoulim F, *et al.* How viral genetic variants and genotypes influence disease and treatment outcome of chronic hepatitis B. *J Hepatol* 2017;67:1281–97.
- 4 Lumley SF, Delphin M, Mokaya JF, *et al.* A systematic review and meta-analysis of the risk of hepatitis B virus (HBV) genotypic resistance in people treated with Entecavir or tenofovir. *Infectious Diseases (except HIV/AIDS)* 2023.
- 5 Mokaya J, Maponga TG, McNaughton AL, *et al.* Evidence of tenofovir resistance in chronic hepatitis B virus (HBV) infection: an observational case series of South African adults. *J Clin Virol* 2020;129:104548.
- 6 Lumley SF, Jennings D, Waddilove E, *et al.* Pan-genotypic probe-based enrichment to improve efficiency of hepatitis B virus sequencing. *bioRxiv* 2023.
- 7 NICE. Guidance. Hepatitis B (chronic): diagnosis and management, 2017. Available: <https://www.nice.org.uk/guidance/CG165/chapter/1-Recommendations> [Accessed 30 Apr 2021].
- 8 European Association for the Study of the Liver (EASL). Clinical practice guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017;67:370–98.
- 9 Im YR, Said K, Martyn E. Social, clinical, and biological barriers to hepatitis B virus suppression with nucleos(t)ide analogue therapy: who is at risk and what should we do about it. *Sex Transm Infect* 2024;100:259–63.
- 10 Zoulim F, Locarnini S. Hepatitis B virus resistance to Nucleos(t)ide analogues. *Gastroenterology* 2009;137:1593–608.