SOME ASPECTS OF GONORRHOEA IN THE FEMALE—WITH SPECIAL REFERENCE TO INFECTION OF THE RECTUM*

BY

C. S. NICOL

*Medical Officer in Charge, Venereal Diseases Department, St. Bartholomew's Hospital;
and Chief Assistant Medical Officer, Whitechapel Clinic*

Gonococcal infection in the female presents many difficult problems, both in diagnosis and treatment. It is therefore to be regretted that since the advent of chemotherapy there has been a widespread belief that the problem of the control of this disease has almost ceased to exist. In consequence diagnostic standards are apt to be less stringent, and tests of cure are less exacting in relation both to length and frequency of observation. After an adequate dose of penicillin cure is practically taken for granted and further surveillance concentrates on eliminating the possibility of suppressed syphilitic infection.

It is in diagnosis and by contact tracing that the first and greatest efforts must be made to suppress gonorrhoea. In diagnosis it is essential to include in clinical and bacteriological investigations every method which will increase the chance of detecting gonococcal infection. It is also true that however high the standards of diagnosis may be the venereologist fails in his task if he does not make every effort to trace all contacts of the patient, whether infecting or infected. For this purpose he must make full use of the existing regulations, and of the services of the social service worker if one is attached to the clinic.

To illustrate the importance of these various considerations an analysis has been made of 460 female patients attending the Whitechapel Clinic over a period of twelve months (from July 1, 1946, to June 30, 1947). Details will be given of 229 patients with acute gonorrhoea, of 66 patients in whom only the gonococcal complement fixation test gave a positive finding, of 35 patients with negative tests who were known consorts of males infected with gonorrhoea, and of 130 patients with a trichomonas infection and negative tests for gonorrhoea. It is proposed to discuss the cases under the three following

headings: (1) contact tracing; (2) diagnosis; (3) treatment.

An analysis of the case histories of 229 patients attending with acute gonorrhoea showed that 159 were married women; the remaining 70 were single. All these patients admitted sexual intercourse. One hundred and fifty-nine patients were under thirty years of age, while of the remaining 70 over 30 years of age, 3 were over 50 (Fig. 1).

Contact Tracing

An attempt has been made to analyse the contacts of these cases. The married women have been

FIG. 1

* An Address to the Medical Society for the Study of Venereal Diseases, Oct. 25, 1947.

divided into those living with their respective husbands, and those who had been widowed, divorced, or separated from their husbands. Each of these groups has been further subdivided into three groups: (i) the husband or contact had attended the Whitechapel Clinic; (ii) the husband or contact had attended with gonorrhœa at another clinic; (iii) no definite details had been obtained of the husband or contact. A further analysis has been made in the cases in which the husband or contact attended at Whitechapel, according to whether the man or the woman has attended initially with symptoms and been given a contact slip for the other; and finally details of the diagnosis of the contact or husband have been collected. The same comparable method of analysis has been used for the single patients (Table I). It will be seen that efforts to trace the husband or contact were unsuccessful in only 57 cases of the series.

Of the 110 married patients living with their husbands only 5 admitted to extramarital intercourse. In some cases the husband admitted extramarital intercourse, but in most instances neither husband nor wife made such an admission. Of the 70 single patients only 2 admitted intercourse with more than one contact, and the 49 patients separated from their husbands all admitted relations with only one sexual partner.

Table II illustrates the findings in all cases in which contact slips from the female side of the clinic were brought by male patients, and vice versa. Thirty-two female patients gave contact slips either to husbands or contacts. In 26 cases the male patient had gonorrhœa, while in 6 cases all tests were negative.

One hundred and forty-four male patients with acute gonorrhœa gave contact slips to their wives or contacts, and subsequent examination showed that in 99 cases the woman had acute gonorrhœa,

TABLE I
CONTACT TRACING

Married : 159	Living with husband : 110	Husband attended White- chapel clinic : 80	Husband gave wife contact slip : 63	Husband Gc+ : 61
			Wife gave husband contact slip : 17	Husband N.V.D. urethritis : 2
		Husband attended with gonorrhœa at another clinic : 18		Husband Gc+ : 13
		No details of husband : 12		Husband N.V.D. : 4
	Separated from husband, widowed, or divorced : 49	Contact attended Whitechapel clinic : 24	Contact gave patient contact slip : 18	Contact Gc+ : 18
			Patient gave contact contact slip : 6	Contact N.V.D. : 0
		Contact attended another clinic with gonorrhœa : 7		Contact Gc+ : 5
		Contact not traced : 18		Contact N.V.D. : 1
Single : 70		Contact attended Whitechapel clinic : 29	Contact gave patient contact slip : 20	Contact Gc+ : 20
			Patient gave contact contact slip : 9	Contact N.V.D. : 0
		Contact attended at another clinic with gonorrhœa : 14		Contact Gc+ : 8
		Contact not traced : 27		Contact N.V.D. : 1

TABLE II

RESULTS OF PATIENTS ATTENDING WITH CONTACT SLIPS

Female contact with gonorrhoea : 32	Male contact in Whitechapel clinic with gonorrhoea... ..	26
	Male contact in Whitechapel clinic with negative tests ..	6
Male contact with gonorrhoea : 144	Female contact in Whitechapel clinic with gonorrhoea ..	99
	Female contact in Whitechapel clinic with positive gonococcal complement fixation test ..	15
	Female contacts in Whitechapel clinic with negative tests :	
	Repeated negative tests over 3 months ..	15
	Salpingitis ..	1
	History of gonorrhoea ..	2
	Defaulted before ..	
	surveillance completed ..	12
		30

in 15 cases only the gonococcal complement fixation test was positive, and in 30 cases all tests were negative. A further analysis of these thirty cases shows that in only 15 was there a full and adequate follow-up with repeated tests. Of the remaining 15, 1 developed salpingitis, 2 gave a history of past gonorrhoea, while 12 defaulted soon after first attendance.

Only three patients of the *whole series* attended as a result of notification under Regulation 33B.

Diagnosis

Symptoms.—One hundred and forty patients attended the clinic on the advice of their male consort, who had developed acute gonorrhoea. At first attendance 76 of these women admitted no

symptoms; 45 complained of vaginal discharge; 15 complained of dysuria; 1 of abdominal pain, and 1 of pain in a joint.

Eighty-nine patients attended the clinic of their own accord. On questioning, 30 admitted no symptoms relating to gonorrhoea, 44 complained of vaginal discharge, 8 of dysuria, 4 of abdominal pain, 2 complained of pain in a joint, and 1 of vulval swelling which proved to be due to Bartholinitis. These figures relate to the presenting symptom in each case; but many patients complained of additional symptoms in varying combination. In Table III the analysis of symptoms is given, indicating their occurrence in patients with and without an additional *Trichomonas vaginalis* infection. The main point which becomes apparent is that 106 patients out of a total of 229 had no significant symptoms; 140 patients out of 229 attended the clinic only at the request of their male consorts.

Smear and Culture Findings.—During the first seven months of the period under consideration hydrocele agar (Thomson, 1923) was used as the culture medium except in the vaginal plate; but, due to difficulties in obtaining hydrocele fluid, serum enriched medium was under trial during the last five months of the period. As subsequent figures will indicate, this medium was not a complete success, and the clinic has again reverted to the original hydrocele medium. Considering the period during which the hydrocele medium was in use, the culture results were comparable with smear results, but nowhere near the level which has recently been claimed by McLeod (1947). Fig. 2 compares smear and culture results, giving the totals of positive findings in the urethra, cervix and vagina, and Fig. 3 shows the number of cases in which the diagnosis was made either by smear or culture alone. It also indicates the number of cases in which positive culture was obtained by the vaginal plate technique (King and Mascall, 1935) alone. As previously stated, hydrocele medium was not used for this purpose. A division has been made to indicate the varying results obtained using the two culture media. The diagnosis

TABLE III

	Patients who attended at request of consort			Patients who attended of their own accord		
	Trichomonas present	Trichomonas absent	Total	Trichomonas present	Trichomonas absent	Total
No symptoms suggesting gonorrhoea ..	33	43	76	10	20	30
Vaginal discharge ..	27	18	45	17	27	44
Dysuria ..	3	12	15	6	2	8
Abdominal pain ..	1	2	3	2	2	4
Arthritis, etc. ..	1	0	1	2*	1	3
Total ..	65	75	140	37	52	89

* 1 Bartholinitis

FIG. 2

FIG. 3

FIG. 4

FIG. 5

of gonorrhœa was made from the first series of tests in 185 cases, from the second series in 28 cases, from the third series in 11 cases, and in subsequent tests in 5 cases, indicating that in 19.1 per cent. of the cases the diagnosis was made by repeating tests which originally gave negative smear and culture results.

Duration of Infection.—Cases have been divided according to the presumed duration of infection, which is often difficult to determine in women. It was decided upon partly by the length of symptoms, and partly by the dates of exposure to infection. In 84 cases (37.7 per cent.) infection was of less than two weeks duration; in 66 (28.8 per cent.) from two to four weeks; in 79 (34.5 per cent.) over four weeks. In an analysis of 544 male patients attending the Whitechapel Clinic in the first half of 1947, 447 (82.2 per cent.) had evidence of infection of less than two weeks duration; 88 (16.1 per cent.) from two to four weeks; and only 9 (1.7 per cent.) over four weeks. This emphasizes the point that the male patient usually reports to a clinic very much sooner after infection than the female. (These figures are shown in Fig. 4, expressed in percentages.)

Fig. 5 indicates the rising incidence of positive results of the gonococcal complement fixation test in relation to the length of infection. This may be compared with the figures which Price quoted in a paper in 1933. He found the gonococcal complement fixation test positive in 46 per cent. of cases after the second week, and in 89 per cent. of cases

TABLE IV

	Patients with positive gonococcal complement fixation test only	Additional trichomonas infection
Given large dose of penicillin for concomitant syphilis	24	12
Previously treated for acute gonorrhœa elsewhere ..	16	4
Less than three local tests performed	11	5
Complications usually associated with gonorrhœa— Salpingitis: 2 .. Arthritis: 3 ..	5	1
Cases fully observed for three months with negative local tests	10	6
Total	66	28

TABLE V

Incidence of trichomonas infection	No. of cases	Per cent.
Gonorrhœa only	127	55.5
Gonorrhœa and trichomonas vaginitis	102	44.5
Syphilis only (early and early latent)	86	80.7
Syphilis and trichomonas vaginitis	28	19.3
Cases with positive gonococcal complement fixation test only ..	38	57.6
Cases with positive gonococcal complement fixation test and trichomonas vaginitis	28	42.4

after the fourth week of the disease, in untreated male and female cases.

Gonococcal Complement Fixation Test.—The finding of a positive gonococcal complement fixation test, without positive results to smear or culture, always presents an interesting problem. An analysis of 66 such patients showed that in 24 cases large doses of penicillin had been given immediately for a concomitant syphilitic infection, in 16 cases there was a recent history of acute gonorrhœa treated at another clinic, in 5 cases there occurred complications which are often associated with gonorrhœa (arthritis in 3 and salpingitis in 2). In 11 cases two series or fewer of local tests had been performed, while in the remaining 10 cases adequate series of local tests over several months were negative. Nevertheless, in 6 of these 10 cases *Trichomonas vaginitis* was found, and of the other 4, 2 were contacts of gonorrhœa and 1 of syphilis. Thus in the majority there was either a history of treated gonorrhœa or evidence of complications associated with the disease; or penicillin treatment for syphilis had inhibited possible positive findings; or an inadequate number of tests had been performed in defaulting patients (Table IV).

Trichomonas Infection.—In 102 of the 229 female patients under review there was an additional infection with *Trichomonas vaginalis*. The diagnosis was made by examination of saline suspensions of vaginal discharge obtained from the posterior fornices. This test was made not only at the patient's first attendance but was repeated on each subsequent examination. In 53 cases the *T. vaginalis* was found before the administration of penicillin for the gonococcal infection, while in 49 cases it was demonstrated only after penicillin therapy. This corresponds with the clinical observation that the patient often complains of increased

TABLE VI

130 FEMALE PATIENTS WITH TRICHOMONAS VAGINITIS AND WITH NO CLINICAL OR LABORATORY EVIDENCE OF GONORRHOEA

Contacts :	
Gonorrhoea	15
Positive gonococcal fixation test	2
Non-specific urethritis	4
Syphilis	11
? a venereal disease at another clinic	5
Negative tests	6
Not traced	87
Associated conditions :	
Syphilis (early)	28
Syphilis (late)	4
Warts	3
Moniliasis	7
Pregnancy	3
Past history of gonorrhoea	14
Nil	71
Symptoms :	
Vaginal discharge	83
Dysuria	3
Vulval irritation	4
No symptoms	40

vaginal discharge after the administration of penicillin. Table V compares the incidence of trichomonas infection in gonorrhoea with its incidence in early and early latent syphilis in the Whitechapel Clinic over the same period, showing in addition its incidence in cases in which only the gonococcal complement fixation test was positive. It will be seen that in both gonorrhoea, and in presumptive gonorrhoea, the incidence is very similar, while in syphilis it is significantly lower. It should be added that at least two examinations for *T. vaginalis* were made in all cases of syphilis, one before and one after penicillin therapy.

On the other hand during the same period 130 female patients attended with a trichomonas infection while the smear and culture tests and fixation tests for gonorrhoea were consistently negative.

A brief analysis of these cases has been made, indicating presenting symptoms, associated conditions, and diagnosis of known contacts (Table VI).

Reaction of Vaginal Secretion.—Using a Universal indicator, initial pH readings of the vaginal fluid were recorded in 54 cases. These are indicated in Fig. 6 and divided into two groups showing the effect of the presence or absence of trichomonas infection, and secondly the effect on the duration of the gonococcal infection. The pH readings have been arbitrarily grouped into those below and those above a pH reading of 5.2. This appears to demonstrate that abnormality of the pH readings of the vaginal fluid depend rather on length of gonococcal infection than the presence or absence of a trichomonas infection. It must be remembered that this is a rough quantitative test and that there are other possible sources of error, such as vaginal fluid mixed with cervical secretion or menstrual blood. The reaction may also be affected by the recent deposit of semen or by the use of a chemical contraceptive.

Moniliasis.—In 18 of the 229 patients with gonorrhoea the yeast forms of monilia were found in the vaginal slide, or the organism was cultured on Sabouraud medium. In five of these cases a clinical monilial vaginitis was diagnosed with the presence of yeast forms and mycelium from a patch of vaginal thrush, or from a scraping of the vulval skin. In four of these five cases there had also been a *Trichomonas vaginitis*.

FIG. 6A

FIG. 6B

Complications.—The following complications were found to be present at the time of diagnosis in the 229 patients with acute gonorrhœa. One patient had skenitis, 5 had bartholinitis, 6 had salpingitis, 2 had arthritis; while in 26 an ano-rectal infection was discovered.

Cases of gonococcal proctitis in women were probably described by Hecker (1801). Gonococci were first demonstrated in a rectal smear by Bumm in 1884 (Yeomans, 1929). In the past fifteen years Martin (1935), Brunet and Salberg (1936), Ault (1937) in America, and Ruys (1934, 1939) in the Netherlands have all discussed this condition. Papers have also been published in Germany, France, Sweden, and Russia. Little interest appears to have been taken in the subject in this country since Clements and Hughes' (1935) excellent paper describing 69 cases with positive rectal findings out of 160 female patients tested. They made the following main points:

1. In the majority of cases there were no signs or symptoms.
2. Cultural methods were important. (They used hydrocele agar at pH of 7.6. either on slopes or in plates.)
3. In seven cases gonococci were found only in the rectum.

4. In eight cases gonococci were found in the rectum at final tests. A proctoscope was not used.

The technique used in this clinic has been to pass a proctoscope, of the Graeme Anderson type, using a little vaseline as a lubricant, after routine tests, making certain that no vaginal discharge flows across the perineum before it is passed. Specimens have been taken with a platinum loop for smear and cultural examinations. Occasionally faeces in the lower rectum obscure the examination, and ideally the bowel should have been previously emptied, but this is not always possible with out patients. As the tests are being taken the rectal wall—as far up as the proctoscope reveals it—should be carefully observed; because, although in some cases the infection is obvious, with pus present on the rectal mucosa—which appears red and congested—in the majority of cases the mucosa appears normal; and only on careful examination is a small flake of stringy mucus picked off in the loop. The anal canal and ano-rectal junction usually appear normal.

Tests taken in this series were examined by smear and culture. In the smear a positive diagnosis should only be made when gram-negative intracellular diplococci are seen in the presence of numerous pus cells. In many positive slides there are surprisingly few other organisms present. This appears to be particularly so in the acute stage of the infection. Serum-enriched medium on slopes or in plates was used for culture. After incubation

TABLE VII

Case No.	Rectal symptoms	Vaginal trichomonas	Vaginal discharge	Rectum		Urethra smear	Cervix smear	Vaginal plate	Urethra culture	Cervix culture	G.F.T.
				Smear	Culture						
1	Nil	—	Nil	+	—	+	—	+	+	+	—
2	Discharge	—	+	+	—	+	—	—	+	+	±
*3	Discomfort	—	—	+	—	—	—	—	—	—	—
4	Discharge	+	+	+	—	+	+	—	—	—	—
5	Discharge	+	+	+	+	+	+	—	—	+	—
*6	Nil	—	+	+	—	—	—	—	—	—	+
*7	Nil	+	+	+	—	—	—	—	—	—	+
*8	Nil	+	—	+	—	—	—	—	—	—	—
9	Nil	+	—	+	—	—	+	—	—	—	±
10	Nil	+	+	+	—	+	+	—	+	—	++
11	Discharge	—	—	+	—	—	—	—	—	+	+
12	Nil	+	+	+	—	+	+	—	—	—	—
13	Nil	+	+	+	—	+	+	—	—	+	—
14	Nil	+	+	+	—	+	+	—	—	—	—
15	Nil	+	—	—	+	—	—	+	+	—	—
16	Discharge	—	+	+	—	+	+	—	+	+	++
17	Nil	+	+	+	—	+	+	—	—	—	++
*18	Nil	—	—	+	—	—	—	—	—	—	+
19	Nil	—	+	+	—	+	—	—	—	—	±
20	Nil	—	+	—	+	—	—	+	—	+	+
21	Discharge	—	+	+	—	+	+	—	—	—	++
22	Nil	—	—	—	+	—	—	+	+	+	—
23	Nil	—	—	+	—	+	—	—	—	—	±
24	Nil	—	+	+	—	+	+	—	—	—	+
25	Bleeding	+	+	+	+	+	—	—	+	—	++
26	Nil	—	—	+	—	—	+	—	—	—	++

* Diagnosis of gonorrhœa made from the rectal findings alone

colonies were first differentiated by the oxidase reaction, and smears of oxidase-positive colonies examined. Subculture was attempted in cases with positive results, and if successful in pure culture sugar fermentation reactions were performed.

Seventy-four consecutive patients in the latter part of the main series were examined rectally by this technique. In 2 patients both smear and culture findings were positive, in 21 the smear only was positive, while in 3 the culture only was positive.

Only 8 of the 26 patients with proctitis admitted to any symptoms, and in all these cases the symptoms were admitted only when the patient was questioned again after the diagnosis of proctitis had been made, so it must be assumed that they were of minor severity. Patients complained either of rectal discharge and irritation, or of discomfort or bleeding when the bowels were opened.

In 12 of the 26 patients there was an additional vaginal trichomonas infection. In 13 cases vaginal discharge was profuse, while in the other 13 it was minimal. Fifteen patients were nulliparous, while of the remaining 11 only 4 had had more than one confinement. Thus it does not appear that the presence of trichomonas infection, or a profuse vaginal discharge, or the after-effects of childbirth are very important factors in the production of rectal infection. Table VII shows these findings in full, and it will be observed that in 5 cases the diagnosis of gonorrhoea was made from the rectal findings alone. In 3 of these 5 cases the gonococcal complement fixation test was positive, and in the other 2 cases the male contact had gonorrhoea, and in 1 of these there was a history of another male contact contracting gonorrhoea one and a half years previously. In 2 cases in the series there was a past history of gonorrhoea.

In an additional 14 patients of the 74 examined there was some evidence of rectal infection, as indicated by the presence of leucocytes in the rectal

smear, but smears and cultures were negative. Table VIII summarizes these figures, and also indicates the incidence of proctitis in relation to length of infection. It appears that the incidence tends to rise the longer patients remain undiagnosed.

Treatment

A routine treatment consisting of 30,000 units of sodium penicillin in water given intramuscularly at two-hourly intervals for five injections was given to 154 of the 229 patients with acute gonorrhoea; 17 patients were given no penicillin because they defaulted; 3 defaulted after only one injection, while 8 were given sulphathiazole. Thirty-two patients were given larger doses of penicillin up to 2.4 mega units because of complications of gonorrhoea (arthritis and salpingitis), or concomitant syphilis. The remaining fifteen patients were given penicillin in ethyl-oleate and beeswax in a single dose of 200,000 units, as they could not attend several times during the day.

It is intended only to give details of results of treatment in the main group of cases, that is, 144, excluding 10 cases with a past history of gonorrhoea, while a further analysis of the treatment results of the patients with proctitis will be made, irrespective of which treatment group they represent.

Tests of Cure.—Smears and cultures from the cervix

TABLE IX

RESULTS OF TREATMENT

Surveillance	Gonorrhoea only		Gonorrhoea with trichomonas vaginitis	
	Pus cells — or ± in smear	Pus cells + or ++ in smear	Pus cells — or ± in smear	Pus cells + or ++ in smear
Defaulted immediately ..	Nil	7	Nil	4
Less than 3 months ..	5	11*	2	14 (6)
3 to 6 months ..	4	10	3	10*
Over 6 months ..	8	9†	6	13†(7)
? Reinfection ..	4		5	
? Relapse ..	7		8	
Gonococcal complement fixation test persistently positive ..	2		3	
Developed syphilis — further penicillin ..	1		—	
Oxidase-positive organism —not gonococci ..	4		4	

() still trichomonas present.

* 1 admitted sexual intercourse

† 2 admitted sexual intercourse

TABLE VIII
RECTAL GONORRHOEA

No. of cases tested	74
Rectal smear positive	21
Rectal culture positive	3
Rectal smear and culture positive ..	2
Pus cells but no gonococci in smear, and negative culture	14
Negative smear and culture	34

Duration of infection	Rectal tests positive	Rectal tests negative
Under 2 weeks	7	21
2 to 4 weeks	8	20
Over 4 weeks	11	7

and urethra, and vaginal cultures were taken twice weekly for a month after treatment; then tests were repeated immediately after the menstrual period each month for a further two months, and at the end of the sixth month, with Wassermann, Kahn, and gonococcal complement fixation tests at six weeks, three months, and six months. In the cases with rectal gonorrhœa, rectal smear and culture were also taken at the first test after treatment, and at subsequent tests as indicated clinically; and as a routine after three months and six months. Recently rectal smears and cultures have been taken as a routine at the final tests of all patients.

The 144 patients have been divided into two groups; 72 cases with an additional trichomonas infection. In Table IX each group is divided according to the length of observation, showing in addition whether the number of leucocytes in the various smears were minimal or increased in number. Those cases with only the gonococcal complement fixation test remaining positive, and with supposed re-infection and supposed relapse are also indicated.

During tests of cure in 6 cases not included under any of these headings an organism morphologically

similar to the gonococcus was grown on culture, only differing in that it was rather small. However, pure subcultures in 5 of these cases gave negative sugar fermentation reactions. A similar organism has been described in the past by Thomson.

There were 24 treatment failures in this series; in 19 cases the gonococcus was found in the tests; salpingitis followed in 4 cases and Bartholinitis in 1. Table-X shows these cases in detail, indicating supposed re-infection or relapse. In 16 of the 24 cases the male contacts were also attending the clinic, so closer correlation was possible.

It is impossible to distinguish re-infection from relapse of a gonococcal infection, except by the patient's statement in relation to further sexual intercourse. But even if further sexual contact has occurred it may act merely as a provocation to a latent infection rather than as a cause of fresh infection. It is interesting to note that in cases of supposed relapse the diagnosis is almost invariably made by culture, although the reason for this is by no means clear. Thus in the tests of cure in this

TABLE X

Gonorrhœa only					
	Case No.	Exposure	Gonococci present	Complications	Interval since penicillin
? Reinfection	1	Admitted	—	Bartholinitis	1 month
	2	"	In smear	—	4 months
	3	"	" *	—	2½ months
	4	"	In culture	—	3 months
? Relapse	1	Denied	In culture	—	1 week
	2	"	"	—	1 week
	3	"	"	—	2 months
	4	"	"	—	3 weeks
	5	"	"	—	3 weeks
	6	"	Nil	Salpingitis	7 weeks
	7	"	"	"	6 weeks
Gonorrhœa and trichomonas infection					
? Reinfection	1	Admitted	In smear	—	4 months
	2	"	"	—	3 weeks
	3	"	In culture	—	3 weeks
	4	Not admitted†	In smear	—	3 months
	5	Admitted	In culture	—	1 month
? Relapse	1	Denied	In culture‡	—	2 weeks
	2	"	In smear	—	1 month
	3	(Alcohol+) Denied	In smear	Salpingitis	3 weeks
	4	"	In culture	—	6 months
	5	"	—	Salpingitis	6 weeks
	6	"	In culture	—	5 days
	7	"	"	—	3 weeks
	8	"	"	Salpingitis	5 weeks

* Rectal smear

† Social service worker reported numerous contacts

‡ Rectal culture

series culture methods have shown a return of gonococcal infection in 12 out of 24 cases, while diagnosis by smear occurred in only 8, of which 6 were probably re-infections. The majority of the supposed relapses occurred in the first 6 weeks after treatment. Gonococci were found in the rectal smear in 1 case of supposed re-infection, and by rectal culture in 1 case of supposed relapse. In neither of these cases had gonococci been found in the rectum in earlier tests, and they are not included in the 26 cases already described.

Of these 26 patients with rectal gonorrhoea, 2 defaulted before treatment, 18 were treated with the routine dosage of 150,000 units of penicillin, 5 were treated with 2.4 mega units of penicillin for concomitant syphilis, while one was given 200,000 units of penicillin in ethyl-oleate. This last patient defaulted immediately, so in Table XI only the first two groups are considered. These have been classified by the same method as in the main treatment series.

Discussion

Contact Tracing.—This all-important aspect of the disease requires the services of a social service worker, who should interview every new patient. In this clinic it would have been impossible to obtain the results described without this aid. Ideally there should be more than one such worker, so that interviewing in the clinic and visiting duties could be carried out at the same time.

Diagnosis.—The fact shown in this paper, that gonococcal infection in the female is so often symptomless, has been stressed by writers many times in the past, but it still appears that some practitioners do not appreciate this very important fact. It may be added, also, that signs of the disease are often minimal, even though the patient remains infectious.

On this account it is vital to use every means

available in a routine of testing to diagnose gonococcal infection. The results obtained by culture are of the utmost importance. The difficulties in achieving a first-class standard of culture do not fall within the scope of this paper, but they are only too well-known to all venereal disease pathologists. During the period which this paper covered the results with an experimental medium were not good, but they were considerably better when a hydrocele agar medium was used.

The incidence of trichomonas infection in the gonococcal cases of this series is very similar to that quoted in a recent paper by King and Gallagher (1946), in which they found that 135 out of 266 female patients with gonorrhoea also had a trichomonas infection.

The search for clinical evidence of various complications by examination of the external genitalia and by bimanual examination, which indicates examination by smear or cultural methods from the glands of Skene and Bartholin, is part of the routine of all clinics. But examination of the anal canal and lower rectum by proctoscopy, and the taking of rectal smears and cultures is rarely performed. Table XII indicates that the findings in this paper confirm those of earlier workers that gonococcal proctitis in women is a very common complication. It is generally accepted that this is due to a direct spread from the vagina, but it should be remembered that in some cases rectal coitus is the cause of the complication, and may well be the cause of the earlier infections. However, admission of rectal intercourse is never easy to elicit from a female patient. In three cases of gonococcal proctitis after this series was completed rectal coitus was admitted, and in one of these cases the male consort gave the history of symptoms appearing after a single rectal coitus with a woman with whom he had cohabited for three months. She was found, on examination, to have gonorrhoea limited to the ano-rectal region. It is considered

TABLE XI
RECTAL GONORRHOEA: TREATMENT OF TWENTY-SIX CASES

1. Penicillin in water 30,000 units in five injections at 2-hourly intervals .. 18							
2. Penicillin in water 40,000 units in sixty injections at 3-hourly intervals 5							
3. Penicillin in oil 200,000 units in a single injection 1							
4. No penicillin 2							
No. of patients	Defaulted immediately	Under three months	Three to six months	Over six months	G.F.T. persist. +	?Reinfection	? Relapse
1. 18	2	6 (1)	3 (1)	1 (1)	1*	1	4
2. 5	1	—	—	1	1	1	1
3. 1	1	—	—	—	—	—	—

() Still pus in rectal smear

* Rectal ulceration lymphogranuloma inguinale

TABLE XII
ANO-RECTAL INFECTION

	No. of cases examined	Cases with positive results (%)	Culture	Method
Martin (1935)	111	30	Not done	By urethroscopy
Brunet and Salberg (1936) ..	250	38	Not done	By proctoscopy
Clements and Hughes (1935) ..	160	43	Done	By platinum loop
Ruys (1934)	201	55	Done	By platinum loop
This paper	74	37	Done	By proctoscopy

that the use of a proctoscope is essential, both in obtaining adequate specimens of material, and in assessing the condition of the rectal mucosa. It can be passed with the minimum of discomfort to the co-operative patient.

In this series the most important justification for routine examination and testing of the ano-rectal region is that the diagnosis of the disease itself depended upon these rectal findings in 5 of the 26 cases. Culture results in relation to rectal infection were disappointing, but it was unfortunate that these coincided with a time when the general culture standard was not good, and they compare unfavourably with those obtained by Clements and Hughes. The problem is an important one, because in the tests of cure culture methods are essential. A summary is given of the main diagnostic points concerning rectal infection in the female.

Gonococcal Infection of the Ano-Rectal Region

1. Common in female. Rare in male.
2. Method of infection : (a) direct spread from urino-genital tract ; (b) rectal coitus ; (c) accidental infection.
3. Infection in some cases before second week ; in most cases after second week.
4. The majority of patients complain of no symptoms. Symptoms are only elicited on direct questioning. They are usually rectal irritation or discharge, or bleeding per rectum.
5. The signs are usually minimal, with only a small piece of muco-pus on a normal mucosa. In a few cases there is a large amount of pus on the rectal wall, with congestion of rectal mucosa.
6. Ulceration of the rectal mucosa is rare. The possibility of lymphogranuloma venereum must be remembered.
7. Diagnosis by smear is usually not difficult ; in many cases there are few secondary organisms present. A positive diagnosis should be made only in the presence of numerous pus cells.
8. Culture diagnosis is important, especially after treatment ; but the technique of culture is difficult.

9. Examination by proctoscope is the best method ; In a co-operative patient it is not difficult to pass.

10. Diagnosis of gonorrhœa is made in some cases by means of the rectal findings only.

11. Rectal infection is probably not always self-limiting. It may remain as a focus for re-infection of the urinogenital tract, or for metastatic infection.

12. The clinical results with parenteral penicillin appear to be initially satisfactory, but tests of cure show that in some cases the infection persists. Better cultural methods might reveal a higher percentage of failure.

13. Rectal testing should always be a part of the tests of cure.

14. Additional local treatment may be necessary, but this has not been assessed in the present series.

Treatment.—The results of treatment of a group of female patients with a first attack of gonorrhœa using a routine dosage of penicillin have been described. It is always extremely difficult to give a correct interpretation of treatment results. The treatment analysis in this paper is primarily intended to stress the following facts, and to illustrate how impossible it is to give an exact percentage cure rate.

1. Cultural diagnosis is never a simple matter even for the most experienced pathologist. An oxidase-positive organism, very similar morphologically to the gonococcus, may be differentiated by the sugar fermentation reactions—always provided the colonies can be successfully subcultured in pure culture.

2. When the gonococcus is found in smear or culture after treatment it is really impossible to prove that this is due to a relapse and not to a re-infection, unless the patient be a prison inmate.

3. A trichomonas vaginitis which is also present may resist treatment, so that the assessment of the tests of cure for gonorrhœa is difficult.

4. In the absence of a trichomonas infection, tests of cure may give negative culture findings, but the smears may contain numerous leucocytes either

in the cervix, urethra, or rectum. The patient may also admit further sexual intercourse.

5. The gonococcal complement fixation test may remain positive.

6. There is always the possibility of variation in the potency of the penicillin used.

Thus the results of treatment might be open to many various interpretations. Taken at the worst the results are not impressive, indicating a high failure rate.

The results of treatment in the small series of cases with proctitis are affected by the above-mentioned variations. It can only be said that rectal testing is as vital in the tests of cure as in diagnosis, and that there is great scope for the improvement of such methods.

Summary and Conclusions

A study is made of various groups of patients attending the Whitechapel Clinic over a period of one year.

These cases are analysed under the main headings of (1) contact tracing, (2) diagnosis, and (3) treatment.

Special reference is made to the importance of gonococcal infection of the ano-rectal region.

I should like to thank Mr. A. J. King, the Director of the Whitechapel Clinic, for his encouragement and help during the writing of this paper, and to acknowledge the co-operation of Dr. I. N. Orpwood Price and Dr. A. E. Wilkinson in the laboratory work involved.

REFERENCES

- Ault, G. W. (1937). *Amer. J. Syph.*, **21**, 430.
 Brunet, W. M., and Salberg, J. B. (1936). *Ibid.*, **20**, 37.
 Clements, P. A., and Hughes, K. E. A. (1935). *Lancet*, **2**, 18.
 Hecker, A. F. (1801). "Deutliche Anweisung die venerischen Krankheiten genau zu erkennen und richtig zu behandeln, Erfurt." G. A. Keyser.
 King, A. J., and Gallagher, E. (1946). *Lancet*, **1**, 916.
 —and Mascal, W. N. (1935). *Ibid.*, **1**, 1492.
 McLeod, J. W. (1947). *Brit. J. vener. Dis.*, **2**, 53.
 Martin, C. L. (1935). *J. Amer. med. Ass.*, **104**, 192.
 Price, I. N. O. (1933). "The Complement Fixation Test for Gonorrhoea." London County Council.
 Ruys, A. C. (1934). *Ned. Tijdschr. Geneesk.*, **78**, 5533.
 —(1939). *Indian J. vener. Dis.*, **5**, 81.
 Thomson, D. (1923). "Gonorrhoea." London. Henry Froude and Hodder and Stoughton.
 Yeomans, F. C. (1929). "Proctology." New York and London. D. Appleton Co.

DISCUSSION ON GONORRHOEA IN THE FEMALE

DR. G. L. M. McELIGOTT said he was delighted to hear of Dr. Nicol's success with the contact slip method of contact tracing, which he probably saw for the first time at St. Mary's Hospital and from which very good results were still obtained. One of the reasons why the results of the treatment of gonococcal proctitis with penicillin in Dr. Nicol's cases were so depressing might be the fact that a very large amount of penicillinase was produced in the rectum by the *B. Coli* and it was almost an indication for concurrent sulphonamide treatment in such cases.

DR. ROBERT LEES said that Dr. Nicol's case follow-up and contact-tracing aroused his admiration. It was very difficult indeed, even with the help of an experienced social worker, to get contact-tracing such as had been reported.

He had occasion recently to question a small boy of 12 who had contracted acute gonorrhoea. This led to proof that his consort was a small girl of 9 years. The girl's mother was an old patient of the clinic; she was probably a prostitute, and she had also a lodger, a man who was suffering from both gonorrhoea and syphilis. The story was that the lodger had interfered with the small girl and she had subsequently seduced the small boy.

He was interested to find out who questioned the

patients, as female patients were often reluctant to talk to a man about their sexual affairs; many of them would admit to one sexual exposure or one sexual partner, but it was exceedingly difficult to get them to admit to the promiscuity to which one was accustomed in the male, although one was often satisfied that they were completely "farmyard" in their behaviour.

Regulation 33B was not mentioned, and it coincided with his experience that this regulation was not effective in tracing contacts.

A positive contribution of the paper was the emphasis on the high incidence of anal and rectal infections, and the fact that these could be present but unsuspected and could only be found after extensive examinations. He had not found the same percentage that was described, probably because he had not looked for it. Anal infection was common in vulvovaginitis in young girls. For ano-rectal infections he had always used local treatment in addition to specific treatment, using irrigations with potassium permanganate and a suppository containing a flavine compound. These were well tolerated and results of treatment were satisfactory.

MR. A. J. KING was sorry to hear that Dr. Lees did not use the method of vaginal plate culture. It was ten years since that method was recommended by Dr. Mascal and himself, and the test had received less attention than