PROSTITUTION AND VENEREAL DISEASES IN MANCHESTER*

BY

SYDNEY M. LAIRD

Director, St. Luke's Clinic and V.D. Department, Royal Infirmary, Manchester

The relationship between prostitution and the spread of venereal disease probably varies from country to country and, indeed, between the various communities of one country. Many believe that prostitution plays little part in the spread of venereal infection in England (British Social Biology Council, 1955), although Nicol (1956) is of the opinion that the "reservoir" of undetected gonorrhoea in females comprises both the prostitute and the promiscuous "amateur". My own experience suggested that prostitution is an insignificant factor in the smaller provincial towns and rural communities, but it assumes a significant role in the largest cities especially when they are also seaports. This impression was supported by the relevant data from the Gonorrhoea Study of the British Cooperative Clinical Group (B.C.C.G.) modified in Table I.

TABLE I
MALE CASES OF GONORRHOEA INFECTED BY
PROSTITUTES (MODIFIED FROM TABLE IV OF
THE B.C.C.G. GONORRHOEA STUDY)

Source of Infection	Group				Total
	I. Ports	II. Inland Industrial Cities	III. London	IV. Quiet Areas	
All Cases ..	2,926	962	2,001	115	6,004
No. Infected by Prostitutes ..	1,332	69	462	30	1,893
Infected by Prostitutes (per cent.)	45	7	23	26	31

The total of the four groups in Table I also shows that 31 per cent. of 6,004 male cases of gonorrhoea were infected by prostitutes. This figure may, however, be significantly weighted by seafarers infected by prostitutes abroad and may not reflect fairly the position in the ports in which these seafarers were treated. The B.C.C.G. Gonorrhoea Study was a retrospective analysis and details of the sources of infection had not been fully recorded in all cases.

PRESENT STUDY

A detailed study of the epidemiological and social factors relating to venereal diseases was planned in Manchester in 1954 primarily to collect information

on which to develop greater control activities. As full information as possible has been recorded for each person with recent sexual exposure attending the two largest V.D. Clinics in Manchester—St. Luke's Clinic and the V.D. Clinic of the Manchester Royal Infirmary. This information, in addition to being entered on the patient's records, has been transcribed to a special register to facilitate subsequent analysis. The data collected during the year ending June 30, 1956, have been studied, and information regarding the role of the prostitute in the spread of venereal disease in Manchester is presented below.

Syphilitic infections have been excluded from the study as the number is so small. However, minor conditions requiring treatment and those cases seeking reassurance after exposure to infection, which together form a not unimportant part of the work of the V.D. clinics, have been included with gonorrhoea and non-gonococcal urethritis. The data have been analysed quarterly for St. Luke's Clinic (Table II, overleaf) Manchester Royal Infirmary Clinic (Table III, overleaf), and for Manchester for the year ending June 30, 1956 (Table IV, overleaf).

RESULTS

The experience in the two clinics is essentially uniform and there is little difference between the last three quarters of each year. The third quarter of each year (July to September inclusive) is usually the busiest quarter, and the figures for this quarter in 1955 are not exceptional. Of a total of 3,325 persons, 876 (26 per cent.) had consorted with a prostitute. As some men hesitate to admit to association with a prostitute, this percentage is regarded as a minimum figure. The term "prostitute" is used in its narrowest sense and was recorded only in cases in which money was stated to have been given to the woman. (It is interesting to note that the Special Committee of the British Medical Association, responsible for the B.M.A. Memorandum of Evidence (1955) for the Departmental Committee recently set up by the Government, defines female prostitution as "habitually consorting with a variety of men for sexual purposes and for gain in money or kind". I agree with this definition and believe that promiscuous women accepting kind rather than

* Received for publication July 27, 1956.

TABLE II
ST. LUKE'S CLINIC, MANCHESTER

Type of Infection	3rd Quarter, 1955			4th Quarter, 1955			1st Quarter, 1956			2nd Quarter, 1956		
	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)
Gonorrhoea	311	108	34	187	57	30	171	34	20	249	83	33
Non-Gonococcal Urethritis	139	57	41	97	16	16	107	15	14	135	22	16
Other conditions requiring treatment*	127	41	33	126	31	25	113	30	26	126	28	22
Other cases <i>not</i> requiring treatment	171	60	35	157	42	26	152	40	26	185	38	20
Totals	748	266	35	577	146	25	543	119	22	695	171	25

* All syphilis cases excluded.

TABLE III
MANCHESTER ROYAL INFIRMARY CLINIC

Type of Infection	3rd Quarter, 1955			4th Quarter, 1955			1st Quarter, 1956			2nd Quarter, 1956		
	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)
Gonorrhoea	102	31	30	90	20	22	74	20	27	73	23	31
Non-Gonococcal Urethritis	26	8	30	19	8	42	26	3	13	43	8	19
Other conditions requiring treatment*	17	4	23	22	3	14	22	3	14	15	4	26
Other cases <i>not</i> requiring treatment	57	9	16	71	13	18	58	8	14	57	9	16
Totals	202	52	25	202	44	22	180	34	19	188	44	23

* All syphilis cases excluded.

TABLE IV
MANCHESTER FOR THE YEAR ENDING JUNE 30, 1956

Type of Infection	St. Luke's Clinic			Royal Infirmary Clinic			Totals		
	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)	Total Cases	Exposed with Prostitute	Exposed with Prostitute (per cent.)
Gonorrhoea	918	282	30	339	94	28	1,257	376	30
Non-Gonococcal Urethritis ..	478	110	23	114	27	24	592	137	23
Other conditions requiring treatment*	492	130	28	76	14	18	568	144	24
Other cases <i>not</i> requiring treatment	665	180	27	243	39	16	908	219	24
Totals	2,553	702	27	772	174	23	3,325	876	26

* All syphilis cases excluded.

money are prostitutes in effect although not in law; in relation to the spread of venereal disease they play an important part which will be discussed in a later communication.) In many cases the actual sum was noted: it varied from 2s. to £6; exposures outside cost up to £2 while exposures inside in a hotel bedroom, the woman's apartment, or a rented

room varied between £2 and £6.

The 876 exposures to risk of infection from prostitutes resulted in 376 infections with gonorrhoea, 137 infections with non-gonococcal urethritis, and 144 minor conditions requiring some treatment. The remaining 219 cases attended for exclusion of infection and reassurance (Table V, opposite).

TABLE V
RESULTS OF EXPOSURE WITH PROSTITUTES IN 876 CASES

Clinic	Period	Number of Exposures	Gonorrhoea	Non-Gonococcal Urethritis	Other Conditions Requiring Treatment	Other Cases <i>not</i> Requiring Treatment
St. Luke's Clinic	3rd Quarter, 1955	266	108	57	41	60
	4th Quarter, 1955	146	57	16	31	42
	1st Quarter, 1956	119	34	15	30	40
	2nd Quarter, 1956	171	83	22	28	38
	Totals	702	282	110	130	180
Manchester Royal Infirmary Clinic	3rd Quarter, 1955	52	31	8	4	9
	4th Quarter, 1955	44	20	8	3	13
	1st Quarter, 1956	34	20	3	3	8
	2nd Quarter, 1956	44	23	8	4	9
	Totals	174	94	27	14	39
Grand Totals		876	376	137	144	219

Further analysis with regard to place of infection has shown that 85 per cent. of these exposures with prostitutes took place in Manchester. One half of the remainder (7 per cent.) occurred in other English cities, chiefly London and Liverpool, and the remaining 8 per cent. in foreign cities.

DISCUSSION

The data presented above show that in Manchester a quarter of all male patients attending the V.D. clinics do so after exposure with a prostitute. Furthermore, 30 per cent. of male cases of gonorrhoea and almost a quarter of the cases of non-gonococcal urethritis are infected by prostitutes. Some prostitutes attend the V.D. clinics at irregular intervals on their own initiative but the number is small. It is usually impossible to bring the source of infection of a male case of gonorrhoea to treatment when he has been exposed with a prostitute as the economic basis of the association does not encourage acquaintance with those personal particulars which would assist in contact tracing. There is no doubt that the prostitute contributes substantially to the "reservoir" of untreated female gonorrhoea and that treatment of the prostitute would assist significantly in the control of gonorrhoea in Manchester. Police action against soliciting is handicapped by the unsatisfactory state of the law and, on conviction, the imposition of the maximum fine of forty shillings sets the woman free to recoup her fine at the expense of the health of the public. It would be logical and helpful if Magistrates were to recommend attendance at the V.D. clinics for examination and treatment, perhaps at special sessions of the clinic. This is currently done throughout Canada and in many

cities in the U.S.A., the examination room often being situated in the same building as the Court. It is sincerely hoped that the Report of the Departmental Committee, which is at present considering the law and practice relating to homosexuality and prostitution, will include recommendations on this aspect of the problem of prostitution, and that realistic legislation will permit police and Magistrates to discourage prostitution and assist in the control of venereal disease.

SUMMARY

Data from the special study, which is being made of the epidemiological and social aspects of venereal diseases, show that in Manchester 30 per cent. of male cases of gonorrhoea and the attendance of a quarter of all male patients at the major V.D. clinics result from exposure with prostitutes. Altogether 85 per cent. of these exposures with prostitutes took place in Manchester.

As it is usually impossible to effect the attendance of the prostitute at the V.D. clinic through the infected male client, it is urged that police action against prostitution should be encouraged by more realistic legislation, and that Magistrates should recommend the prostitute to report to the V.D. clinic for examination.

I wish to thank Mr. J. Payton and Mr. A. Tattersall of St. Luke's Clinic for their work in transcribing the information to the special register.

REFERENCES

- B.M.A. Memorandum (1955). *Brit. med. J.*, 2, Suppl., 169.
- British Clinical Cooperative Group Gonorrhoea Study (1956). *British Journal of Venereal Diseases*, 32, 21.
- British Social Biology Council (1955). "Women of the Streets: A Sociological Study of the Common Prostitute", p. 93. Ed. by C. H. Rolph. Secker and Warburg, London.
- Nicol, C. S. (1956). *British Journal of Venereal Diseases*, 32, 27.