Isolation of *Neisseria meningitidis* and *Neisseria catarrhalis* from the genitourinary tract and anal canal

CAROLINE BLACKWELL, HUGH YOUNG, AND SHEILA S. R. BAIN*

From the Department of Bacteriology, University of Edinburgh, and the University Department of Venereology,* Edinburgh Royal Infirmary

SUMMARY During a 12-month period 285 isolates of *Neisseria* species, other than *Neisseria gonorrhoeae*, were cultured from patients attending the Department of Venereology, Royal Infirmary, Edinburgh. There were eight patients in whom genitourinary or rectal isolates of *Neisseria meningitidis* or *Neisseria catarrhalis* were found. Differences between data from our series and from previous reports are discussed with particular regard to sites routinely cultured in female patients. We also report *in vitro* inhibition of *N. gonorrhoeae* by an isolate of *N. meningitidis* cultured from the cervix of a patient.

Introduction

Neisseria species other than *Neisseria gonorrhoeae* have been found to cause symptomatic infections of the genitourinary tract and the anal canal. A report of an increase in incidence of isolation as well as a brief review of the literature on *Neisseria meningitidis* infections of these sites has been recently published (Givan *et al.*, 1977). In the present study we isolated 285 non-gonococcal species of *Neisseria*. Among these there were four *N. meningitidis* isolates and five *Neisseria catarrhalis* isolates from genitourinary or rectal sites. Our findings for *N. meningitidis* differ from those previously reported—in particular, the number of male patients in the series from whom the organism was isolated, overt signs of infection, autoinoculation from the nasopharynx as a source of the organism. The possibility of inhibitory substances produced by *N. meningitidis* preventing infection by *N. gonorrhoeae* in one patient is discussed.

Materials and methods

PATIENTS

The patients in this study attended the Department of Venereology, Edinburgh Royal Infirmary between 1 July 1976 and 30 June 1977. During this

period there were approximately 2000 women and 2500 men.

In men, if a urethral discharge was present, or if there was presumptive evidence of contact with gonorrhoea, a Gram-stained film of urethral discharge was examined microscopically and material inoculated directly on to culture plates. In the case of homosexuals, material from the urethra, rectum and pharynx was cultured routinely, and if negative on the first occasion, repeated twice at weekly intervals.

In women, Gram-stained films of material from the urethra and cervix, and cultures from the urethra, cervix, rectum and, if indicated, from the pharynx were taken. Generally, if the first cultures were negative for *N. gonorrhoeae*, these were repeated twice at weekly intervals.

CULTURE AND IDENTIFICATION

During the first six months of the study cultures were made on Thayer-Martin medium containing the antibiotics vancomycin (4.0 µg/ml), colistin (6.0 µg/ml), and nystatin (10.0 µg/ml). During 1977, cultures were made on modified New York City Medium (Young, 1978) containing the antibiotics lincomycin (1.0 µg/ml), colistin (6.0 µg/ml), amphotericin B (1.0 µg/ml), and trimethoprim lactate (6.5 µg/ml).

Cultures were incubated at 36°C in a carbon dioxide enriched (10%) atmosphere. After incubation for 24 hours, plates were examined and any suspect colonies tested by the oxidase reaction and Gram's stain. Oxidase positive Gram-negative diplococci

Address for reprints: C. Blackwell, Department of Bacteriology, University of Edinburgh, Medical School, Teviot Place, Edinburgh EH8 9AG

Received for publication 22 July 1977

were identified by the rapid carbohydrate utilisation test and direct immunofluorescence using Difco fluorescein-labelled anti-gonococcal conjugate as described by Young *et al.* (1976). Negative cultures were re-examined after 48 hours of incubation.

Demonstration of the inhibitory substance from *N. meningitidis* was by means of the soft agar method for bacteriocin production (Fredericq, 1957).

Results

During the 12-month period from 1 July 1976 to 30 June 1977 approximately 4500 patients attended the Department of Venereology, Royal Infirmary, Edinburgh. There were 350 specimens of *N. gonorrhoeae* isolated from the 2000 female patients and 480 specimens isolated from the 2500 male patients.

The *Neisseria* isolates, other than *N. gonorrhoeae*, from these patients are summarised in Table 1.

Table 1 Isolation of *Neisseria* species other than *N. gonorrhoeae* over a 12-month period

Neisseria	Total no. of isolates	Site of isolation		
		Throat	Genito-urinary tract	Anal canal
<i>N. meningitidis</i>	250	246	2	2
<i>N. lactamica</i>	23	23	0	0
<i>N. catarrhalis</i>	7	2	1	4
<i>N. flava</i>	3	3	0	0
<i>N. perflava</i>	2	2	0	0

N. meningitidis was the species most often isolated (250); among these there were two from the anal canal and two from the genitourinary tract. We found *N. lactamica* (23), *N. flava* (3), and *N. perflava* (2) only in throat cultures. Of the seven *N. catarrhalis*, two were from throat cultures, one was from the genitourinary tract, and four from the anal canal.

There were four patients from whom *N. meningi-*

tidis was isolated. The information available for each patient is summarised in Table 2. The only additional systemic complaints from Patient 1 were a loss in weight of 7 lb (3.175 kg), dysuria, and frequency of micturition. *N. gonorrhoeae* were not isolated at this time nor on the three subsequent sets of cultures. She had had a regular consort for the past two years and the only previous history of a sexually transmitted disease (STD) was warts in 1975 and genital candidosis in 1974. A second rectal isolate was from Patient 2 who had attended the clinic four months previously, at which time *N. meningitidis* was isolated from the throat. The only systemic complaint was eczema on face, trunk, and limbs and there was no previous history of STD. Patient 3, a bisexual male with no previous history of STD, had attended the clinic two months earlier. At that time *N. meningitidis* was isolated from the throat but no *Neisseria* had been found in cultures of rectum or urethra.

The only patient in the series from which we isolated *N. meningitidis* and, later, *N. gonorrhoeae* was Patient 4. She was seen in the clinic as a gonorrhoea contact; her husband had become infected from an extramarital consort. Microscopical examination of material from the cervix revealed large numbers of polymorphonuclear leucocytes with intracellular Gram-negative diplococci. She was treated with co-trimoxazole as there was a possibility of penicillin allergy. The culture of cervical material grew only *N. meningitidis* and no gonococci were isolated from any other site. A month later gonococci were cultured from the cervix and she was treated with another course of co-trimoxazole. Disc sensitivity tests revealed both the previously isolated meningococcus and the gonococcus to be sensitive to trimethoprim-sulphamethoxazole (disc containing 23.5 µg sulphamethoxazole, 1.5 µg trimethoprim). At the time it was serotyped, the meningococcus was

Table 2 Isolation of *N. meningitidis*

Patient	Sex	Site of isolation	Throat culture	Serotype	Reason for attending STD clinic	Diagnosis, treatment, and follow-up
1	F	Rectum	—	B	Warts	Cautery and liquid nitrogen for warts; 3 subsequent sets of cultures negative
2	F	Rectum	Not done	B	Frequency of micturition; warts	Podophyllin and cautery for warts; no follow-up cultures subsequent to isolation of <i>N. meningitidis</i> from anal canal
3	M	Urethra	Not done	NT	Syphilis contact	No treatment; no follow-up cultures
4	F	Cervix	—	NT	Gonorrhoea contact	Asymptomatic gonorrhoea (only <i>N. meningitidis</i> cultured); co-trimoxazole 5 days; 2 subsequent sets of cultures negative; <i>N. gonorrhoeae</i> cultured 3 times during next 3 months

—=no *Neisseria* isolated

NT=non sero-typable *N. meningitidis*

further tested and found to grow on 0.1 µg/ml sulfadiazine but not on 1 µg/ml. She denied any sexual contact with her husband during this period. The soft agar method for demonstrating sensitivity to bacteriocins showed the meningococcus elaborated a substance which greatly inhibited the gonococci from Patient 4 and several other *N. gonorrhoeae* isolates from different sources. It was later discovered that the gonococcal infection was probably not a treatment failure or inhibition by the meningococcal substance but infection from an extramarital contact. There was no systemic upset in Patient 4, but her extramarital contact and his wife developed systemic infections and arthritis.

There were four female patients from whom *N. catarrhalis* was isolated (Table 3). None of these had any systemic upset and no gonococci were cultured at follow-up visits.

Discussion

Givan *et al.* (1977) have reviewed case reports of isolation of *N. meningitidis* from the genitourinary tract and anal canal. Our data from the past 12 months provide a somewhat different view. These differences may be owing to routine culturing of urethra, cervix, and rectum of all female patients. Previous reports mention only cervical cultures of women, with additional blood or joint cultures from two systemic infections (Keys *et al.*, 1971; Morgan *et al.*, 1976). Other series contain a predominance of men, 58 of the 84 patients in the literature; in our study three of the four patients from whom we isolated *N. meningitidis* and all of the patients from whom we isolated *N. catarrhalis* were women.

There were no overt symptoms of these meningococci causing infection in any of the patients, again in contrast to the findings of others. The main reason for attending the Department of Venereology was genital warts (four out of eight patients). Three were either syphilis or gonorrhoea contacts, and the fourth was referred for follow-up of a gonococcal

infection. There was no evidence of rectal infections with *N. meningitidis* associated with systemic infection. Currently there is no evidence for treating rectal carriers, but follow-up cultures to monitor any spread to the urethra or the cervix may be of value.

It has been suggested (Givan *et al.*, 1977) that there may be autoinoculation of these areas from the nasopharynx. In their series all six of the rectal carriers from whom there were throat cultures had meningococcus in the pharynx. In our series, none of the patients from whom throat cultures were taken was carrying *N. meningitidis* in the nasopharynx. It is also interesting to note that meningococci were not isolated from either contact of Patient 4.

The significance of the inhibitory effect of *N. meningitidis* from Patient 4 on *N. gonorrhoeae* is difficult to assess. A similar instance of apparent protection has been reported for a male patient with asymptomatic meningococcal urethritis (Volk and Kraus, 1973). The role of this inhibitory effect in preventing infection from her initial exposure is not clear. The meningococcus would probably have been eliminated by the first co-trimoxazole treatment before her second exposure and subsequent infection with *N. gonorrhoeae*.

These observations underline the necessity of correct and full identification of all neisserial isolates, even when a selective medium is used. These non-gonococcal *Neisseria* represent approximately 2% of the anogenital isolates in women and 0.2% of those from men.

This work is part of a project supported by a grant from the Scottish Home and Health Department, Medical Research Committee.

Serotyping of meningococci and sulphadiazine sensitivity testing was kindly carried out by Dr R. J. Fallon, Department of Laboratory Medicine, Ruchill Hospital, Glasgow.

Table 3 Isolation of *N. catarrhalis*

Patient	Sex	Site of isolation	Throat culture	Reason for attending STD clinic	Diagnosis, treatment, and follow-up
5	F	Rectum	—	Gonorrhoea contact	No treatment; 3 subsequent sets of cultures negative
6	F	Rectum	Not done	Warts	Podophyllin for warts; 3 subsequent sets of cultures negative
7	F	Rectum	Not done	Follow-up treatment for gonorrhoea	Non-gonococcal vaginal discharge; nystatin for discharge
8	F	Urethra Rectum	Not done	Warts	Podophyllin for warts; 4 subsequent sets of cultures negative

— = no *Neisseria* isolated

References

- Fredericq, P. (1957). Colicins. *Annual Review of Microbiology*, 11, 7–22.
- Givan, K. F., Thomas, B. W., and Johnston, A. G. (1977). Isolation of *Neisseria meningitidis* from the urethra, cervix, and anal canal: Further observations. *British Journal of Venereal Diseases*, 53, 109–112.
- Keys, T. F., Hecht, R. H., and Chow, A. W. (1971). Endocervical *Neisseria meningitidis* with meningococemia. *New England Journal of Medicine*, 285, 505–506.
- Morgan, J. R., Fedder, J., and Smith, J. A. (1976). Postpartum infection with meningococemia. *Canadian Medical Association Journal*, 114, 294.
- Volk, J., and Kraus, S. J. (1973). Asymptomatic meningococcal urethritis. Possible protective value against gonococcal infection by bacteriocin production. *British Journal of Venereal Diseases*, 49, 511–512.
- Young, H. (1978). Cultural diagnosis of gonorrhoea with modified New York City (MNYC) medium. *British Journal of Venereal Diseases*, 54, 36–40.
- Young, H., Paterson, I. C., and McDonald, D. R. (1976). Rapid carbohydrate utilisation test for the identification of *Neisseria gonorrhoeae*. *British Journal of Venereal Diseases*, 52, 172–175.