



One

Two



Free!

Canesten clears vaginal candidiasis in three days: and speed is essential since almost 50% of women may fail to complete the 14 days demanded by conventional therapy.¹ In a recent study, 103 patients were treated for three days with Canesten. Four weeks later, 89% were clinically and mycologically clear of candidiasis.²

Three-day Canesten is economical, has a low relapse rate, rarely causes irritation and does not stain.

A Canesten Duopak is available containing vaginal tablets for the patient and cream to treat her vulvitis or her partner to help prevent reinfection.

Canesten[®] clears vaginal clotrimazole candidiasis in three days

1. *Current Medical Research and Opinion*, 3, 83-88, 1975. 2. *British Journal of Venereal Diseases*, 53, 126-128, 1977. Canesten is available as vaginal tablets, each containing 0.1g clotrimazole, in packs of six with applicator. Canesten Duopak contains six vaginal tablets with applicator plus 20g cream (1% clotrimazole). Dosage - vaginal tablets: Two vaginal tablets should be inserted daily for three consecutive days. Alternatively, one vaginal tablet daily for six days. Duopak: Vaginal tablets as above. The cream should be applied night and morning to the vulva and surrounding area and/or to the sexual partner's penis to prevent re-infection. Side effects: Rarely patients may experience transient local mild burning or irritation. N.H.S. costs: 6 vaginal tablets with applicator £1.92; Duopak £3.13. Product Licence numbers: Vaginal tablets: PL0010/0015; Duopak: PL0010/0016 & PL0010/0015. Bayer UK Limited, Pharmaceutical Division, Haywards Heath, West Sussex RH16 1TP. Telephone: Haywards Heath (0444) 57911. © Registered Trade Mark of Bayer Germany.

