

## Correspondence

TO THE EDITOR, *British Journal of Venereal Diseases*

### Lack of effect of bicarbonate on the survival of *Treponema pallidum* (Nichols) in vitro

Sir,  
No published studies have reported the effect of bicarbonate ( $\text{HCO}_3^-$ ) on the survival of *Treponema pallidum* in vitro.

Carbon dioxide ( $\text{CO}_2$ ) has been shown to be beneficial to *Neisseria gonorrhoeae*, *N meningitidis*, some staphylococci,<sup>1</sup> and other bacteria.<sup>2,3</sup>  $\text{CO}_2$  serves as a carboxylating agent<sup>4</sup> and as an essential precursor for the synthesis of certain amino acids.<sup>5</sup> It is also required for fatty acid synthesis,<sup>1,6</sup> biosynthesis of purine ribonucleotides, which are precursors of DNA and RNA and are an integral part of many coenzymes.<sup>6</sup>

Different concentrations of sodium bicarbonate ( $\text{NaHCO}_3$ ) were used in both a cell-free and cell-culture system to determine whether bicarbonate had any beneficial effect on the survival of *T pallidum* in vitro.

The medium in the cell-free system<sup>7</sup> was slightly modified by replacing the sodium thioglycollate with 2 mmol/l dithiothreitol (DTT). In the cell-culture system, Eagle's minimal essential medium with 10% heat-inactivated fetal calf serum, 0.5 mmol/l DTT, and 20 mmol/l N-2-hydroxyethyl-piperazine-N-2-ethane-sulphonic-acid (HEPES) was used.

*T pallidum* was harvested from infected rabbit testes.<sup>7</sup> Anaerobic incubation has been described.<sup>7</sup> For microaerophilic incubation, tubes with cottonwool plugs were equilibrated with 3%  $\text{O}_2$ <sup>7</sup> for two days before inoculation with treponemes. Treponemes were then added to an approximate final density of  $5 \times 10^6$ /ml, followed by  $\text{HCO}_3^-$  to give a final concentration of from  $10^{-5}$  to  $10^{-2}$  mol/l. The cottonwool plugs were immediately changed to rubber stoppers to prevent escape of  $\text{CO}_2$ . The pH remained at  $7.3 \pm 0.2$  in the cell-free system. The incubation temperature was 34°C.

The isolation and subculture of the rabbit testicular cell lines have been described.<sup>8</sup>

Viability of *T pallidum* in vitro was determined by the percentage motility under darkfield microscopy and their virulence in rabbits.<sup>7</sup> In the cell-culture

system, attached treponemes were separated from tissue cells<sup>8</sup> before virulence was tested.

Ten experiments were performed in a cell-free medium under anaerobic conditions; a representative experiment is shown in the figure. There was no enhancement by  $\text{HCO}_3^-$  ( $10^{-2}$  to  $10^{-5}$  mol/l) on the retention of motility of *T pallidum* (figure; a); the percentage motility in the presence of  $10^{-3}$  and  $10^{-2}$  mol/l  $\text{HCO}_3^-$  was consistently lower than the control. Similarly,  $\text{HCO}_3^-$  ( $10^{-2}$  to  $10^{-5}$  mol/l) had no beneficial effect on the retention of virulence of *T pallidum* in vitro (figure; b).

Bicarbonate ( $10^{-5}$  to  $10^{-2}$  mol/l) showed no enhancement of treponemal survival in two experiments under microaerophilic conditions with 3%  $\text{O}_2$ . The percentage motility (figure; c) and latent periods (figure; d) were virtually the same at all concentrations tested and were not appreciably different from the control tubes lacking bicarbonate.

In three experiments in cell-cultures, the pH of the medium dropped to 6.8 at day 4 in the control tubes while remaining at 7.2 in the presence of  $\text{HCO}_3^-/\text{CO}_2$ . When the pH of the medium was regularly adjusted, however,  $\text{HCO}_3^-$  ( $10^{-4}$  to  $10^{-2}$  mol/l) had no enhancing effect on the retention of treponemal motility (figure, e) or virulence (figure, f) in the presence of tissue culture cells. Thus, in the media tested,  $\text{HCO}_3^-/\text{CO}_2$  may serve mainly as a buffering system rather than as a reactant in any essential metabolic process.

Possible reasons why  $\text{HCO}_3^-/\text{CO}_2$  had no enhancing effect on treponemal survival in vitro may be: (a)  $\text{CO}_2$  requirements were satisfied by its endogenous metabolism, such as degradation of pyruvate and glucose.<sup>9,10</sup> (b) *T pallidum* under conditions which do not allow multiplication may have no capacity to use  $\text{CO}_2$  even if it normally utilises it. (c) Even if *T pallidum* had the capacity to incorporate  $\text{CO}_2$ , it may be unnecessary. Since the medium was extremely rich, the presence of these nutrients may have shut down certain biosynthetic pathways. The incorporation of  $\text{NaH}^{14}\text{CO}_3$  was insignificant when compared to total glucose incorporated.<sup>10</sup>

(d)  $\text{CO}_2$  would not be used for fatty acid synthesis since *T pallidum*, like other treponemes, cannot synthesise fatty acids.<sup>11</sup> (e)  $\text{HCO}_3^-/\text{CO}_2$  may serve as a buffer in a

closed system, but with HEPES (20 mmol/l) serving this purpose  $\text{HCO}_3^-/\text{CO}_2$  was unnecessary in both the cell-free and cell-culture systems.

We gratefully acknowledge the excellent technical assistance provided by Mrs L Drummond and thank Professor S Fine and Dr Ben Adler for constructive critical evaluation. This work was supported by grants from Australian National Health and Medical Research Council (NH&MRC); the Utah Foundation, the Heizer Program, and Monash University.

Yours faithfully,  
G H W Wong  
B M Steiner  
S R Graves

Department of Microbiology,  
Monash University,  
Victoria,  
Australia.

### References

- Davis BD, Dulbecco R, Eisen HN, Ginsberg HS. *Microbiology*. Harper International Edition (3rd edition). Maryland, USA: Harper & Row, 1980.
- Mickelson MN. Chemically defined medium for growth of *Streptococcus pyogenes*. *J Bacteriol* 1964; **88**: 158-64.
- Lachina RVF, Hartman PA. Carbon dioxide fixation and the synthesis of aspartic acid by *Streptococcus faecium* var *durans*. *Biochem Biophys Res Commun* 1968; **32**: 691-5.
- Lyman CM, Mosely O, Wood S, Butter B, Hale F. Some chemical factors which influence amino acid requirements of lactic acid bacteria. *J Biol Chem* 1947; **167**: 177-87.
- Hancock R, McManus F. Carbon dioxide fixation in the synthesis of aspartic acid by a strain of *Staphylococcus aureus*. *Biochim Biophys Acta* 1960; **42**: 152-4.
- Lehninger AL. *Biochemistry*, 2nd ed. New York: Worth, 1976.
- Steiner B, Mclean I, Graves S. Repox potential and survival of virulent *Treponema pallidum* under microaerophilic conditions. *Br J Vener Dis* 1981; **57**: 295-301.
- Fitzgerald TJ, Miller JN, Sykes SA. *Treponema pallidum* (Nichols strain) in tissue cultures: cellular attachment, entry, and survival. *Infect Immun* 1979; **11**: 1133-40.
- Barbieri JT, Cox CD. Pyruvate oxidation by *Treponema pallidum*. *Infect Immun* 1979; **25**: 157-63.
- Barbieri JT, Cox CD. Glucose incorporation by *Treponema pallidum*. *Infect Immun* 1979; **24**: 291-3.
- Smbirt RM. Spirochaetales, a review. *Crit Rev Microbiol* 1973; **2**: 491-552.

Br J Vener Dis: first published online 1 April 1982. Downloaded from <http://journals.cambridge.org/hyg>. Archived from <http://www.cambridge.org/core>. Created by copyright.