Analysis of immune responses in genital tracts of mice immunised with purified ribosomal fractions of *Neisseria gonorrhoeae*

EIJI KITA AND SHUZO KASHIBA

From the Department of Bacteriology, Nara Medical College, Kashihara City, Japan

SUMMARY Immunisation of ddY mice with the purified ribosomal fraction of *Neisseria gonorrhoeae* was found to protect against intravaginal challenge with homologous organisms. This protection correlated with the presence of bactericidal antibody to purified ribosomal fraction in serum as well as in vaginal secretions. Analysis of the vaginal fluids from control mice and those immunised with purified ribosomal fraction showed that the enhanced elimination of gonococci in immune mice might be because of an early response of leucocytes generated by the reaction mediated by antibody and complement. Absorption studies showed that there was at least one major protective antigen in purified ribosomal fraction, other than cell surface substances such as lipopolysaccharide, outer membrane proteins, and pili. Bactericidal assays mediated by antibody and complement showed that matched samples of serum and vaginal fluid from immune mice had comparable gonococcal activity, which was augmented by the effect of progesterone. Although delayed hypersensitivity was produced in immune mice that were resistant to *N. gonorrhoeae*, the exact role of cellular immunity could not be clarified in this study. These results suggest that antibody to purified ribosomal fraction plays a major part in protection against gonococcal infection in the genital tract, and that such protection may entail both cellular immunity and hormonal changes.

Introduction

The development of practical models of infection with *Neisseria gonorrhoeae* in laboratory animals^{1, 2} and in chimpanzees³ has appreciably advanced our understanding of the immunobiology of this organism. We have reported a mouse model for the study of gonococcal genital infection,⁴ and showed that female ddY mice were sensitive to genital infection with gonococci during the preovulatory period of the oestrous cycle. In recent years, the subcutaneous chamber model in guinea pigs has been the most widely used to monitor the immune response to gonococcal infection.^{5, 6} Although the subcutaneous chambers are sites at which humoral immunity to experimental infection can be analysed, the site of infection is very different from that of natural infection in man. One of the important factors affecting the susceptibility to enhanced infection in women is the effect of reproductive hormones in the

genital tract.⁷ Our previous work using the mouse uterus as the site of infection showed the preovulatory stage to be the most suitable for intravaginal inoculation of gonococci, because at this stage the effects of polymorphonuclear leucocytes and progesterone are absent.

The present study was undertaken to investigate the genital immune responses in mice immunised with ribosomal vaccine.

Materials and methods

MICE

Female ddY mice 5 to 6 weeks old (Shizuoka Agricultural Cooperative Association for Laboratory Animals, Shizuoka, Japan) were used for all experiments. They were fed a commercial laboratory animal diet and water ad libitum. The stages of their oestrous cycles were determined by conventional vaginal smear techniques^{8, 9} as reported before.⁴

BACTERIA

The PH2 (P + + Tr) strain of *N. gonorrhoeae* used in this study was a penicillin sensitive isolate from the human genital tract. Characteristics of strain PH2

Address for reprints: Dr E Kita, Department of Bacteriology, Nara Medical College, 840 Shijyocho, Kashihara City, Nara 634, Japan

Accepted for publication 13 December 1983

have been described previously.⁴ Gonococci were passed serially through mice to maintain their virulence. They were also cultured on GC agar base supplemented with isovitalax (BBL, Microbiology Systems, Cokeysville, Maryland, USA) for about 20 hours at 37°C in a moist candle jar. Identification of cultures of *N gonorrhoeae* by colony morphology, oxidase reaction, and Gram stain was confirmed by sugar fermentation tests. Gonococci from plates with more than 95% type 2 colonies, as determined by the technique and criteria of Kellogg *et al.*,¹⁰ were used for challenge. Organisms were preserved by the gelatin disc method as described by Yamai *et al.*¹¹

We grew gonococci for experiments in liquid medium containing 2% (w/v) tryptose (Difco Laboratories, Detroit, Michigan, USA), 0.4% dipotassium phosphate, 0.1% potassium biphosphate, 0.5% sodium chloride, and 0.1% soluble starch (Difco). After autoclaving and immediately before use we added 1% (w/v, final concentration) glucose, 0.04% sodium bicarbonate, and 0.01% cocarboxylase (thiamine pyrophosphate chloride, Sigma Chemical Co, St Louis, Missouri, USA) and sterilised the medium by filtration through a type HA 0.45 µm grid membrane filter (Millipore, Bedford, Massachusetts, USA). Broth cultures were grown at 37°C with continuous revolution after the culture bottle had been tightly capped with a rubber stopper.

PREPARATION OF THE PURIFIED RIBOSOMAL VACCINE

We prepared the ribosomal vaccine by the method of Kita and Kashiba,¹² and purified it by affinity chromatography using Sepharose conjugates of antibody to outer membrane complex as previously described.¹³ Briefly, the antigens that cross reacted with antibodies to outer membrane complex were removed from the ribosomal fraction by passing it over a 15 ml column containing conjugated Sepharose 4B. The fraction was then pooled, dialysed, lyophilised, and stored at -70°C until used. The ribosomal fraction obtained by affinity chromatography was designated the purified ribosomal fraction. Antiserum to purified ribosomal fraction prepared in rabbits did not react with envelope components of *N gonorrhoeae* strain PH2 in the immunodiffusion test.

The ratio of ribonucleic acid (RNA) to protein was 2:14, and endotoxin was not detected by a limulus lysate coagulation test. The 2-keto-3-deoxyoctonate was not detectable in the purified ribosomal fraction by the method of Karkhians *et al.*¹⁴

GONOCOCCAL INFECTION IN THE GENITAL TRACT

Gonococci grown in the liquid medium were suspended at a concentration of 10⁸ colony forming

units (cfu)/ml in trypticase soy broth prewarmed at 37°C, and kept at 37°C until inoculated. The gonococcal suspension (0.01 ml) was inoculated into the vagina with a sterile tip on a micropipette (Pipetman) as previously described.⁴ Mice were inoculated in the late proestrous or early oestrous stages, and one week after inoculation were killed for cultures of uterine specimens. The uterine bodies were removed, resected, and homogenised in 0.1 ml sterile trypticase soy broth. The homogenate was centrifuged at 100 × g for five minutes to sediment cell debris. The supernatant (0.05 ml) was cultured on GC agar plates with and without 1% VCN (vancomycin-colistin-nystatin). After overnight incubation at 37°C in a candle jar, the colonies formed on plates were identified as *N gonorrhoeae* by Gram stain and oxidase test, and their identity was confirmed by a sugar fermentation test.

PREPARATION OF ANTIGENS USED FOR PHA

Outer membrane was isolated by the method described by Wolf-Watz *et al.*¹⁵ Crude outer membrane fraction obtained by centrifuging at 280 000 × g for 70 hours using sucrose density gradient (65%-40%) was pooled after centrifuging several times, and was lyophilised after dialysis. Gonococcal pilus preparation was isolated by the standard purification method described by Brinton *et al.*¹⁶ Lipopolysaccharide was extracted from whole cells by the hot phenol method of Westphal and Jann,¹⁷ digested with pronase (50 mg/ml protease from *Streptomyces griseus*, Sigma) at 40°C for four hours and heated at 100°C for five minutes to stop the reaction. The reacted lipopolysaccharide was centrifuged at 100 000 × g for 30 minutes, washed several times, and finally suspended in saline.

IMMUNISATION WITH RIBOSOMAL VACCINE

Mice were immunised subcutaneously with an appropriate amount of purified ribosomal fraction.

SERUM ANTIBODY TITRES

Specimens of serum were obtained from mice on days 0, 7, 14, and 28 after the final immunisation. Serum collected from each mouse was frozen and stored at -70°C until tested. Each serum sample was examined for the presence of antibodies against gonococcal antigens, and for the bactericidal activity against the challenge organism. Antibody titres were measured by the passive haemagglutination test as previously described,¹³ using sheep red blood cells coated with purified ribosomal fraction, pili, outer membrane, and lipopolysaccharide as passive haemagglutination test antigens.

Bactericidal activity was measured by the method of Wannemuehler *et al.*⁶ with a slight modification. *N gonorrhoeae* PH2 type 2 was grown in the liquid

medium, and cells were then washed with warm (37°C) phosphate buffered saline (PBS) containing 0.5% glucose, 0.01 mol/l magnesium chloride, and 0.01% fetal bovine serum. The cells were diluted to 10^6 cfu/ml in RPMI 1640 (Gibco Laboratories, Grand Island, New York, USA). After it had been inactivated at 56°C for 30 minutes serum was serially (twofold) diluted in RPMI 1640 by using microtitre U plates, and the same volume of the cell suspension (50 µl) was added to each well. A 50 µl volume of a 1/10 dilution of normal guinea pig serum used as a source of complement was added to the reaction mixture, which was then incubated at 37°C in a carbon dioxide chamber for 90 minutes while being shaken. Samples (25 µl each) were plated in triplicate on agar plates. All plates were incubated at 37°C in a candle jar for 24 hours. Bactericidal activity was expressed as the maximum dilution yielding a 90% reduction in cfu when compared with corresponding controls, which lacked complement.

VAGINAL POLYMORPHONUCLEAR LEUCOCYTE RESPONSE DURING INFECTION

Total leucocyte counts in vaginal washing fluid were made with a haemocytometer at 12 hour intervals after challenge. The vaginal wash fluid was obtained from immune mice and pooled for measurement of bactericidal activity. The pooled fluid was centrifuged at $400 \times g$ for 10 minutes, concentrated by ultrafiltration, and sterilised by a membrane filter. Bactericidal assay of the vaginal wash fluid was performed in the same way as with the serum.

DELAYED FOOTPAD TEST

Mice immunised with ribosomal vaccine were tested by the footpad test for delayed hypersensitivity. Ribosomal proteins extracted by the method of Fogel and Sypherd¹⁹ were used as the eliciting antigen, and a 25 µl volume of solution containing protein 100 mg/l was injected intradermally into the right hind footpad. Footpad thickness was measured 48 hours later, and any increase in thickness was corrected for by deducting the increase in thickness of the control left hind footpad, which had been injected with saline.

ABSORPTION STUDY

Samples of serum and vaginal fluid obtained from immune mice were absorbed by incubation with sheep red blood cells coated with antigen suspended in two volumes of serum or vaginal fluid for one hour at 37°C with slow shaking, and were then kept at 4°C overnight. The absorbed samples were centrifuged at $1200 \times g$ for 15 minutes, and the absorption procedure was repeated twice.

Results

PROTECTION AGAINST GONOCOCCAL UTERINE INFECTION BY PURIFIED RIBOSOMAL FRACTION
Our previous work showed that an inoculum size of 10^6 gonococci was optimal for the production of uterine endometritis in mice, and that the infectivity, as measured by the percentage of culture positive uterine specimens one week after challenge, was almost 100%, regardless of the stage of the oestrous cycle. In this study the minimum inoculum dose required to establish uterine endometritis in mice was found to be 10^6 cfu in the late proestrous stage (data not shown).

Mice were immunised subcutaneously with different amounts of purified ribosomal fraction and intravaginally challenged 10 days after immunisation with 10^6 cfu of strain PH2. Table I shows that the minimum effective dose required to achieve 90% protection was 40 µg per mouse.

TABLE I Protection of purified ribosomal fraction against uterine infection

Immunising dose* (µg protein)	Protection (%)† on day 7
5	25 (p>0.1)
10	50 (p>0.01)
20	65 (p<0.01)
40	90 (p<0.001)
60	100 (p<0.001)
Saline (control)	5

*Each group consisted of 20 mice immunised subcutaneously with the indicated dose. Mice in the late proestrous stage were challenged intravaginally with 10^6 cfu of strain PH2 10 days after immunisation. Data were obtained from two separate experiments.

†No. of mice with uterine specimens yielding negative cultures, expressed as percentage of mice studied. P values with respect to controls (saline) were calculated by Fisher's exact test.

HUMORAL IMMUNE RESPONSE

Subcutaneous immunisation with 20 µg of purified ribosomal fraction elicited a high antibody response as early as 10 days after a single injection of the fraction. Activity of antibody to purified ribosomal fraction reached a maximum two to three weeks after immunisation. Antibodies against outer membrane and pilous antigens could not be measured in serum samples of mice immunised with purified ribosomal fraction. Samples of serum and vaginal fluid obtained from immune mice gave a single precipitin band which formed a line of identity when tested by immunodiffusion against purified ribosomal fraction. No visible line was formed against lipopolysaccharide, pilous, or outer membrane preparations.

BACTERICIDAL ACTIVITY OF IMMUNE SERUM AND VAGINAL FLUID

Table II shows bactericidal activity of samples of serum and vaginal fluid from mice immunised with

TABLE II *Bactericidal activity of vaginal fluid and serum from mice immunised with gonococcal purified ribosomal fraction*

Immunogen*	Oestrous stage when samples taken	Bactericidal activity†		Protection (%)‡ on day 7
		Serum	Vaginal fluid	
Purified ribosomal fraction	Late proestrous	1/8	1/16	60 (p<0.001)
	Early oestrous	1/8	1/16	65 (p<0.001)
	Early metoestrous	1/16	1/32	90 (p<0.001)
	Dioestrous	1/32	1/64	100 (p<0.001)
Saline (control)	Late proestrous	1/2	1/2	0
	Early oestrous	1/2	1/2	0
	Early metoestrous	1/4	1/4	5
	Dioestrous	1/4	1/8	5

*Groups of 40 mice were immunised subcutaneously with 20 µg purified ribosomal fraction and serum and vaginal fluid collected from 20. The other 20 were challenged intravaginally with 10⁶ cfu of strain PH2 and protection was calculated by uterine culture on day 7. Data were obtained from two separate experiments.

†Bactericidal activity expressed as the maximum dilution yielding a 90% reduction in cfu when compared with corresponding controls lacking complement.

‡No of mice with uterine specimens yielding negative cultures, expressed as percentage of mice studied. P values with respect to the corresponding controls (saline) were calculated by Fisher's exact test.

purified ribosomal fraction. Serum samples from immunised mice were bactericidal at titres ranging from 1/8 to 1/32, while those from control mice injected with saline had titres of less than 1/4. The degree of bactericidal activity of immune serum correlated with the stage of oestrous. Serum from mice in early metoestrous or dioestrous stages showed more than a twofold increase in bactericidal activity compared with that from mice in late proestrous or early oestrous stages. These findings were true of the bactericidal activity of vaginal fluid, and the degree of bactericidal activity in it was a little higher than in serum. Protection was paralleled by the bactericidal activity in samples of serum and vaginal fluid. These data indicated that vaginal fluid as well as serum from mice immunised with purified ribosomal fraction had sufficient bactericidal activity to reduce infectivity. The fact that the degree of bactericidal activity changes with mouse oestrous cycles, however, might suggest that gonococcal activity, which was conferred by antibody to purified ribosomal fraction, entailed hormonal activity.

Table III shows absorption study results with appreciably reduced bactericidal activity in both samples after they had been absorbed with purified ribosomal fraction. Absorption with lipopolysaccharide, pilous, or outer membrane antigens did not affect the bactericidal activity of either sample. This strongly indicated that purified ribosomal fraction was one of the major antigens capable of evoking the gonococcal antibody that was secreted into the genital tract.

ANALYSIS OF CELLULAR RESPONSES IN THE GENITAL TRACT

To assess initial changes in the genital tract of mice immunised with purified ribosomal fraction or saline, we studied vaginal polymorphonuclear

TABLE III *Bactericidal activity of serum and vaginal fluid after absorption with gonococcal antigens*

Samples*	Absorbed with†	Bactericidal activity‡
Serum	Purified ribosomal fraction	≤ 1/8
	Lipopolysaccharide	1/32
	Outer membrane	1/32
	Pili	1/32
	Not absorbed	1/32
Vaginal fluid	Purified ribosomal fraction	≤ 1/16
	Lipopolysaccharide	1/64
	Outer membrane	1/64
	Pili	1/64
	Not absorbed	1/64

*Each sample was collected from mice in the dioestrous stage.

†Each sample was absorbed with sheep red blood cells coated with purified ribosomal fraction, lipopolysaccharide, outer membrane and pili respectively. Absorption was carried out three times.

‡Bactericidal activity expressed as the maximum dilution giving 90% reduction in cfu compared with corresponding controls without complement.

leucocyte responses after inoculation. As shown in the figure, infiltration by polymorphonuclear leucocytes was rather poor in control mice when challenged at late proestrous or early oestrous stages. The mean number of polymorphonuclear leucocytes in the vaginal fluid of mice at these oestrous stages was $1.5 \times 10^6/1$ 24 hours after inoculation, while in mice at the dioestrous stage it ranged from $50 \times 10^6/1$ to $500 \times 10^6/1$. In immunised mice, polymorphonuclear leucocyte response was very high regardless of the stage of oestrus at which mice were challenged, with a mean greater than $100 \times 10^6/1$ vaginal fluid 24 hours after inoculation. Two distinct differences in polymorphonuclear leucocyte response between immune mice and controls were as follows: (a) responses became appreciable in immune mice as soon as 12 hours after inoculation, and (b) the level of response in immune mice became more than 10

FIGURE 1. Kinetics of vaginal leucocyte responses of ddY mice (a) controls and (b) immunised mice) inoculated intravaginally with gonococci. Mice in various oestrous stages were challenged with 10^6 cfu strain PH2, and vaginal fluids were collected in 1 ml saline. Leucocyte responses were estimated by numbers of polymorphonuclear leucocytes (PMNL) ($\log_{10} \times 10^6$ /l of fluid).

times as great as that in controls. The number of polymorphonuclear leucocytes in immune mice at the dioestrous stage was not as large 24 hours after inoculation as that in non-immune mice at the same stage.

DELAYED HYPERSENSITIVITY

Table IV shows the effect on infectivity of delayed hypersensitivity to ribosomal proteins. On days 14 and 21, when footpad responses to ribosomal proteins were maximum in immunised mice, the protection afforded was greatly enhanced. During the second to third week after immunisation, however, activity of antibody to purified ribosomal fraction in the serum of immune mice was found to be highest. These data indicated that both cellular and humoral factors contributed to mouse protection afforded by purified ribosomal fraction.

Discussion

The sensitivity to gonococcal infection in the genital tract varies between strains of mice, and the ddY strain can be the most susceptible to vaginal inoculation of gonococci. Our preliminary study showed that not only ddY but also ICR mice were susceptible to genital infection with *N. gonorrhoeae*. CF1 mice were also available for the study of gonococcal

TABLE IV. Effect of delayed hypersensitivity to ribosomal proteins on mouse protection

Days after immunisation	Immunogen*	Mean (SD) footpad increases (mm)†	Protection (%)‡
0	Ribosomal	0.29 (0.12)	5
	Saline	0.28 (0.15)	0
7	Ribosomal	0.64 (0.05)	35
	Saline	0.27 (0.06)	5
14	Ribosomal	0.85 (0.08)	90
	Saline	0.29 (0.05)	0
21	Ribosomal	0.92 (0.06)	75
	Saline	0.28 (0.04)	5

*Each group consisted of 20 mice receiving 40 μ g purified ribosomal fraction or 0.2 ml saline.

†Footpad response was measured 48 hours after an injection of antigens and expressed as the mean values of 10 mice. Data were obtained from two separate experiments.

‡The other 10 mice were challenged intravaginally with 10^6 cfu of strain PH2 in the late prooestrous or early oestrous stages. No of mice with uterine specimens yielding negative cultures, expressed as percentage of mice studied. P values with respect to controls (saline) were calculated by Fisher's exact test.

genital infection but the response of polymorphonuclear leucocytes in the vagina of this mouse strain was poor compared with reactions in the other two strains. This ddY strain was originally produced in West Germany and imported to Japan. We have no genetic information on this strain, but it is very susceptible to bacterial infections such as salmonellosis and listeriosis. The activity of complement in the serum of ddY mice has not been examined in detail. The haemolytic activity of whole complement for sensitised sheep erythrocytes can be measured, but the characteristics of each component are less well understood. It has been reported that mouse strains which lack the fifth component of complement ($C5^-$) are more susceptible to gonococcal infection.¹⁹ In this study we saw that the peritoneal macrophages activated by the crude ribosomal fraction of gonococci showed an immediate uptake of gonococci because of the presence of lipopolysaccharide, unlike the $C5^-$ macrophages which, although they were activated, showed a less efficient rate of phagocytosis. Because of these findings the ddY strain was not thought to be $C5^-$.

The most important factor affecting the infectivity of the gonococcal strains used in this model is the preservation of the gonococci (T1 or T2) by the gelatine disc method.¹¹ We have examined 30 gonococcal strains (T1 or T2) and have shown that only strains preserved by this method can survive in the mouse uterus. Strain PH2, which was used in this study, was not different from other T2 gonococcal strains, except that it had been preserved by the gelatine disc method since its isolation from the patient.

Immunisation with 40 µg purified ribosomal fraction could not completely inhibit gonococcal invasion of the uterine epithelial layer, but gave 90% protection. Cooper *et al.*^{20, 21} reported that protection against subcutaneous infection with *N gonorrhoeae* in guinea pigs correlated with the presence of bactericidal and agglutinating antibodies in the serum of immune animals, and that lipopolysaccharide was the major antigen responsible for the bactericidal activity seen in serum immune to ribosome.⁶ In the present study the ribosomal fraction was purified by affinity chromatography using antibodies to outer membrane complex conjugated to Sepharose 4B to remove contaminants that react with outer membrane proteins and lipopolysaccharides. Mouse serum with antibodies to purified ribosomal fraction did not react with gonococcal lipopolysaccharides or outer membrane complex preparations in the immunodiffusion test, and gave one major line of precipitation against purified ribosomal fraction preparation. The results of bactericidal assays have shown that vaginal fluid had higher levels of bactericidal activity than matched serum from immune mice. Samples of serum and vaginal fluid from control mice in early metoestrous or dioestrous stages had low, but detectable, bactericidal activity, although those from control mice in late proestrous or early oestrous stages had no detectable bactericidal activity.

The bactericidal activity of vaginal fluid from control mice in the dioestrous stage was comparable with that from immune mice in late proestrous or early oestrous stages. Infectivity in control mice at the dioestrous stage, however, was significantly higher than in immune mice at late proestrous or early oestrous stages, when control mice were the most susceptible to gonococcal infection. This finding indicates two possibilities. The first is that antibody to purified ribosomal fraction may serve as an opsonin and enhance the uptake of gonococci by polymorphonuclear leucocytes. Although it is uncertain whether antibodies to microsomal antigens (such as antibody to purified ribosomal fraction) can react with intact bacteria, the effect of reproductive hormones (especially progesterone) has been reported to enhance cell leakage resulting from the binding of progesterone by membrane proteins and phospholipids.²² This phenomenon may lead to the direct interaction between ribosomal substances and anti-serum to purified ribosomal fraction. We based this theory on our preliminary study which showed that the incubation with progesterone of *N gonorrhoeae* labelled with ³H-uracil enhanced the release of ³H-labelled RNA or nucleotides from cells (unpublished observation). The second possibility is that the expression of protection in the mouse may entail

non-humoral defence mechanisms. In man, anti-gonococcal antibodies in genital secretions belong to immunoglobulin G and A, and an appreciable amount of specific IgG is produced.²³ Our previous work showed that antibody to gonococcal purified ribosomal fraction was of the IgG class.¹³ The absorption tests showed that the higher gonococcal activity of serum and vaginal fluid obtained from immune mice was attributable to the presence of antibody to purified ribosomal fraction. There is a great difference between the genital tract and the subcutaneous chamber in the major antigen responsible for the bactericidal activity of fluids obtained from the site of inoculation.⁶

Reproductive hormones are another important humoral factor in the genital tract. The bactericidal activity of serum and vaginal fluid obtained from mice in early metoestrous and dioestrous stages may be enhanced by the effect of progesterone, although the concentration of hormones was not measured in this study. This assumption is based on the report by Salt showing that oestrogen enhanced gonococcal growth, whereas progesterone greatly inhibited it.²⁴ Furthermore, Rank *et al.* showed that oestradiol treatment produced chlamydial genital infection of greater intensity and longer duration than in control guinea pigs.²⁵ Previous work in our laboratory has shown that female mice were more susceptible to genital infection with *N gonorrhoeae* at late proestrous or early oestrous stages, when oestrogen levels are highest.⁴ As many factors are related to hormonal activity, the exact relation between it and defence systems in the genital tract of mice requires further study.

Cellular responses in immune mice after gonococcal inoculation were different from those in control mice. Infiltration of polymorphonuclear leucocytes was seen much earlier in immune mice than in control mice, and immune mice showed about a hundredfold increase in the total leucocyte counts in the genital tract 24 hours after challenge. In addition, immunisation with purified ribosomal fraction induced delayed hypersensitivity, as measured by footpad responses. This could be detected one week after immunisation, and infectivity was greatly reduced after delayed hypersensitivity had been induced. These findings suggest a role for cellular immunity in protection of the mouse against gonococcal genital infection. The actual contribution of delayed hypersensitivity to an early response in the genital tract of immune mice, however, is not clear in this study. This is because the chemotactic factor for polymorphonuclear leucocytes, as one of the T cell products in a delayed hypersensitivity reaction has not been identified, as it has for mononuclear cells. It is easier to understand

that the rapid response of polymorphonuclear leucocytes in the genital tract of immune mice after gonococcal inoculation may be due to the chemotactic factor (C5a) produced by the complement mediated reaction, not only because of the presence of C5 in this mouse strain but also because there is high activity of antibody to purified ribosomal factor in the serum of immune mice.

Although there are some questions about immunity to gonococcal infection in the genital tract of mice immunised with purified ribosomal fraction, this study shows that an early polymorphonuclear leucocyte response mediated by antibody and complement interaction can reduce infectivity, and that the reproductive hormones may be responsible for enhancing gonococcal activity in serum and vaginal fluid. IgG antibody to purified ribosomal factor is present in vaginal secretion and plays an important part in gonococcal activity in the genital tract. Thus the major factor in defence mechanisms of the genital tract is very different from those of the subcutaneous chamber. This difference indicates that studies on infection with and immunity to *N. gonorrhoeae* should concentrate on the genital tract because the subcutaneous chamber is not the natural site of infection with *N. gonorrhoeae*.

Studies are in progress to identify the mechanisms by which antibody to purified ribosomal fraction kills gonococci, the active moiety of purified ribosomal fraction, and the effect of reproductive hormones on the defence systems against gonococcal infection in the genital tract.

References

- Arko RJ. *Neisseria gonorrhoeae*: experimental infection of laboratory animals. *Science* 1972; **177**:1200-1.
- Faster RS, Vinson JW. Chicken embryo as an animal model for gonorrhea. *Infect Immun* 1977; **16**:568-74.
- Lucas CT, Chandler F Jr, Martin JE Jr, Schmale JD. Transfer of gonococcal urethritis from man to chimpanzee. An animal model for gonorrhea. *JAMA* 1971; **216**:1612-4.
- Kita E, Matsuura H, Kashiba S. A mouse model for the study of gonococcal genital infection. *J Infect Dis* 1981; **143**:67-70.
- Turner WH, Novotony P. The inability of *Neisseria gonorrhoeae* pili antibodies to confer immunity in subcutaneous guinea pig chambers. *J Gen Microbiol* 1976; **92**:224-8.
- Wannemuehler JM, Miller RD, Cooper MD. Characterization of the immune response in subcutaneous chambers of guinea pigs immunized with a ribosomal preparation from *Neisseria gonorrhoeae*. *Infect Immun* 1982; **37**:469-73.
- Hilton AL, Richmond SJ, Milne JD, Hindley F, Clarke SKR. Chlamydia A in the female genital tract. *Br J Vener Dis* 1974; **50**:1-9.
- Allen E. The oestrous cycle in the mouse. *Am J Anat* 1922; **30**:297-349.
- Thung PH, Boot LM, Muhlbock O. Senile changes in the oestrous cycle and ovarian structure in some inbred strains of mice. *Acta Endocrinol (Copenh)* 1956; **23**:8-32.
- Kellog DS Jr, Peacock WL, Deacon WE, Brown L, Pirkle CI. *Neisseria gonorrhoeae*. I. Virulence genetically linked to clonal variation. *J Bacteriol* 1963; **85**:1274-9.
- Yamai S, Obara Y, Nikkawa T, Shimoda Y, Miyamoto Y. Preservation of *Neisseria gonorrhoeae* by the gelatin-disc method. *Br J Vener Dis* 1979; **55**:90-3.
- Kita E, Kashiba S. Immunogenicity of the ribosomal fraction of *Salmonella typhimurium*: analysis of humoral immunity. *Infect Immun* 1980; **27**:197-203.
- Kita E, Kashiba S. Passive hemagglutination test for detection of antibody to gonococcal ribosomal antigen in sera from patients with asymptomatic gonorrhea. *J Clin Microbiol* 1982; **15**:558-676.
- Karkhanis VK, Anderson RL, Zekner JY, Carlo DJ, Stoudt TH. Antigenic polypeptide complex from the Melvin strain of *Neisseria gonorrhoeae*: isolation and properties. *Infect Immun* 1979; **25**:635-44.
- Wolf-Watz H, Normark S, Bloom GD. Rapid method for isolation of large quantities of outer membrane from *Escherichia coli* K-12 and its application to the study of envelope mutants. *J Bacteriol* 1973; **115**:1191-7.
- Brinton CC, Bryan J, Dillon JA, et al. Uses of pili in gonorrhea control: role of bacterial pili in disease, purification and properties of gonococcal pili, and progress in the development of a gonococcal pilus vaccine for gonorrhea. In: Brooks GF, Gotschlich EC, Holmes KK, Sawyer WD, Young FE, eds. *Immunobiology of Neisseria gonorrhoeae*. Washington, DC: American Society for Microbiology, 1978:155-78.
- Westphal O, Jan K. Bacterial lipopolysaccharides. *Methods in Carbohydrate Chemistry* 1965; **5**:81-91.
- Fogel S, Sypherd PS. Extraction and isolation of individual ribosomal proteins from *Escherichia coli*. *J Bacteriol* 1968; **96**:358-64.
- Cooper MD, Floyd SA. In vitro kinetics of phagocytosis and intracellular killing of gonococci by peritoneal macrophages from mice deficient in complement component 5. *Infect Immun* 1982; **36**:363-70.
- Cooper MD, Tewari RP, Bowers DV. Immunogenicity of ribosomal preparations from *Neisseria gonorrhoeae*. *Infect Immun* 1980; **28**:92-100.
- Cooper MD, Wannemuehler MD, Miller RD, Fedyk MF. Role of outer envelope contamination in protection elicited by ribosomal preparations against *Neisseria gonorrhoeae* infection. *Infect Immun* 1981; **32**:173-9.
- Morse SA, Fitzgerald TJ. Effect of progesterone on *Neisseria gonorrhoeae*. *Infect Immun* 1973; **10**:1370-7.
- Tramont CE, Ciak J. Antigonococcal antibodies in genital secretions. In: Brooks GF, Gotschlich EC, Holmes KK, Sawyer WD, Young FE, eds. *Immunobiology of Neisseria gonorrhoeae*. Washington DC: American Society for Microbiology, 1978:274-8.
- Salt IE. The differential susceptibility of gonococcal opacity variants to sex hormones. *Can J Microbiol* 1982; **28**:301-6.
- Rank RG, White HJ, Hough Jr, Pasley JN, Barron AL. Effect of estradiol on chlamydial genital infection of female guinea pigs. *Infect Immun* 1982; **38**:695-701.