

IMUNOVIRTM

inosine pranobex

the body's ally
against viral infection



The immunopotentiator proven in herpes

Prescribing Information

Dosage: Two tablets q.d.s. for 7 to 14 days.

Presentation: Each tablet of IMUNOVIR contains 500 mg. inosine pranobex. **Uses:** Imunovir is an agent demonstrating anti-viral activity and possessing immunopotentiating action in viral diseases. **Indications:** Imunovir is indicated in the management of mucocutaneous infections due to herpes simplex virus (type I and or type II). **Contra-Indications:** None known. **Warnings:** As the inosine component of Imunovir is metabolised to uric acid, it should be used with caution in patients with renal impairment, a history of gout or hyperuricaemia. **Pregnancy:** Although animal tests have shown no teratogenic effect, the use of Imunovir in women where pregnancy is suspected or confirmed, should be avoided. **Adverse Effects:** Side effects are rare and usually of a mild and transitory nature. The only commonly associated adverse

effects occurring during treatment with Imunovir are elevated serum and urinary concentrations of uric acid. These return to normal once treatment is withdrawn. **Overdose:** There has been no experience of overdosage with Imunovir. However, serious adverse effects, apart from increased levels of uric acid in the body, seem unlikely in view of the animal toxicity studies. Treatment should be restricted to symptomatic and supportive measures. **Product Licence No:** 0748/0049. **Basic N.H.S. Price:** 100 £38.00.



Edwin Burgess Limited
Longwick Road, Princes Risborough
Aylesbury, Bucks. HP17 9RR

TM Trademark of the Licensor, Newport Pharmaceuticals Int. Inc.

1 IS ENOUGH



BAYER

**3-day action against thrush
in a single tablet**

Canesten 1
clotrimazole

"Treatment of vaginal candidosis with a single 500-mg clotrimazole pessary [Canesten 1] was found to be as effective as previous treatment regimens of longer duration."

Br J Ven Dis 1982;58:124-6.

Prescribing information: Presentation: Canesten 1 is available as a single 500mg vaginal tablet with applicator. **Dosage:** A single vaginal tablet to be inserted preferably at night. **Side effects:** Rarely patients may experience transient local mild burning or irritation. **Basic NHS Cost:** 1 x 500mg vaginal tablet with applicator: £2.63. **Product Licence Number:** Vaginal Tablets 500mg, PL0010-0083.



Further information is available from:
Bayer UK Limited Pharmaceutical Division
Bayer House Strawberry Hill
Newbury Berkshire RG13 1JA
Telephone: Newbury 06357 39000

* Registered trade mark of Bayer, Germany

M.S.S.V.D.

The Medical Society for the Study of Venereal Diseases

President: Dr. B. A. Evans

The object of the Society is to bring together those interested or engaged in the diagnosis, treatment, and public health aspects of venereal diseases. The Society meets regularly in London for the reading of papers, discussions, and presentation of cases. An out-of-London meeting is held in the Spring.

Annual subscription for membership of the Society is £35 for all members.

Applications for membership are very welcome and forms can be obtained from the Honorary Secretary, Dr. M. A. Waugh, Department of Genito-Urinary Medicine, The General Infirmary, Great George Street, Leeds LS1 3EX.

Please note—all correspondence regarding MSSVD to Hon. Sec.

Programme—1985

Nevin Lecture Theatre, St. Thomas' Hospital, London SE1 7EH, 19.00 hours.

Friday 29th March 1985

Symposium on Recent Aspects of STD

- (1) Plasmids in the Gonococcus—Dr. A. E. Jephcott, Bristol
- (2) Chlamydial Cervicitis: Findings on Colposcopy and Biopsy—Dr. E. Dunlop, London
- (3) Cerebro-spinal Fluid Changes in Syphilis—Dr. E. J. Thompson, London


Friday 26th April 1985

Presidential Address: Dr. B. A. Evans, London

16th-19th May 1985

Spring Meeting in conjunction with Scandinavian Societies, Dept. of Dermatology and Venereology, University Hospital, Uppsala, Sweden

This publication is available in microform.



University
Microfilms
International

University Microfilms International reproduces this publication in microform: microfiche and 16mm or 35mm film. For information about this publication or any of the more than 13,000 titles we offer, complete and mail the coupon to: University Microfilms International, 300 N. Zeeb Road, Ann Arbor, MI 48106. Call us toll-free for an immediate response: 800-521-3044. Or call collect in Michigan, Alaska and Hawaii: 313-761-4700.

Please send information about these titles:

Name _____


Company/Institution _____

Address _____

City _____

State _____ Zip _____

Phone () _____



The growing risk of Chlamydial infection

In men – Chlamydial infection is implicated in over half the cases of non-gonococcal urethritis (NGU).

Doxycycline – effective in curing Chlamydial urethritis

Pharmatherapeutica 1980; 2 (5): 341

In women – it is responsible for an increasing number of cases of pelvic inflammatory disease (PID). If untreated, the infection may cause endometritis or salpingitis.

Doxycycline – eradicates urinary symptoms, pyuria, and infecting micro-organisms.

New Engl. J. Med., 1981; 304: 956

DOXATET ROUTINELY



PRESCRIBING INFORMATION. Uses: infections caused by a wide range of organisms, including respiratory tract, urinary tract, soft tissue and gastro-intestinal infections. **Dosage:** two tablets (200 mg) first day, followed by 1 tablet/day. In severe infections give 200 mg daily, see detailed dosage in data sheet. **Note:** absorption is virtually unaffected by food or milk. **Contra-indications:** hypersensitivity to tetracyclines. **Side effects:** occasionally, gastro-intestinal effects, skin rashes. **Tooth discolouration** is a possible sequel of treatment in the latter half of pregnancy or in infancy – up to the age of 12 years. **Note:** Tetracyclines are also found in the milk of lactating women receiving therapy. **Packs:** Blister packs of 10 tablets (each containing doxycycline 100 mg) Basic NHS cost £4.25 per 10 tablets. **Product licence number:** 0142/0186. **Legal category:** POM.

Full information on request: Cox Pharmaceuticals, Whiddon Valley, Barnstaple, Devon EX32 8NS Tel: (0271) 75001

