Brachioproctic eroticism and transmission of retrovirus associated with acquired immune deficiency syndrome (AIDS)

BASIL DONOVAN,* BRETT TINDALL,† AND DAVID COOPER‡ (FOR THE SYDNEY AIDS STUDY GROUP)

*From the *Taylor Square Private Clinic, the †Centre for Immunology, St Vincent's Hospital, and the ‡National Health Medical Research Centre for Clinical and Epidemiological AIDS Research, University of New South Wales, Sydney, Australia*

SUMMARY A case of transmission of the acquired immune deficiency syndrome (AIDS) associated retrovirus (ARV) by brachioproctic eroticism is described.

Introduction

The lymphocytotropic retroviruses variously known as lymphadenopathy associated virus (LAV),¹ human T cell lymphotropic virus type III (HTLV-III),² and the acquired immune deficiency syndrome (AIDS) associated retrovirus (ARV)³ are now well documented as the cause of AIDS and related conditions. These viruses have been isolated from blood,¹⁻³ semen,⁴ saliva,⁵ breast milk,⁶ and tears.⁷ The major routes of transmission documented to date include anal intercourse,⁸ vaginal intercourse,⁹ transfusion of blood¹⁰ and blood products,¹¹ sharing of needles and syringes by intravenous drug users,¹² transmission from mother to child in the perinatal period,¹³ and artificial insemination.¹⁴ Reports of needlestick transmission in

a health care worker¹⁵ and suggestive of transmission via breast feeding¹⁶ have also appeared. We describe a probable case of transmission of the AIDS associated retrovirus (ARV) by brachioproctic eroticism ("fisting").

Case report

A 43 year old homosexual man was recruited into the Sydney AIDS prospective study¹⁷ in August 1984. His medical history included solar keratoses with ulceration on his face and the back of his hands for many years, acute hepatitis B in 1972, and a syphilis chancre on the back of his left hand in 1982. He had no other history of sexually transmissible diseases (STDs). On enrolment and in April 1985 he was symptomless with no lymphadenopathy or hepatosplenomegaly. On both occasions antibodies to ARV by immunofluorescence (IF), enzyme linked immuno-sorbent assay (ELISA), and radioimmuno-precipitation assay (RIPA) were not detected in his serum, and his T cell subsets were normal.

In mid June 1985 the patient suffered sudden onset of lethargy and passed dark urine and pale stools for one day. Eleven days later he developed generalised myalgias and arthralgias, fevers, night sweats, severe malaise, and a diffuse swelling of the right buccal mucosa. These symptoms settled during three weeks, but he became depressed and irritable and developed a non-productive cough and nasal congestion that persisted for seven weeks. In August 1985 he developed generalised lymphadenopathy, and on review in September physical examination showed hepatosplenomegaly, several crusted solar keratoses and an atrophic chancre scar on the left hand, truncal

Address for reprints: Dr B Donovan, Taylor Square Private Clinic, 302 Bourke Street, Sydney, New South Wales 2010, Australia

Accepted for publication 18 March 1986

SYDNEY AIDS STUDY GROUP

Consultants: A I Adams, D G Fox, G Berry, J Dykes.

Project principal investigators: D A Cooper, J Gold.

Centre for Immunology, St Vincent's Hospital, Sydney: R Penny, P Borrow, G Chapman, L Hurren, A Imrie, C Katakouzinos, S Kehrer, H McCabe, A Malcolm, M North, E Schreurs, G Smith, H Smith, B Tindall, R Wilson

School of Public Health and Tropical Medicine, Sydney University: J Burcham

Albion Street Centre, Sydney Hospital: M Anns, D Grant, C Kronenberg, M Perdices, P Todd.

STD Centre, Sydney Hospital: R Philpot, R Jones, J Roberts

Participating medical practitioners: T G Barnes, P Brooke, B Donovan, R Finlayson, P Haynes, R Lovett, H M Michelmores, K Mutimer, R Price.

prurigo, and patches of atrophic papillae on his tongue. Investigations in early July showed mild lymphopenia, an erythrocyte sedimentation rate (Westergen) of 14 mm in the first hour, slightly raised transaminase activity, and a negative test for antibodies to ARV. In August his serum was positive for antibodies to ARV (by ELISA and IF). The blood film showed eosinophilia ($870 \times 10^6/l$) and atypical mononuclear cells. Heterophilic antibodies were detected, but IgM antibodies to cytomegalovirus, Epstein-Barr virus, and *Toxoplasma gondii* were not found. The laboratory data showing changes in ARV antibody and T cell subsets in this patient are presented in the table.

The patient's regular sexual partner had a continual history of lethargy and night sweats from 1983. In 1983 he was noted to have generalised lymphadenopathy and hepatosplenomegaly. He was treated for anal condylomata in 1985. Testing in 1983 had shown anergy, an inverted ratio of T4 to T8 cells (0.80), and moderately raised serum transaminase activity. In August 1984 antibodies to ARV were detected in his serum by IF, ELISA, and RIPA, and the results of T cell subset testing are shown in the table.

The couple's sexual practices since early 1984 consisted of "dry" kissing, mutual masturbation to ejaculation, and brachioproctic eroticism (insertion of the patient's unprotected left hand at least 15 cm into his partner's rectum). His partner had noted occasional slight rectal bleeding after this practice, which occurred about once a week. The patient disliked anogenital, orogenital, and oroanal sexual practices and had not engaged in these for more than 18 months

before seroconversion. He has neither used intravenous drugs nor received blood products.

Discussion

We think that the patient probably contracted ARV infection through the practice of brachioproctic eroticism between April and June 1985, after a long potential exposure to the retrovirus. The acute illness, which began in June 1985, had similar features to other illnesses associated with seroconversion.¹⁸

Though insertive and receptive brachioproctic eroticism have been significantly associated with ARV seropositivity in homosexual men,^{19,20} most men who engage in this practice also engage in anogenital and orogenital sex. Brachioproctic eroticism may indicate a high degree of involvement in homosexual practices. This patient, however, had avoided penetrating sexual practices other than brachioproctic eroticism for a long time, which was confirmed by the negative history of STDs except for hepatitis B and the chancre on his hand.

Though epidemiological evidence strongly suggests that ARV is not transmitted through intact skin, this patient had persistent keratotic ulcers on his hand, which had been frequently and directly exposed to his partner's anorectal mucosa and probably blood for almost two years. In addition, he had a history of transmission of *Treponema pallidum* by identical means. It is not impossible, however, that ARV could have been transmitted in this instance by kissing or by contact with his partner's semen either through his hand lesions or through accidental mucosal contact.

TABLE Laboratory data on reported patient and his sexual partner and 63 controls

Date	Antibodies to ARV	Lymphocyte count ($\times 10^6/l$)	T4 cells ($\times 10^6/l$)	T8 cells ($\times 10^6/l$)	Ratio of T4 to T8
Patient					
August 1984	Negative	2400	1100	340	3.20
April 1985	Negative	2300	920	300	3.10
*					
July 1985	Negative	1100	NT	NT	NT
September 1985	Positive	3200	540	510	1.10
December 1985	Positive	2500	710	560	1.30
Partner					
1983	NT	NA	NA	NA	0.80
August 1984	Positive	1200	290	440	0.65
Controls					
mean (SD)	Negative	2100 (330)	930 (230)	410 (100)	2.40 (0.55)

ARV = AIDS associated retrovirus; NT = not tested; NA = not available.

*Onset of symptoms in June 1985.

On the basis of this report, we recommend covering any skin lesions that might be exposed directly to body fluids or persons with or at risk of infection with ARV. It would be prudent for anybody engaging in brachio-proctic eroticism to wear latex gloves.

References

1. Barre-Sinoussi F, Chermann JC, Rey F, *et al.* Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 1983;**220**:868-71.
2. Gallo RC, Salahuddin SZ, Popovic M, *et al.* Frequent detection and isolation of cytopathic retroviruses (HTLV-III) from patients with AIDS and at risk for AIDS. *Science* 1984;**224**:500-3.
3. Levy JA, Hoffman AD, Kramer SM, *et al.* Isolation of lymphocytopathic retroviruses from San Francisco patients with AIDS. *Science* 1984;**225**:840-3.
4. Zagury D, Bernard J, Leibowitch J, *et al.* HTLV-III in cells cultured from semen of two patients with AIDS. *Science* 1984;**226**:449-51.
5. Groopman JE, Salahuddin SZ, Sarngadharan MG, Gonda M, Sliiski A, Gallo RC. HTLV-III in saliva of people with AIDS-related complex and healthy homosexual men at risk for AIDS. *Science* 1984;**226**:447-9.
6. Thiry L, Sprecher-Goldberger S, Jonckheer T, *et al.* Isolation of AIDS virus from cell-free breast milk of three healthy virus carriers. *Lancet* 1985;ii:891-2.
7. Fujikawa LS, Salahuddin SZ, Palestine AG, Masur H, Nussen Blatt RB, Gallo RC. Isolation of human T-lymphotropic virus type III from the tears of a patient with the acquired immunodeficiency syndrome. *Lancet* 1985;ii:529-30.
8. Goedert JJ, Sarngadharan MG, Biggar RJ, *et al.* Determinants of retrovirus (HTLV-III) antibody and immunodeficiency conditions in homosexual men. *Lancet* 1984;ii:711-6.
9. Redfield RR, Markham PD, Salahuddin SZ, Wright DC, Sarngadharan MG, Gallo RC. Heterosexually acquired HTLV-III/LAV disease (AIDS related complex and AIDS). Epidemiologic evidence for female-to-male transmission. *JAMA* 1985;**254**:2094-6.
10. Curran JW, Lawrence DN, Jaffe H, *et al.* Acquired immunodeficiency syndrome (AIDS) associated with transfusions. *N Engl J Med* 1984;**310**:69-75.
11. Lederman MM, Ratnoff OD, Evatt BL, McDougal JS. Acquisition of antibody to lymphadenopathy-associated virus in patients with classic haemophilia (factor VIII deficiency). *Ann Intern Med* 1985;**102**:753-7.
12. Friedland GH, Harris C, Butkus-Small C, *et al.* Intravenous drug abusers and the acquired immunodeficiency syndrome (AIDS). Demographic, drug use and needle-sharing patterns. *Arch Intern Med* 1985;**145**:1413-7.
13. Rubinstein A, Sicklick M, Gupta A, *et al.* Acquired immunodeficiency with reversed T4/T8 ratios in infants born to promiscuous and drug-addicted mothers. *JAMA* 1983;**249**:2350-6.
14. Stewart GJ, Tyler JPP, Cunningham AL, *et al.* Transmission of human T cell lymphotropic virus type III (HTLV-III) by artificial insemination by donor. *Lancet* 1985;ii:581-4.
15. Anonymous. Needlestick transmission of HTLV-III from a patient infected in Africa. *Lancet* 1984;ii:1376-7.
16. Ziegler JB, Cooper DA, Johnston RO, Gold J. Postnatal transmission of AIDS-associated retrovirus from mother to infant. *Lancet* 1985;ii:896-7.
17. Sydney AIDS Study Group. The Sydney AIDS Project. *Med J Aust* 1984;**141**:569-73.
18. Cooper DA, Gold J, Maclean P, *et al.* Acute AIDS retrovirus infection. Definition of a clinical illness associated with seroconversion. *Lancet* 1985;ii:537-40.
19. Nicholson JKA, McDougal JJ, Jaffe W, *et al.* Exposure to human T-lymphotropic virus type III/lymphadenopathy associated virus and immunologic abnormalities in asymptomatic homosexual men. *Ann Intern Med* 1985;**103**:37-42.
20. Jeffries E, Willoughby B, Boyko W, *et al.* The Vancouver lymphadenopathy-AIDS study: 2. Seroepidemiology of HTLV-III antibody. *Can Med Assoc J* 1985;**132**:1373-7.