Mucopurulent cervicitis: a clinical entity?

F E WILLMOTT

From the Department of Genitourinary Medicine, Royal South Hants Hospital, Southampton

SUMMARY Of 297 women attending a sexually transmitted disease clinic who were examined for the presence of mucopurulent cervicitis, 96 (32%) satisfied the diagnostic criteria. Mucopurulent cervicitis was strongly associated with the isolation of *Chlamydia trachomatis* and *Neisseria gonorrhoeae*. It was also associated with bacterial vaginosis, the use of oral contraceptives, and sexual contact with men who had non-gonococcal urethritis.

Conversely, the presence of opaque cervical secretions did not show these associations, and the results indicated no difference in genital infections in women with clear cervical secretions whether or not more than 10 polymorphonuclear leucocytes per field (at a magnification of $\times 1000$) were present.

The findings support the suggestion that mucopurulent cervicitis is a definite clinical entity that requires investigation and treatment.

Non-gonococcal urethritis (NGU) is recognised by the presence of five or more polymorphonuclear leucocytes (PMNL) per field at a magnification of $\times 1000$ (high power field (HPF)) in a Gram stained smear.¹ The equivalent condition in women is termed non-specific genital infection (NSGI), but specific criteria for the diagnosis have not been defined. The clinical diagnosis of cervicitis is subjective, and differentiation between columnar cell ectopy (erosion) and cervicitis is difficult, if not impossible, with the naked eye. Cervicitis may be indicated by finding an excess of PMNL in a Gram stained smear of cervical secretions, but the number of polymorphs considered to be normal has not been defined. A condition of mucopurulent cervicitis has been described and designated the female equivalent of male NGU.² The diagnosis of mucopurulent cervicitis is based on two findings: visible mucopurulent (yellow) cervical secretions, and more than 10 PMNL/HPF in the cervical secretions. Mucopurulent cervicitis has been associated particularly with the presence of *Chlamydia trachomatis* but not *Neisseria gonorrhoeae* or herpes simplex virus.² Further studies have shown an association with *Ureaplasma urealyticum*, *Gardnerella vaginalis*, *Trichomonas vaginalis*, and the clinical diagnosis of bacterial vaginosis.³

The present study was undertaken to assess the presence of mucopurulent cervicitis in women attending a sexually transmitted diseases (STD) clinic in the United Kingdom and the validity of the definitive

criteria, particularly the importance of more than 10 PMNL/HPF in a Gram stained smear of the cervical secretions.

Patients, materials, and methods

Women attending the STD clinic of this hospital as new patients were studied. They were excluded if they were menstruating or if the cervical smear was not satisfactory (more than 100 squamous cells were present on the slide⁴). A routine history including details of contraceptive practice was taken. I examined all the patients as described by Brunham *et al.*² and noted the presence of columnar cell ectopy or cervicitis (defined as erythema or oedema within the zone of ectopy), or both. The appearance of the cervical secretions was classified as mucopurulent (yellow), opaque (white or cloudy), or clear. The presence of any bleeding on swabbing the cervix was also recorded.

Endocervical mucus was collected with a cotton wool swab rolled on to a glass slide, which was Gram stained and examined under oil immersion at a $\times 1000$ magnification. A wet smear of vaginal secretions was examined for trichomonads, yeasts, and clue cells. Cultures for *N gonorrhoeae* were taken from the urethra and cervix on Transgrow medium (blood agar containing vancomycin, colistin, and nystatin). Vaginal secretions were transported in Amies' medium and cultured for *Candida albicans* on Sabouraud's medium. Material for culture for chlamydiae was taken from the cervix, placed in sucrose phosphate broth, and transported to the laboratory within four hours. It was then stored at -70°C until the day of culture, which was performed twice weekly. The

Address for reprints: Dr F E Willmott, Department of Genitourinary Medicine, Royal South Hants Hospital, Graham Road, Southampton SO9 4PE

Accepted for publication 27 October 1987

Table 1 Findings related to age of 297 women

Age	No of women with:		No of all women (n = 297)
	Mucopurulent cervicitis (n = 96)	Chlamydial infection (n = 57)	
Under 25	64	46	172
Over 25	32	11	125

specimens were cultured in BGMK (baboon green monkey kidney) cells inhibited by cycloheximide, and were examined by indirect immunofluorescence. A specimen for culture for herpes simplex virus was taken when appropriate. The diagnosis of bacterial vaginosis was made using the criteria of Amsel *et al.*⁴

Results

Of 297 women investigated, 96 (32%) had mucopurulent, 57 (19%) opaque, and 144 (49%) clear cervical secretions. Gram stain of the cervical smears showed > 10 PMNL/HPF in all 96 with mucopurulent secretions, 55 out of 57 with opaque secretions, and 43 out of 144 with clear secretions. Induced cervical bleeding was present in 19 of all patients, but in 11 of the 96 women with mucopurulent cervicitis, eight of whom yielded chlamydiae. Clinical cervicitis was present in all 96 patients with mucopurulent secretions, in 38/57 with opaque secretions, and in 5/144 with clear secretions.

AGE

Table 1 shows that patients aged under 25 made up 58% (172/297) of all patients, but 67% (64/96) of patients with mucopurulent cervicitis, which was a significant difference ($\chi^2 = 4.4$; $p < 0.05$).

CONTRACEPTION

Table 2 shows that 50% (149) of the women used oral contraceptives, 6% (18) used intrauterine contraceptive devices (IUCDs), 9% (27) used a barrier method (sheath or diaphragm), and 35% (104) used no contraception. Of the 96 women with mucopurulent cervicitis, 69 (72%) used oral contraceptives ($\chi^2 = 26.7$; $p < 0.001$).

Table 3 Cervical mucus related to menses in 262 women

Phase of cycle	Mucopurulent mucus (n = 88)	Opaque mucus (n = 52)	Clear mucus (n = 122)	
			> 10 PMNL/HPF	< 10 PMNL/HPF
Postmenstrual	28	19	10	26
Mid-cycle	28	15	14	38
Premenstrual	32	18	14	20

PMNL = polymorphonuclear leucocytes; HPF = high power ($\times 1000$ magnification) field.

Table 2 Methods of contraception used by 297 women

Appearance of cervical mucus	No using:				Total
	Oral	IUCD	Barrier	None	
Mucopurulent	69	5	4	18	96
Opaque	29	6	3	19	57
Clear	51	7	20	66	144
Total	149	18	27	103	297

IUCD = intrauterine contraceptive device.

MENSTRUAL CYCLE

The type of cervical mucus was related to the time of the last menstrual period (35 patients with amenorrhoea of more than four weeks' duration were excluded). It was classified as postmenstrual (proliferative), mid-cycle (ovulatory), or premenstrual (secretory). Table 3 shows that mucopurulent, opaque, or clear secretions containing more than 10 PMNL/HPF were not associated with any phase of the menstrual cycle. The presence of clear mucus containing fewer than 10 PMNL/HPF appeared to be most common in the mid-cycle phase, but this was not significant.²

ASSOCIATED INFECTIONS

Cervical infections with *N gonorrhoeae*, chlamydiae, or herpes simplex virus were present in 54 women with mucopurulent, 15 with opaque, and 27 with clear mucus. Table 4 shows the incidence of individual infections. Mucopurulent cervicitis was significantly associated with gonorrhoea ($\chi^2 = 23.6$; $p < 0.001$), chlamydial infection ($\chi^2 = 34.2$; $p < 0.001$), and bacterial vaginosis ($\chi^2 = 56$; $p < 0.02$), but was negatively associated with candidiasis. Opaque secretions were not associated with any particular infection.

SEXUAL CONTACTS OF MEN WITH NGU

Table 5 shows that of 77 women who were known sexual contacts of men with NGU, 27 yielded chlamydiae and 41 had mucopurulent cervicitis.

EFFECT OF TREATMENT

Of 43 women with mucopurulent cervicitis but without gonorrhoea or chlamydial infection who received tetracyclines or erythromycin, 28 attended follow up and 19 of them showed resolution of the mucopurulent cervicitis.

Table 4 Associated infections in 297 women

Cervical infections	Mucopurulent mucus (n = 96)	Opaque mucus (n = 57)	Clear mucus		Total (n = 297)
			> 10 PMNL/HPF (n = 43)	< 10 PMNL/HPF (n = 101)	
Gonorrhoea	20	3	1	3	27
Chlamydial infection	36	10	4	7	57
Herpes simplex virus	7	4	4	9	24
Candidiasis	17	17	14	30	78
Bacterial vaginosis	40	17	9	30	96
Trichomoniasis	8	5	2	3	18
Warts	11	6	8	13	38

Table 5 Chlamydial infection in 77 female sexual contacts of men with non-gonococcal urethritis (NGU) correlated with appearance of cervical mucus

Appearance of cervical mucus	No of NGU contacts	
	Chlamydia negative (n = 50)	Chlamydia positive (n = 27)
Mucopurulent (n = 41)	22	19
Opaque (n = 16)	12	4
Clear (n = 20)	16	4

Discussion

The object of the study was to assess the importance of the physical appearance of the cervical mucus and of the presence of more than 10 PMNL/HPF in a Gram stained smear of the mucus. In this series 32% of women attending an STD clinic had mucopurulent secretions. Mucopurulent cervicitis was positively associated with the presence of chlamydiae, as shown in previous studies,^{2,3} but I know of no previous study showing an association with gonorrhoea. Bacterial vaginosis was also associated with mucopurulent cervicitis, even when patients with chlamydial infection and gonorrhoea were excluded. As bacterial vaginosis does not cause vaginitis, it is unlikely to cause cervicitis unless it has a specific effect on columnar epithelium. The cause of the cervicitis is not known, but the effect is probably damage to the epithelium that produces an environment favouring the growth of anaerobes, as suggested by Masfari *et al.*⁵ In contrast to other workers, I found no association between *T vaginalis* and mucopurulent cervicitis. As described previously,^{2,3} *C albicans* was negatively associated with mucopurulent cervicitis. The use of oral contraceptives was associated with mucopurulent cervicitis, particularly if chlamydiae were present, but not if gonorrhoea was present.

Mucopurulent cervicitis has been described as the equivalent in women of NGU in men. Of the 77 women who were known sexual contacts of men with NGU, 41 had mucopurulent cervicitis and 27 had chlamydial infection. The response to tetracyclines or erythromycin in 68% (19/28) of the non-chlamydial non-gonococcal group compares with the success of

treatment of non-chlamydial NGU in men.

Opaque cervical secretions were not associated with any infections or any phase of the menstrual cycle. Clear cervical secretions were negatively associated with chlamydial infection and gonorrhoea. Brunham *et al* used 10 PMNL/HPF as the diagnostic criterion for cervicitis.² Other studies have used a different number of polymorphs as indicative of cervicitis, for example Fox used the arbitrary figure of 50 PMNL/HPF.⁶ All patients with mucopurulent mucus had more than 10 PMNL/HPF, as did 96% (55/57) of those with opaque mucus. In the group with clear mucus, however, those with more than 10 PMNL/HPF had no more genital infections than those with fewer than 10 PMNL/HPF, and 7% (3/43) and 12% (12/101) of the two groups were sexual contacts of men with NGU. Further studies are required to assess the number of polymorphs that indicate cervicitis. A more accurate method of assessing the numbers is probably also required.

This study shows that mucopurulent cervicitis is a definite entity that can be diagnosed clinically by the presence of mucopurulent (yellow) cervical mucus, and diagnosis does not depend on the results of a Gram stained smear. As the presence of mucopurulent cervicitis is an indicator of gonorrhoea or chlamydial or non-specific genital infection in most cases, the treatment given should be effective against all these conditions if culture facilities are not available. The male sexual contacts of women with mucopurulent cervicitis should also be traced and examined.

References

- 1 Swartz SL, Kraus SJ, Harman KL, Stargel MD, Brown WJ, Allen SD. Diagnosis and etiology of nongonococcal urethritis. *J Infect Dis* 1978;138:445-54.
- 2 Brunham RC, Paavonen J, Stevens CE, *et al*. Mucopurulent cervicitis—the ignored counterpart in women of urethritis in men. *N Engl J Med* 1984;311:1-6.
- 3 Paavonen J, Critchlow CW, Rouen T, *et al*. Etiology of cervical inflammation. *Am J Obstet Gynecol* 1986;154:556-64.
- 4 Amsel R, Totten PA, Spiegel CA, Chen KCS, Eschenbach D, Holmes KK. Non-specific vaginitis: diagnostic criteria and microbial and epidemiologic associations. *Am J Med* 1983;74:14-22.
- 5 Masfari AN, Duerden BI, Kinghorn GI. Quantitative studies of vaginal bacteria. *Genitourin Med* 1986;62:256-63.
- 6 Fox H. Non-specific genital infection in a general practice. *British Journal of Venereal Diseases* 1974;50:125-31.