by culture five days after initiation of treatment. Non-viable organisms were not detected by ELISA ten days following initiation of therapy or three days following cessation of therapy. Thus, there appeared to be a five-day lag period of chlamydial antigen detection demonstrated by ELISA after culture (McCoy) confirmed cure (fig).

Both culture and EIA are reliable and may be used for tests of cure (TOC) if appropriate timing intervals are observed to avoid false positive TOC results. Since EIA was used by 50% of the United Kingdom clinics for TOC, the results of our investigation may be helpful. A reliable TOC can be obtained utilising EIA three days following completion of treatment (10 days post treatment initiation) instead of one week following treatment as reported by Radcliffe *et al.* Furthermore, other TOC indications may include pregnancy, complicated or severe infections, requests for psychological reassurance or for evidence of persistent cervicitis. However, as eloquently stated by Dr Radcliffe, a routine TOC for chlamydial endocervical infections is presently unessential.

DARON G FERRIS
Department of Family Medicine,
Medical College of Georgia,
Augusta, Georgia 30912, USA

- 1 Radcliffe KW, Rowen D, Mercey DE, *et al.* Is a test of cure necessary following treatment for cervical infection with *Chlamydia trachomatis*? *Genitourin Med* 1990;66:444-6.
- 2 Radcliffe KW, Rowen D, Mercey DE, *et al.* Survey of the management of *Chlamydia trachomatis* infection of the cervix. *Genitourin Med* 1991; 67:41-43.
- 3 Ferris DG, Lawler FH, Horner, *et al.* Test of cure for genital *Chlamydia trachomatis* infection in women. *J Fam Pract* 1990;31:36-41.

Syphilis and the elderly

Dr B Chattopadhyay¹ reports a case of a 91-year-old man found to have positive serological tests for syphilis prior to emigration. He states the difficulties encountered in accurate diagnosis.

Two geriatricians, a psychiatrist and myself² in 1987 reviewed 659 consecutive medical and psychiatric admissions in the elderly to St James's Hospital, Leeds, where standard

serological tests for syphilis were performed.

The ages ranged between 65 and 103 years (mean 81); 210 were men; 449 were women. Positive serological tests for syphilis were found in 23 (3.5%) of the patients, whose ages ranged from 69 to 93 (mean 82.5). Seventeen were women and six were men. Even after extensive investigation 11 (1.7%) could only be diagnosed as latent syphilis.

We were helped in diagnosis in that all the patients were white and a confidential venereal diseases treatment register exists in Leeds going back to 1919 in which we were able to check previous diagnoses.

MA WAUGH
General Infirmary,
Leeds, LS1 3EX, UK

- 1 Chattopadhyay B. Syphilis and the elderly. *Genitourin Med* 1991;67: 270-1.
- 2 Corrado OJ, Bowie PCW, Bagnall WE, Waugh MA. The prevalence of positive serological tests for syphilis among elderly hospital patients. *Age Ageing* 1989;18:407-10.

BOOK REVIEWS

Genital Papillomavirus Infections. Modern Diagnosis and Treatment. Edited by G Gross, S Jablonska, H Pfister, HE Stegner. (Pp 452; Price DM 148) Berlin. Springer-Verlag. ISBN 3-540-52615-3.

This monograph contains the documentation of *The International Symposium on Genital Papillomavirus Infection* held in Hamburg in 1989. It covers epidemiology, molecular biology, diagnosis, clinical aspects, immunological aspects, and therapy. The sixty-two contributors from ten countries are drawn from a wide range

of specialties which deal with papillomavirus infection and the sequelae of such infection.

The sections on molecular biology and diagnosis are especially well presented. The authors ease the uninitiated painlessly into the world of papillomavirus biology. Terminology is explained such that non-specialist readers could readily follow the text without recourse to a dictionary of molecular biology.

The 11 chapters on clinical aspects economically cover a wide subject matter, and are generally well written. Illustrations, particularly the photomicrographs of cervical cytology were of a high quality. My only criticisms of this section concerned the DNA histograms which required further explanation for them to make their full impact, and that the chapter on vulval dystrophies seemed to be misplaced in this book, as papillomavirus is not mentioned at all.

The chapter entitled *Immunology of Genital Papillomavirus Infections* eloquently reviews humoral immunity, cell-mediated immunity, cell-mediated cytotoxicity and local immune mechanisms. In a concise summary the relative importance of these individual components of the immune response to H.P.V., and the implications on future development of therapies is discussed.

I was a little disappointed not to find any information concerning loop excision diathermy in the section concerning therapy, although at the time of this conference this modality of treatment was just emerging. My disappointment was off-set by chapters on the basic science of cryotherapy and laser therapy.

Overall this book is well written and presented. It is not for the molecular biologist, but for the clinician with a interest in papillomavirus infection. The price may limit its sales but any clinician working in the field of such infections would, in my opinion, be advised to ensure that they have access to a copy in their clinics or libraries.

DAVID ROWEN

Economic Aspects of AIDS and HIV Infection. Edited by D Schwefel, R Leidl, J Rovira, M F Drummond (Pp 364, DM 148). Berlin: Springer-Verlag, 1990.