A serological test for granuloma inguinale

A L Freinkel, Y Dangor, H J Koornhof, R C Ballard

Abstract

Objectives—An indirect immunofluorescence technique applied to paraffin embedded tissue sections of lesions containing Donovan bodies was evaluated as a serological test for the diagnosis of granuloma inguinale.

Methods—Sera from patients with proven granuloma inguinale, other sexually acquired genital ulcerations and blood donors from areas where granuloma inguinale is rarely encountered as well as from disease-endemic regions were tested. Sera were tested either unabsorbed or following absorption with whole *Klebsiella pneumoniae* bacteria.

Results—Using unabsorbed sera at a dilution of 1:160 the test was found to have a sensitivity of 100%, specificity of 98%, positive predictive value (PPV) of 89% and negative predictive value (NPV) of 100%. There proved to be no advantage in pre-absorbing sera with *K. pneumoniae* antigen.

Conclusions—In the absence of culture methods for *Calymmatobacterium granulomatis*, an indirect immunofluorescence technique may prove valuable for the diagnosis of individual cases of granuloma inguinale and as an epidemiological tool in studies of the disease.

Introduction

At present, establishment of a diagnosis of granuloma inguinale depends on a high degree of clinical suspicion of the disease and the demonstration of characteristic Donovan bodies in smears prepared from deep scrapings or punch biopsies of lesions,¹ or in histological preparations of biopsy material.² Since routine cultivation of the causative bacterium, *Calymmatobacterium granulomatis*, is at present not possible, a serological test for this disease may prove useful, not only for diagnosis, but also as an epidemiological tool in areas where the disease is suspected to occur.

In general, serological tests require the use of a suitable antigen which is normally obtained from cultures of the causative organism. In 1943 organisms resembling Donovan bodies were cultured in the yolk-sacs of eggs of free-range chickens.³ These organisms had previously been obtained by aspiration of material rich in Donovan bodies from unruptured pseudo-buboes. Antigen was subsequently prepared from such cultures and positive serological reactions were obtained in the majority

of cases of granuloma inguinale when the antigen was used as a component of either precipitin or complement fixation tests.^{4 5} Unfortunately cultures of these organisms are no longer available and therefore this suitable source of antigen cannot be used.

During early studies on granuloma inguinale it was noted that an immunological relationship existed between some strains of *C. granulomatis* and *Klebsiella pneumoniae*.⁶ Maddox *et al.* in Papua-New Guinea,⁷ compared the serological reactions obtained when using a limited supply of *C. granulomatis* antigen with those obtained with an antigen prepared from a strain of *K. pneumoniae*. It was concluded that although less sensitive, the klebsiella antigen cross-reacted with antibody to *C. granulomatis* sufficiently to be useful in the serological diagnosis of granuloma inguinale.

During studies on the aetiology of genital ulcer disease undertaken in southern Africa it was noted that large numbers of Donovan bodies were often present in tissue sections prepared from lesions of granuloma inguinale.^{1 8} It therefore seemed logical that the bodies present in these sections could be used as a source of antigen in an indirect immunofluorescence test. In this study, we have attempted to determine the sensitivity and specificity of the technique in an area where the disease is known to occur. In addition, the effect of absorption of sera with *K. pneumoniae* antigens on the sensitivity and specificity of the test was investigated.

Patients and methods

Sera were obtained from 55 cases of microscopically-proven granuloma inguinale seen either at the Leslie Williams' Memorial Hospital, Carletonville, Transvaal, or various hospitals in the Eastern Transvaal and Swaziland. In addition 289 sera were obtained from patients with sexually acquired genital ulcer disease (GUD) other than granuloma inguinale (mainly chancroid) during an ongoing surveillance programme on GUD undertaken at the Leslie Williams' Memorial Hospital, Carletonville.⁹ A further 79 sera were obtained from presumably healthy blood donors on the Witwatersrand (where granuloma inguinale is infrequently seen) and 413 sera were also collected from blood donors in areas of South Africa (Eastern Transvaal and Natal) where clinical cases of granuloma inguinale have frequently been recorded.^{1 8}

Fifty of the sera obtained from patients with proven granuloma inguinale were tested prior

Department of
Anatomical Pathology
A L Freinkel

Department of
Medical Microbiology
Y Dangor
H J Koornhof
R C Ballard

School of Pathology of
the South African
Institute for Medical
Research and
University of the
Witwatersrand
Medical School,
Johannesburg, South
Africa.

Address for
correspondence
Dr A L Freinkel, South
African Institute for
Medical Research, P O
Box 1038, Johannesburg,
2000 Republic of South
Africa

Accepted for publication
19 February 1992

Fig Donovan bodies present in a lesion of granuloma inguinale appear as clusters of fluorescent oval bodies in a positive immunofluorescence test

to absorption with *K. pneumoniae* antigen and all 55 sera were tested following absorption. Of the remaining sera only those found to have significant titres (>1:160) were absorbed and retested.

The indirect immunofluorescence technique was performed by applying suitable dilutions of test serum to protease-treated, dewaxed and rehydrated tissue sections (5 µm) cut from tissue obtained from a lesion clinically and histologically diagnosed as granuloma inguinale, and incubated at 37°C for 30 minutes. The sera were applied to a series of sections in doubling dilutions commencing with a dilution of 1:20. Both positive and negative control sections were used in each batch of sera tested. The sections were then incubated for a further 30 min at 37°C with sheep antihuman IgG conjugated with fluorescein isothiocyanate (Wellcome Diagnostics, Beckenham, Kent) counter-stained with Evans blue, mounted in an aqueous mounting medium, and examined at ×400 magnification by means of a fluorescence microscope. A positive reaction was defined as clusters of bright fluorescing oval structures of about 1.5 µm within vacuoles in large mononuclear cells (fig). The highest dilution of serum which produced typical fluorescence was regarded as the titre of the serum under consideration.

The test was found to be reproducible, with titres obtained following retesting deviating by

a maximum of one dilution. The test was later modified to include absorption of cross reacting antibodies from the sera by the use of a strain of *K. pneumoniae* in a procedure similar to the absorption of nonspecific treponemal antibodies from sera, as described by Wood *et al.*¹⁰ *K. pneumoniae* strain ATCC 13883 was grown in BHI broth overnight at 37°C, subcultured on agar, washed with phosphate buffered saline (pH 7.2) and centrifuged to form firm pellets. Patient's serum (0.5 ml) was added to 0.2 ml of pellet material, mixed, then recentrifuged to form a pellet and the supernatant serum used in the fluorescent antibody test.

The clinical performance of the serological test was evaluated by applying the predictive value model as described by Galen.¹¹

Results

Prior to absorption, all the sera obtained from patients with granuloma inguinale reacted positively at titres > 1:160 (See table 1). But at titres > 1:160, there were eight sera from presumably healthy blood donors and six sera from patients with a sexually transmitted disease other than granuloma inguinale whose sera were found to be reactive.

After absorption with *K. pneumoniae* antigen all except two sera obtained from confirmed cases of granuloma inguinale remained reactive but at lower titres which varied from 1:10 to 1:1280 (table 2). None of the blood transfusion sera from the Witwatersrand were found to be reactive following absorption but 11 sera obtained from disease-endemic areas remained reactive (titres ranged 1:10-1:320) and one serum obtained from an STD patient with proven chancroid was also reactive at a titre of 1:20.

The sensitivity, specificity, positive and negative predictive values and efficiency of the test, with both unabsorbed and absorbed sera and using the GUD sera as negative controls are shown in table 3.

The efficiency of the test was not enhanced as a result of inclusion of the absorption step. Clearly, the results with the sera obtained from blood donors confirm the clinical impression that granuloma inguinale is not endemic on the Witwatersrand but occurs in the eastern Transvaal and Natal.

Table 1 Distribution of immunofluorescence antibody titres to Donovan bodies in sera from patients with granuloma inguinale, other genital ulcer diseases and blood donors (unabsorbed)

Titre	Proven Granuloma inguinale	STD sera	Blood Donor Sera	
			Witwatersrand (non-endemic)	E. Tol/Natal (endemic)
1:5120	3	—	—	1
1:2560	6	—	—	—
1:1280	8	—	—	1
1:640	17	—	—	—
1:320	10	1	—	—
1:160	6	5	—	6
1:80	—	9	1	7
1:40	—	2	4	18
1:20	—	1	9	66
Neg	—	271	65	316
Total	50	289	79	413

Table 2 Distribution of immunofluorescence antibody titres to Donovan bodies in sera from patients with granuloma inguinale, other genital ulcer diseases and blood donors (sera absorbed with *K. pneumoniae* antigen)

Titre	Proven Granuloma inguinale	STD sera	Blood Donor Sera	
			Witwatersrand (non-endemic)	E. Toll/Natal (endemic)
1:1280	1	—	—	—
1:640	5	—	—	—
1:320	5	—	—	1
1:160	7	—	—	1
1:80	8	—	—	1
1:40	3	—	—	1
1:20	21	1	—	3
1:10	3	—	—	4
Neg	2	288	79	402
Total	55	289	79	413

Table 3 Characteristics of the indirect immunofluorescence test for antibody to Donovan bodies

	Sensitivity (%)	Specificity (%)	PPV (%)	NPV(%)	Efficiency (%)
Unabsorbed (>1:160)	100	98	89	100	98
Unabsorbed (>1:320)	88	100	98	—	—
Absorbed	96	100	98	99	—

PPV = Positive predictive value. NPV = Negative predictive value.

Discussion

The results of these preliminary studies clearly indicate that de-waxed tissue sections from proven cases of granuloma inguinale can be successfully used as an antigen in indirect immunofluorescence tests to detect antibody to *C. granulomatis*. In its original form, with no absorption step, the test clearly differentiated between cases of granuloma inguinale and other causes of genital ulcer disease such as chancroid, syphilis, lymphogranuloma venereum and genital herpes which also occur on the subcontinent. No advantage was gained by including a relatively complicated absorption step using *K. pneumoniae* antigens designed to increase the specificity of the test. However, break-point titres had to be employed in unabsorbed tests in order to achieve satisfactory levels of sensitivity and specificity. If all sera exhibiting antibody titres >1:160 were considered positive the sensitivity was found to be 100% and therefore this could be used as an ideal dilution of serum to screen if granuloma inguinale was being excluded as a diagnosis (NPV=100%). Likewise titres of >1:320 were found to be 100% specific for granuloma inguinale with a PPV of 98%. In practice this could be used as the antibody titre which would be considered diagnostic for the disease.

In this study, the calculations of the characteristics of accuracy of the serological test were made using sera obtained from migrant mine-workers with other proven causes of genital ulceration as negative controls. Approximately 60% of these patients usually live in areas where granuloma inguinale is thought to be endemic but the remainder come from areas remote from the “endemic region”. It could be argued that this group of patients may not be an ideal negative control group for the test; however, their inclusion does indicate that antibody to other causes of genital ulceration does not cross-react in the granuloma inguinale antibody test. Ideally a similar group of STD patients drawn exclusively from granulo-

ma inguinale-endemic areas should be used as the negative control group in order to determine more accurately the specificity of the test.

The patterns of reactivity of the blood donor sera obtained from both endemic and non-endemic areas indicate that exposure to *C. granulomatis* or a closely related organism occurs more frequently in granuloma inguinale-endemic areas. The reactivity detected in sera obtained in the endemic area could reflect either non-specific reactivity, detection of latent or undisclosed disease or previously treated infection. Absorption of the sera with *K. pneumoniae* antigen resulted in lowering of titres in individual cases but the fact that sera remained reactive in the modified (absorbed) test indicates that the reactivity is unlikely to be non-specific.

In this study, we were unable to establish a relationship between duration of infection and presence of elevated antibody levels. In early disease serum antibody may not be detectable, which could limit the diagnostic usefulness of the test. Likewise, further studies on the persistence of antibody to *C. granulomatis* following successful therapy for the disease are clearly indicated. Until *C. granulomatis* organisms can be cultivated in pure culture this serological test could be used to screen relatively large numbers of sera in order to determine whether granuloma inguinale constitutes a significant public health problem in a particular area.

By adapting the test using a known positive serum, the indirect immunofluorescence staining technique could also provide an additional, novel and specific method for demonstrating Donovan bodies in tissue sections or smears obtained from cases of the disease.

1 Freinkel AL. Granuloma inguinale (Donovanosis). *Southern African Journal of Sexually Transmitted Diseases* 1984;4:43-7

2 Freinkel AL. Histological aspects of sexually transmitted genital lesions. *Histopathology* 1987;11:819-31.

- 3 Anderson K. The cultivation from granuloma inguinale of a microorganism having the characteristics of Donovan bodies in the yolk sac of chick embryos. *Science* 1943;97: 560-1.
- 4 Anderson K, Goodpasture EW, deMonbreun WA. Immunologic relationship of *Donovania granulomatis* to granuloma inguinale. *J Exp Med* 1945;81:41-51.
- 5 Goldberg J, Weaver RH, Packer H, Simpson WG. Studies on granuloma inguinale II The complement fixation test in the diagnosis of granuloma inguinale. *American Journal of Syphilis, Gonorrhea and Venereal Diseases* 1953;37:71-6.
- 6 Packer H, Goldberg J. Studies of the antigenic relationship of *D. granulomatis* to members of the tribe eschericheae. *American Journal of Syphilis, Gonorrhea and Venereal Diseases* 1950;34:342-50.
- 7 Maddocks I, Anders EM, Dennis E. Donovanosis in Papua New Guinea. *Br J Venereal Dis* 1976;52:190-6.
- 8 Freinkel AL. Granuloma inguinale of cervical lymph nodes simulating tuberculous lymphadenitis: two case reports and review of published reports. *Genitorin Med* 1988;64:339-43.
- 9 Dangor Y, Fehler G, Exposto F da L MPP, Koornhof HJ. Causes and treatment of sexually acquired genital ulceration in southern Africa. *S Afr Med J* 1989;76:339-41.
- 10 Wood RM, Inouye Y, Argonza W, Bradford L, Jue R, Jeong Y, Puffer J, Bodily HL. Comparison of the fluorescent treponemal antibody absorption and *Treponema pallidum* immobilisation tests on serums from 1182 diagnostic problem cases. *Am J Clin Pathol* 1967;47:521-4.
- 11 Galen RS. Use of predictive value theory in clinical immunology. In: Rose NR, Friedman H, Fahey JL, eds. *Manual of Clinical Laboratory Immunology*. 3rd ed. Washington, American Society for Microbiology, 1986: 966-970.