MATTERS
ARISING

Granuloma annulare of the penis

In the February, 1992 issue of *Genitourinary Medicine* Hillman *et al.*¹ described one case of granuloma annulare (GA) affecting the shaft of the penis. The authors stress the great rarity of this dermatological condition involving the penis and could find only one other example after a careful search of the literature. As a practising dermatologist for some 15 years, I recognised GA in half a dozen or so patients; all were female, their ages varying between 18 and 40 years. In every one of these, the lesion occurred on the dorsum of hand or finger. As a venereologist I saw only one case of genital GA about 25 years ago whilst practising in the Wessex region. The patient was a male aged about 50 years who presented with a single, typical GA lesion on the shaft of his penis. It had been present for over six months and he had attended the VD clinic of a teaching hospital in London without the true diagnosis being made. As in the case studied so fully by Hillman *et al.*, this man was also free from venereal disease.

S M LAIRD
3 Beech Cottages, Lydiat Lane,
Alderley Edge, Cheshire, SK9 7HB, UK

1 Hillman RJ, Waldron S, Walker MM, Harris JRW. Granuloma annulare of the penis. *Genitourin Med* 1992;68:47-9.

World-wide distribution of high level tetracycline-resistant *Neisseria gonorrhoeae*.

We read with interest the paper of Van Dyck *et al.*¹ reporting tetracycline-resistant *Neisseria gonorrhoeae* (TRNG) in Zaire. TRNGs have also been reported from The Netherlands,^{2,3} France,⁴ Spain,⁵ and Canada⁶ following the original isolation of TRNG in the USA.⁷ We have had a long standing interest in high level TRNG having been the first group to isolate and report TRNGs in the UK in June 1987.⁸ This predated the report of Ison *et al* in the *Lancet*⁹ by at least 6 months.

Previous work has also shown that the resistance gene which is of the TetM class can be found on one of two conjugative plasmids, the American type and the Dutch type, so called because of the country of origin of the strains in which they were first observed.¹⁰ We have extended this work to examine isolates acquired in Africa, Asia, Europe and South America.

In the United Kingdom we have observed the importation of tetracycline resistant *N gonorrhoeae* (TRNG) from various parts of the globe and have been able to look at these isolates at a molecular level. The table shows a summary of the regions from which we have either obtained isolates or where the original infection was acquired. The table

also includes their plasmid types (either Dutch or American). The plasmid type was determined as described elsewhere¹⁰ using restriction endonuclease digestion of the plasmid DNA with the enzyme *Bgl* I which easily differentiates the two plasmid types.

The strains which originated in Africa appear to be divided into two groups. Isolates originating from the Cameroon and Gabon on the west coast of Africa have resistance plasmids of the Dutch type, whereas isolates from the rest of the African countries which span central and eastern Africa together with the Seychelles Islands possess the American type. A comparison of the Southern blot hybridisation data of Van Dyck *et al* with that produced by ourselves^{10,11} suggests that the plasmid studied from Kinshasa (Zaire) was of the American and not the Dutch type. This extends the observations that the TRNGs we have examined, acquired from Central Africa, were of the American type. All of the Asian strains examined carried the Dutch type of plasmid as did the strains isolated from patients whose infections were acquired in the South American countries of Brazil and Columbia. There does not appear to be a clear pattern of plasmid types among TRNG found in the European countries. Strains from the Netherlands appear to be predominantly of the Dutch type, one strain from Portugal has the American plasmid and the United Kingdom appears to have both types of plasmid although most isolates acquired in this country carry the American type of plasmid. The West Indies and Jamaica have predominantly American type TRNGs. We have been unable to acquire any recent strains originating in the United States and so we have not been able to include any recent American isolates in this study.

The plasmids encoding high level tetracycline resistance appear to follow closely the epidemiology of the penicillinase-producing *N gonorrhoeae* (PPNG). As with the PPNG there appears to be two plasmid types which appeared independently at approximately the same time. Each plasmid type became established in its own geographical region. We have seen this with the TRNGs although limited numbers of TRNG have been obtained from the different countries. This is probably due to poor surveillance of gonococcal isolates for tetracycline resistance. The presence of TRNG in different countries can only be recognised if there is active screening in laboratories for them and the only method of quantifying the spread of these organisms is for reference laboratories to accumulate data on these strains. Many of the isolates which

we have received were referred because they were resistant to penicillin and not because of tetracycline resistance. The limited numbers of isolates obtained for this study are insufficient for meaningful statistical analysis but indicate the occurrence of an interesting distribution of the different plasmid types in the world.

Many of these isolates have been brought to the United Kingdom by travellers, including those from the business community and tourists. The occurrence of TRNG in these countries may only be indicated by their presence in the communities which come into contact with those travellers. However, these communities can also act as a source of TRNG for those countries as a whole. The importation of TRNG into the United Kingdom has led to the occurrence here of both of the two plasmid types. This may well be an indication that sexually active Europeans travel widely and in consequence sample different bacterial populations. In our study, TRNG/PPNG strains from Asia and West Africa appear to have Asian type β -lactamase-encoding plasmids and strains from Europe, North and South America, Central and West Africa (except one strain from Kenya) have the African type β -lactamase-encoding plasmid. The presence of the African or Asian plasmid does not seem to affect the type of tetracycline resistance plasmid present and this reinforces the reported conjugative spread of the plasmid conferring tetracycline resistance.

D M GASCOYNE-BINZI
PM HAWKEY
J HERITAGE
Department of Microbiology,
The Old Medical School,
University of Leeds, Leeds LS2 9JT
A TURNER
Gonococcus Reference Unit,
Public Health Laboratory,
Myrtle Road, Bristol BS2 8EZ
M NADARAJAH
Department of Pathology,
Singapore General Hospital,
Outram Road, Singapore O316

1 Van Dyck E, Rossau R, Duhamel M, *et al.* Antimicrobial susceptibility of *Neisseria gonorrhoeae* in Zaire: high level plasmid-mediated tetracycline resistance in Central Africa. *Genitourin Med* 1992;68:111-6.
2 Roberts MC, Wagenvoort JHT, Van Klinger B, Knapp JS. *TetM* and β -Lactamase—containing *Neisseria gonorrhoeae* (Tetracycline resistant and penicillinase producing) in the Netherlands. *Antimicrob Agents, Chemother* 1988;32:158.
3 Van Klinger B, Dessens-Kroon M, Verheul M. Increased tetracycline resistance in gonococci in the Netherlands. *Lancet* 1989;ii:1278.

The distribution of tetracycline-resistant *Neisseria gonorrhoea* around the world and the associated plasmid types

Geographical area	Country	Number of strains	Plasmid type	PPNG strains
Africa	Cameroon	1	Dutch	1
	Gabon	1	Dutch	1
	Kenya	4	American	4
	Seychelles	1	American	1
	Zaire	1	American	1
Asia	Indonesia	2	Dutch	2
	Singapore	2	Dutch	2
	South East Asia	1	Dutch	0
	Netherlands	12	Dutch	7
Europe	Portugal	1	American	1
	United Kingdom	2	Dutch	2
	United Kingdom	11	American	4
	Jamaica	3	Dutch	3
North America	Jamaica	12	American	10
	United States	2	American	0
	West Indies	1	American	1
South America	Brazil	1	Dutch	1
	Columbia	2	Dutch	1