

Your resource...

Sexually Transmitted Diseases is your resource for the most current, comprehensive information available on all aspects of sexually transmitted disease. Its unique, multidisciplinary perspective provides insight into the latest advances in the field.

■ **Diagnosis and Treatment**


■ **Etiology** ■ **Pathology**

■ **Epidemiology** ■ **Microbiology**

■ **Public Health and Behavior**


Published bi-monthly, *Sexually Transmitted Diseases* brings you original articles on newly identified syndromes, new investigations into the mechanism of disease, descriptions of clinical experiences and reports on therapeutic trials. Features include notes from the CDC, NIH and ASHA, case reports and AVDA news.

Subscribe to the resource you can rely on for the latest, most authoritative information on STDs!



No-Risk Guarantee!

If you're not completely satisfied, you may cancel your subscription at any time and receive a refund for all unmailed issues.



☎ Call us today TOLL FREE
1-800-777-2295 or FAX your
order 1-301-824-7390

Published bi-monthly
Annual subscription rates:
Individual \$135.00 (U.S.) - \$145.00 (Other countries)
Institution \$187.00 (U.S.) - \$225.00 (Other countries)

Published by

J. B. Lippincott


A **Wolters Kluwer** Company



Journals Division
J. B. Lippincott
227 East Washington Square
Philadelphia, PA 19106-3780

for the latest information on *Sexually Transmitted Diseases*

Journal of the American Venereal Disease Association
Julius Schachter, PhD, Editor



On reflection, no other treatment looks so convenient

Many women with genital warts would welcome the convenience and dignity of treating themselves at home.

New Warticon Fem makes self-treatment practical by the simple inclusion of a mirror in every pack.

Proven to be effective¹ and well tolerated² when used in this way,

Warticon Fem is now a clear first choice treatment.

ABBREVIATED PRESCRIBING INFORMATION Warticon, Warticon Fem: Podophyllotoxin 0.5% w/v. **Presentation:** An acidic ethanolic solution of 0.5% w/v podophyllotoxin. **Uses:** For the topical treatment of condyloma acuminata affecting the penis or the female external genitalia. **Dosage and Administration:** The affected area should be thoroughly washed with soap and water, and dried prior to application. Using the applicator provided, the warts should be painted twice daily for 3 days. The treated area should be allowed to dry. If residual warts persist, this 3-day treatment may be repeated, at weekly intervals, if necessary, for a total of 4 weeks of treatment. The majority of patients will not require in excess of 30 loops for each application, however a maximum of 50 loops per application (equivalent to 250µl of Warticon Solution) may be applied. Where lesions are greater in area than 4cm², it is recommended that treatment takes place under the direct supervision of medical staff. **Contra-Indications, Warnings, etc:** Open wounds following surgical procedures should not be treated with podophyllotoxin. Hypersensitivity to podophyllotoxin is a contra-indication. Avoid contact with the eyes. In the event of the preparation entering the eye, the eye should be thoroughly bathed in water. **Side effects:** Local irritation may occur on the second or third day of application associated with the start of wart necrosis. In the majority of cases the reactions are mild. Tenderness, smarting, erythema, superficial epithelial ulceration and balanoposthitis have been reported. Local irritation decreases after treatment. Use in **Pregnancy:** Do not use during pregnancy or lactation. **Overdosage:** There have been no reported overdoses with Warticon Solution. No specific antidote is known. Following accidental spillage, wash the skin well with soap and water. In the event of accidental ingestion give emetic or stomach washout. Treatment should be symptomatic and in severe oral overdose ensure the airway is clear and give fluids, check and correct electrolyte balance, monitor blood gases and liver function. Blood count should be monitored for at least five days. **Pharmaceutical Precautions:** Product should be stored at room temperature. **Legal Category:** POM. **Package Quantities:** Each bottle contains 3ml of Warticon Solution. Plastic applicators are also enclosed in each pack. Each loop will carry a volume of approximately 5µl Warticon Solution. Warticon Fem also contains a mirror to facilitate accurate application. **Basic NHS Cost:** Warticon 3ml £14.50, Warticon Fem 3ml £14.50. **Product Licence Number:** PL3863/0007 **Date of Preparation:** March 1994.


NEW
Warticon
fem
Podophyllotoxin 0.5% w/v

Clearly simple enough for
self-application in female genital warts

 Perstorp Pharma

Further information is available from: Perstorp Pharma Ltd, Intec 2, Wade Road, Basingstoke, Hants RG24 8NE. Tel: 0256 477868.

REFERENCES 1. Baker D. A. et al. *Obstet Gynaecol* 1990; 76: 656-9. 2. Kinghorn G. R. et al. 1993; In press.

 and Warticon are registered trademarks.