Neutrophil enzymes in urine for the detection of urethral infection in men

P A Fraser, J Teasdale, K S Gan, R Eglin, S C Scott, C J N Lacey

Abstract

Objectives—To determine if assaying the neutrophil enzymes, neutrophil elastase (NE) and myeloperoxidase (MPO) in the urine of men attending a genitourinary medicine clinic could identify those with *Neisseria gonorrhoeae* or *Chlamydia trachomatis* infections, and those with urethritis (with or without an identified infection with either organism), and to compare the new assays with the performance of the leucocyte esterase test (LET).

Method—100 men had urethral specimens taken for Gram-stained urethral smear, culture for *N gonorrhoeae*, and for *C trachomatis* testing by enzyme immunoassay. First-voided urines were tested for leucocyte esterase by commercial dipstick (positives were defined as greater than "trace") and then frozen at -20°C prior to being assayed for NE and MPO.

Results—Five patients had gonorrhoea, six had chlamydia and none had both. Evidence of urethritis (>5 polymorphonuclear leucocytes in four $\times 1000$ fields) was found in 29 men. The results of the urine assays showed MPO levels to be non-discriminatory; however NE levels were significantly elevated in patients with proven infection or urethritis or both. Using NE values from men with no infection or urethritis an upper limit for normal was defined. Utilising this, the sensitivity of the elastase assay was calculated and found to be superior to the sensitivity of LET for detecting proven infection (64% vs 36%) and urethritis (52% vs 31%).

Conclusions—Further studies of neutrophil elastase in the pathogenesis, diagnosis and treatment of urethritis are indicated.

(*Genitourin Med* 1995;71:176-179)

Keywords: neutrophil elastase; urethritis

Chlamydia trachomatis and *Neisseria gonorrhoeae* are common sexually transmitted pathogens which may be associated with serious long-term sequelae, particularly in women.¹ Men with genital infections caused by these organisms often develop urethral discharge or dysuria, but asymptomatic carriage is not uncommon and may represent a reser-

voir of infection in the community. Treatment is cheap but diagnosis is expensive and uncomfortable, since it usually involves taking urethral swabs, which many men find unacceptably invasive. The leucocyte esterase test (LET) is a urine dipstick test which detects esterases present in polymorphonuclear leucocytes² (PMNLs) and since men with urethral infection usually have pyuria, LET has been investigated as a cheap, non-invasive screening test for asymptomatic *C trachomatis* and *N gonorrhoeae* infections. Initial studies in adolescent males were promising, showing sensitivities for the LET test in detecting urethral infection of 72-100%.³⁻⁵ However, subsequent studies in adult males have shown sensitivities of 41-70%⁶⁻⁸ with one of the studies showing the test to be more sensitive in younger men. We decided to extend this strategy by investigating the levels of two other neutrophil enzymes in urine and whether these might be better markers of infection or urethritis. We therefore investigated assays of neutrophil elastase (NE) and myeloperoxidase (MPO), two enzymes which are found in the azurophil granules of PMNLs. MPO reacts with H_2O_2 and chloride to produce hypochlorous acid, an oxidant which is thought to be bactericidal,⁹ whilst elastase is a non-specific proteinase, capable of cleaving a wide range of proteins, including human elastin and collagen. NE is a powerful enzyme; its activity is regulated by alpha-1 antitrypsin, a protease inhibitor secreted mainly by the liver. The nature of unregulated neutrophil elastase activity is illustrated by hereditary deficiency of alpha-1 antitrypsin where those affected usually develop pulmonary emphysema.¹⁰ NE is also thought to play a significant role in the lung injury which occurs in the adult respiratory distress syndrome.¹¹

We investigated urine samples of symptomatic and asymptomatic men attending a genitourinary medicine clinic to determine if those with infections could be identified by the new assays. LET was also performed to compare its sensitivity and specificity.

Materials and methods

Any male undergoing full genital screening tests in the Department of Genito-urinary Medicine, whether symptomatic or asymptomatic, was considered eligible for the study. One hundred males (age range 18-68 years) were entered into the study. First voided urine samples were collected into sterile containers

Department of Genitourinary Medicine

P A Fraser
K S Gan
C J N Lacey

Department of Haematological Malignancy Diagnostic Service, Leeds General Infirmary

J Teasdale
S C Scott

Department of Virology, Public Health Laboratory Service, Seacroft Leeds

R Eglin

Address correspondence to:
Dr C J N Lacey,
Department of
Genitourinary Medicine,
Leeds General Infirmary,
Great George Street, Leeds
LS1 3EX.

Accepted for publication
19 January 1995

without preservatives or boric acid and were tested for leucocyte esterase with a commercially available dipstick (Multistix 8 SG; Bayer Diagnostics). Results were recorded on a five-point colour scale (negative, trace, small, moderate and large) with LET positive urines being defined as those with more than a trace reaction. All urines were then frozen at -20°C . Routine sexually transmitted disease (STD) investigations carried out for these patients included Gram staining and microscopy of a urethral smear to document the number of PMNLs and the presence or absence of gonococci, followed by culture on vancomycin-colistin-amphotericin-trimethoprim medium for *N. gonorrhoeae*. A further urethral swab was tested for *C. trachomatis* by EIA (VIDAS—bioMerieux) and blood was taken for syphilis serology.

Urine enzyme assays

The methods used for determining the urinary levels of NE and MPO were optimised in a series of preliminary experiments which determined assay pH, substrate concentration, assay time and assay volume. For MPO, the optimised assay procedure was as follows. Aliquots (50 μ l) of buffered urine samples, prepared by mixing urines with an equal volume of 0.5M phosphate buffer pH 5.0, were

pipetted into microtitre plate wells and pre-warmed to 37°C. O-phenylenediamine substrate solution (100 µl) (4.2 mM substrate dissolved in 50 mM citrate-phosphate buffer pH 5.0 containing 0.3 mM hydrogen peroxide prepared immediately before use) was then added and the absorbance at 492 nm immediately determined (that is, at time 0 hours) in relation to a substrate blank (that is, 100 µl substrate plus 50 µl buffer). The reaction mixtures were then incubated at 37°C in the dark for three hours after which the A₄₉₂ values were measured again against the substrate blank (note that the reactions were not stopped with sulphuric acid, as preliminary studies had demonstrated significant colour changes occurring in some urine samples because of an apparent reaction between the acid and the urine pigments). The change in A₄₉₂ over the three hour period was calculated for each urine and the result expressed as arbitrary units (AU).

For the assay of urinary neutrophil elastase, 50 μ l of each urine sample was mixed in microtitre plate wells with 100 μ l of 0.1M Tris-HCl buffer pH 8.3 containing 0.96M NaCl and the microtitre plates then prewarmed to 37°C. To the buffered urines was then added 50 μ l of working substrate solution and the absorbance at 405 nm immediately determined (time 0) in relation to a substrate blank (that is, 50 μ l substrate plus 150 μ l buffer). The plates were then incubated at 37°C for three hours in the dark and the A_{405} measured again against the substrate blank. The change in A_{405} over the three hour period was calculated for each urine and the result expressed as arbitrary units (AU). The L-pyroglutamyl-L-prolyl-L-valine-p-nitroanilide (Novabiochem; Switzerland) substrate used in this study is highly specific for neutrophil elastase.¹² A stock 10 mM solution was made by dissolving 25 mg substrate in 5.6 ml dimethylsulphoxide and this was stored at 4°C; a working solution was subsequently prepared immediately before use by diluting the stock solution 1:5 with distilled water.

Results

We first looked at two main outcomes: proven infection, defined as a positive culture for *N gonorrhoeae* or a positive EIA test for *C trachomatis*, and evidence of urethritis, defined as > 5 PMNLs seen in oil immersion microscopy of a Gram-stained urethral smear in four fields at $\times 1000$ magnification. In total, 11 patients had a proven infection, 5 with *N gonorrhoeae* and 6 with *C trachomatis*. There were no dual infections. All 5 of those with gonorrhoea, but only 2 of the *C trachomatis* positive group, had urethritis. In addition, 22 men had urethritis but no defined infection making a total of 29 with urethritis altogether. The remaining 67 men had neither an infection nor urethritis.

Urine enzyme activities

Analysis of the urines for NE revealed that the majority of urine enzyme activities were below

Figure 1 Urinary neutrophil elastase activities expressed as arbitrary units (AU).

Figure 2 Urinary myeloperoxidase activities expressed as arbitrary units (AU).

the 0.1 AU level (fig 1). The Mann Whitney test was used to compare the NE urine activities of the men with proven infection with the urine activities of those with no infection or urethritis; NE levels were significantly higher in the infection group ($p = 0.003$). Comparison was also made between the NE activities of the urethritis group and the group with no infection or urethritis; the difference was highly significant ($p < 0.001$).

Analysis of the urines for MPO revealed a wide range of activities (fig 2). The Mann Whitney test showed no significant difference between the urine MPO activities of men with infection, men with urethritis and men with neither infection nor urethritis.

Performance of LET

LET was positive in 10 cases; 4 of 5 of those with gonorrhoea, none of those with *C trachomatis* infection, 5 of 22 of those with urethritis but no proven infection and 1 of 67 of those with no infection and no urethritis. The sensitivity, specificity, positive predictive and negative predictive values of LET for detecting proven infection were 36%, 93%, 40% and 92% respectively. The corresponding values for detecting urethritis were 31%, 98%, 90% and 78% (table 1).

Table 1 Performance of the leucocyte esterase test

	Proven infection*	Urethritis†
Sensitivity	36%	31%
Specificity	93%	98%
Positive predictive value	40%	90%
Negative predictive value	92%	78%

*Defined as a positive culture for *N. gonorrhoeae* or a positive ELISA test for *C. Trachomatis*
†Defined as > 5 PMNLs per $\times 1000$ field seen on oil immersion microscopy of a Gram stained urethral smear.

Table 2 Performance of the elastase assay

	Proven infection	Urethritis
Sensitivity	64%	52%
Specificity	90%	97%
Positive Predictive Value	44%	88%
Negative Predictive Value	95%	83%

Performance of the elastase assay

We took the whole group of elastase values for the patients with neither infection nor urethritis and using ranks calculated the 95th centile value. This was calculated as 0.195 AU and was defined as the upper limit of the normal range.¹³ We then calculated sensitivity, specificity, positive and negative predictive values for patients with proven infection or urethritis (table 2). The sensitivity of the elastase assay was considerably superior to the LET for detecting proven infection (64 vs 36%) and urethritis (52 vs 31%).

We then looked at the sub-group of men in the study population who were asymptomatic (that is, they did not complain of urethral discharge or dysuria). In total 72 men were asymptomatic. Of these, 2 were found to have chlamydia infections but no urethritis and 7 had evidence of urethritis but no identified infection. LET was negative in the 2 chlamydia cases but positive in 1 of the 7 urethritis cases and in one additional man who had no infection or urethritis.

The elastase assay was positive in 1 of the chlamydia cases, 2 of the urethritis cases and in an additional 3 men with no infection or urethritis. The sensitivity of LET for detecting urethritis or infection in this subgroup was 11% compared to 33% for the NE assay, with specificities, positive predictive values and negative predictive values being very similar for the two tests. However, the small numbers of men with infection/urethritis means that these figures should be interpreted with caution.

We determined whether the elastase values might be correlated with age of the study subjects. Forty four men were < 25 years and 56 men > 25 years of age. There was no statistical difference in elastase values between the two groups (Mann Whitney). This was also the case when this analysis was limited to the 29 men with urethritis. Numbers of men with proven infection were too small to make such an analysis valid.

Discussion

Cheap and simple screening tests for bacterial STDs are urgently needed, particularly in developing countries where STDs can

increase the risk of acquiring or transmitting HIV three to five fold.¹⁴ The performance of LET as a screening test in males has been variable, with good results being reported in adolescents, but other authors have described lower sensitivities in adults,⁶⁻⁸ as we have found in this study. However, the concept of a urine test for neutrophil enzymes which could be used to screen a population is an appealing one and prompted us to evaluate MPO and NE as possible candidates.

We have shown in our pilot study that a population of symptomatic and asymptomatic men had a wide range of urinary MPO levels without any discriminative value. In contrast, urinary NE values were significantly elevated in patients with a proven infection and/or urethritis, compared to those with neither. By using the elastase values of the men without infection or urethritis to define a normal range we have shown that the elastase assay was considerably better than LET in sensitivity. We do, however, accept that these data may be influenced by our use of EIA alone for defining chlamydial infection (which the manufacturers state has an 80% sensitivity with a 98% specificity for male urethral specimens). Further studies to more accurately determine the sensitivity of urinary NE assays should ideally incorporate immunofluorescence, cell culture or PCR for *Chlamydia trachomatis* and PCR for *Mycoplasma genitalium*.¹⁵

The divergent behaviour of myeloperoxidase and elastase in their association with urethritis is worthy of further comment. Both enzymes are derived from the primary azurophil granules of neutrophils. Myeloperoxidase though is capable of directly reacting with superoxide and neutralising free radicals generated during the inflammatory response. Consequently MPO levels within urethral secretions might therefore be "self-regulating" and not show disease association. Elastase, however, is clearly capable of inducing endothelial injury, especially in the presence of free radicals or neutrophils which are already activated.¹¹ This present study is the first to report significantly increased neutrophil elastase activity in material derived

from men with urethritis, and it is tempting to speculate that urinary NE could be directly involved in the pathogenesis. Further studies should be directed towards evaluating its use in the diagnosis of urethritis. One could also postulate that inhibitors of neutrophil chemotaxis¹⁶ or elastase might have activity in therapy of pathogen-negative chronic urethritis.

We are grateful to Dr M A Waugh, Dr E F Monteiro, and Dr J Wilson for allowing us to study their patients. We also thank Vanessa Dixon for secretarial support.

- Weström LV, Mårdh PA. Acute pelvic inflammatory disease. In: KK Holmes, P-A Mardh, PF Sparling, PJ Weisner eds, *Sexually Transmitted Diseases*, 2nd ed. New York, McGraw-Hill, 1990. pp. 593-613.
- Kusumi KR, Grover PJ, Kunin CM. Rapid detection of pyuria by leucocyte esterase activity. *JAMA* 1981; 245:1653-5.
- Sadof MD, Woods ER, Emans SJ. Dipstick leucocyte esterase activity in first-catch urine specimens. *JAMA* 1987;258:1932-4.
- O'Brien SF, Bell TA, Farrow JA. Use of a leucocyte esterase dipstick to detect *Chlamydia trachomatis* and *Neisseria gonorrhoeae* urethritis in asymptomatic adolescent male detainees. *Am J Public Health* 1988; 78:1583-4.
- Shafer MA, Schachter J, Moscicki AB, et al. Urinary leucocyte esterase screening test for asymptomatic chlamydial and gonococcal infections in males. *JAMA* 1989; 262:2562-6.
- McNagny SE, Parker MR, Zenilman MJ, Lewis JS. Urinary leucocyte esterase test: a screening method for the detection of asymptomatic chlamydial and gonococcal infections in men. *J Infect Dis* 1992;165:573-6.
- White DJ, Malet RM, Bignell CJ. Non-invasive sampling method for detecting *Chlamydia trachomatis*. *Lancet* 1989;i:96-7.
- Patrick DM, Rekart ML, Knowles L. Unsatisfactory performance of the leukocyte esterase test of first voided urine for rapid diagnosis of urethritis. *Genitourin Med* 1994;70:187-90.
- Smolen JE, Boxer LA. Functions of neutrophils. In: WJ Williams, E Beutler, AJ Erslev, MA Lichtman eds, *Haematology*, 4th ed, New York, McGraw-Hill, 1992. pp. 780-94.
- Segal AW, Walport MJ. Neutrophil Leucocytes. In: J O'D McGee, PG Isaacson, NA Wright eds, *Oxford Textbook of Pathology, Volume 1*. Oxford, Oxford University Press, 1992. pp. 321-9.
- Repine JE. Scientific perspectives on adult respiratory distress syndrome. *Lancet* 1992;339:466-9.
- Kramps JA, Van Twisk CH, Van Der Linden AC. L-Pyroglutamyl-L-prolyl-L-valine-p-nitroanilide, a highly specific substrate for granulocyte elastase. *Scand J Clin Lab Invest* 1983;43:427-32.
- Altman DG. *Practical Statistics for Medical Research*. London, Chapman and Hall, 1990. pp. 19-47.
- Berkley S. Diagnostic tests for sexually transmitted diseases: a challenge. *Lancet* 1994;343:685-6.
- Horner PJ, Gilroy CB, Thomas BJ, Naidoo ROM, Taylor-Robinson D. Association of *Mycoplasma genitalium* with acute non-gonococcal urethritis. *Lancet* 1993; 342:582-5.
- Lomas DA, Natin D, Stockley RA, Shahmanesh M. Chemotactic activity of urethral secretions in men with urethritis and the effect of treatment. *J Infect Dis* 1993;167:233-6.