

# Health care: the way ahead

## Controversies in Health Care Policies: Challenges to Practice

Edited by Marshall Marinker

Can health policy ever be completely right?

This book explores how to get it right in six of today's most important issues: rational prescribing; primary care following Tomlinson; public access to medical information; the meaning of aging; medical practice variation; and risk in medicine.



EDITED BY  
**MARSHALL MARINKER**

BMJ  
Publishing Group

ISBN 0 7279 0894 4 144 pages

November 1994

UK £12.95; Overseas £15.00

(BMA members £11.95; £14.00)


## Outcomes into Clinical Practice

Edited by Tony Delamothe

Why do outcomes research?  
How should it be done?

These crucial questions are addressed in this ground breaking book, based on the conference held by the BMJ, BMA, and UK Clearing House for Information on the Assessment of Health Outcomes. In it, experts in the field cover a wide range of topics including acute care; cost effectiveness; death rates; health gains; and general practice.

- Essential reading for all involved in making and implementing health care policies.
- Documents the work of six think tanks of clinical experts, GPs, managers, and patient representatives
- Includes input from the Kings Fund, NAHAT, Age Concern, RCGP, National Medical Adviser Support Centre, Social Market Foundation
- Offers innovative and realistic ideas for the way forward
- Shows how information on outcomes can be used to improve care
- Gives examples of outcomes research at its best
- Looks at the subject from all perspectives in the health service
- Provides an ideal starting point for all members of clinical teams, both in hospitals and general practice, and is invaluable for health care purchasers and clinical audit officers.



ISBN 0 7279 0888 X 184 pages

November 1994

UK £17.95; Overseas £20.00

(BMA members £16.95; £19.00)

Also available  
**A Manual for Assessing Health Practices and Designing Practice Policies: the explicit approach**  
By David M Eddy

Published by the American College of Physicians

ISBN 0 943126 18 5 100 pages 1992 UK £19.95; Overseas £21.00 (BMA members £17.95; £19.00)

## ORDER FORM

Available from: BMJ Publishing Group, P.O. Box 295, London WC1H 9TE (Tel: 0171 383 6185/6245),  
medical booksellers or the BMJ bookshop in BMA House

Please send me the following books

\_\_\_\_\_

Name \_\_\_\_\_

(Print Clearly)

Address \_\_\_\_\_

Postcode \_\_\_\_\_

BMA Membership No. \_\_\_\_\_

Cheque enclosed (made payable to British Medical Journal) £ \_\_\_\_\_  
Debit my AMERICAN EXPRESS VISA MASTERCARD \_\_\_\_\_

Card No. \_\_\_\_\_ Exp. \_\_\_\_\_

Signature \_\_\_\_\_

Please send me a BMJ PUBLISHING GROUP CATALOGUE

**BMJ**  
Publishing  
Group

“Something for the next five years sir?”



SmithKline Beecham

VACCINES



SmithKline Beecham Pharmaceuticals  
Welwyn Garden City, Hertfordshire AL7 11Y.

**Presentation:** Each 1 ml of 'Engerix B' hepatitis B vaccine (rby), PL 10592/0015, contains 20 micrograms of hepatitis B surface antigen protein, together with thiomersal 1:20,000. Pack of 1 (1 ml) prefilled syringe containing 20 micrograms, £12.13; pack of 10 (1 ml) prefilled syringes each containing 20 micrograms, £121.30; pack of 1 (1 ml) vial containing 20 micrograms, £11.95; pack of 3 (1 ml) vials each containing 20 micrograms, £35.85; pack of 10 (1 ml) vials, £119.50; pack of 1 paediatric (0.5 ml) vial containing 10 micrograms, £8.96.

**Uses:** Active immunisation against infections caused by hepatitis B virus.

**Dosage and administration:** For intramuscular use only. Shake well and inspect before use. Three doses should be given, the

second one month and the third six months after the initial dose. For more rapid immunisation the third dose can be given two months after the initial dose with a booster at 12 months.

**Adults and children over 12 years:** 20 micrograms (1 ml) given intramuscularly.

**Neonates and children 12 years and under:** 10 microgram (0.5 ml) given intramuscularly.

Administer in the deltoid region, though the antero-lateral thigh is the preferred site for infants. 'Engerix B' should not be administered in the buttock since this may result in lower immune response. In neonates of HBsAg positive mothers, give hepatitis B immunoglobulin at the same time as vaccine at different sites within a few hours of birth.

'Engerix B' isn't just something for the weekend. It provides up to five years' protection against hepatitis B; which means you don't have to rely on your patients using a condom every time they have sex.

So who's at risk? People who are sexually active, either with multiple partners, or who travel abroad and have casual, unprotected sex.

The fact is hepatitis B can be contracted in the same way as AIDS, but it's 100 times more infectious. Worst still, it has been found in body fluids such as sweat, saliva, even tears.

It's quite reassuring to know then that, world-wide, 'Engerix B' has protected more people against the hepatitis B virus than any other vaccine.

You can order 'Engerix B' in pre-filled syringes, by calling SmithKline Beecham on 0181-913 4290. So, even if you can't prevent your patients from picking up every sexually transmitted disease, you can give them five years' protection against hepatitis B.



**Contra-indications:** Hypersensitivity to any component of the vaccine. Severe febrile infections.

**Precautions:** Response may be impaired in renal dialysis patients or those who are immunocompromised. Adrenaline 1:1000 should be available in case of anaphylaxis. Use in pregnancy: see Data Sheet.

**Adverse reactions:** Mild transient local soreness, erythema and duration at the injection site. Occasionally low grade fever, malaise, fatigue, arthralgia, arthritis, myalgia, headache, dizziness, syncope, nausea, vomiting, diarrhoea, abdominal pain, lymphadenopathy, abnormal liver function tests, rashes rarely with urticaria. Exceptionally, severe skin disorders such as erythema multiforme. Very rarely one week or more after

injection, transient arthralgia, pruritus or urticaria, but no causal relationship established.

Neurological manifestations in temporal association with the vaccine, including very rarely paraesthesia and extremely rarely paralysis, neuropathy and neuritis (including Guillain-Barré syndrome, optic neuritis and multiple sclerosis). No causal relationship established.

Early onset allergic-type reactions reported rarely.

**Legal category** POM. 11.8.94.

'Engerix B' is a trade mark.

© 1995 SmithKline Beecham Pharmaceuticals

**ENGRIX B**  
**Hepatitis B Vaccine (rby)**


0695 EB:AD/5/051

# How to do it even better!

The highly popular *How To Do It* series is a collection of expert advice on an enormous range of topics that doctors need to know about that are not necessarily covered in formal training.

All three volumes have now been completely updated, with new chapters, and a user friendly format where subjects are grouped into themes. Whether you want to chair a committee, improve your counselling skills with cancer patients, or even survive a formal dinner, your questions will be answered in these entertaining and practical guides.


A must for all doctors, from students through to the retired.



### How to Do It Volume 1

- Management ● Employment
- Counselling


ISBN 0 7279 0848 0 272 pages April 1995  
UK £12.95; Overseas £15.00  
(BMA members £11.95; £14.00)



### How to Do It Volume 2

- Examinations ● Academia
- Research and Communication

ISBN 0 7279 0895 2 240 pages April 1995  
UK £12.95; Overseas £15.00  
(BMA members £11.95; £14.00)



### How to Do It Volume 3

- Writing ● Publishing and Publicity
- Something Different

ISBN 0 7279 0896 0 264 pages April 1995  
UK £12.95; Overseas £15.00  
(BMA members £11.95; £14.00)

## Purchase the complete set at a reduced price!

ISBN 0 7279 0906 1 UK £34.95; Overseas £39.00 (BMA members £31.95; £36.00)


**BMJ**  
Publishing  
Group

**Also available: How to Write a Paper** Edited by G M Hall

"Highly recommended as a practical and readable guide to getting published."

*Journal of the Institute of Health Education*

ISBN 0 7279 0822 7 128 pages 1994 UK £8.95; Overseas £11.00 (BMA members £8.45; £10.50)



### Order Form

Available from: **BMJ Publishing Group, P.O. Box 295, London WC1H 9TE, (Tel: 0171 383 6185/6245), medical booksellers or the BMJ bookshop in BMA House**

Please send me \_\_\_\_\_ copy/ies of HOW TO DO IT VOLUME 1

Please send me \_\_\_\_\_ copy/ies of HOW TO DO IT VOLUME 2

Please send me \_\_\_\_\_ copy/ies of HOW TO DO IT VOLUME 3

Please send me \_\_\_\_\_ COMPLETE SET/S OF HOW TO DO IT

Please send me \_\_\_\_\_ copy/ies of HOW TO WRITE A PAPER

BMA Membership No. \_\_\_\_\_

Name \_\_\_\_\_  
(Print Clearly)

Address \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_ Postcode \_\_\_\_\_

Cheque enclosed (made payable to British Medical Journal) £ \_\_\_\_\_

Debit my AMERICAN EXPRESS/VISA/MASTERCARD

Card No. \_\_\_\_\_ Exp \_\_\_\_\_

Signature \_\_\_\_\_

Please send me a BMJ PUBLISHING GROUP CATALOGUE

# NEW WARTICON CREAM



## La Crème de la Crème

New Warticon Cream is the first ever podophyllotoxin cream for the self-treatment of genital warts.

The cream combines the well proven efficacy<sup>1</sup> and good tolerability<sup>2</sup> of podophyllotoxin in an easy to use formulation.

Men and women will find that new Warticon Cream, with its discreet tube and simple application, adds convenience and dignity to the self-treatment of genital warts.

**NEW**  
**Warticon**<sup>®</sup>  
*cream*

Podophyllotoxin 0.15% w/w

### The cream of self-treatment for genital warts


#### Abbreviated Prescribing Information

**Warticon Cream Presentation:** White homogeneous cream containing 0.15% w/w podophyllotoxin.  
**Uses:** For the topical treatment of condyloma acuminata affecting the penis, and the female external genitalia. **Dosage and Administration:** The affected area should be thoroughly washed with soap and water, and dried prior to application. Using a fingertip, Warticon Cream is applied twice daily for 3 days using only enough cream to just cover each wart. The hands should be thoroughly washed after each application. Residual warts should be treated with further courses of twice daily applications for 3 days at weekly intervals, if necessary, for a total of 4 weeks of treatment. Where lesions are greater in area than 4cm<sup>2</sup> it is recommended that treatment takes place under the direct supervision of medical staff. **Contraindications, Warnings etc:** Open wounds, hypersensitivity to podophyllotoxin. Avoid contact with the eyes. In the event of the preparation entering the eye, the eye should be thoroughly bathed in water. Prolonged contact with healthy skin should be avoided, as the cream contains an active pharmaceutical substance that could be harmful to healthy skin. **Side Effects:** Local irritation

may occur on second or third day of application associated with the start of wart necrosis. In the majority of cases the reactions are mild (see Data Sheet). **Use in Pregnancy:** Do not use during pregnancy or lactation. **Overdosage:** There have been no reported overdosages with Warticon Cream. No specific antidote is known. In the event of accidental ingestion give emetic or stomach washout. Treatment should be symptomatic and in severe oral overdose ensure the airway is clear and give fluids, check and correct electrolyte balance, monitor blood gases and liver function. Blood count should be monitored for at least five days. **Pharmaceutical Precautions:** Product should be stored at room temperature. **Legal Category:** POM. **Package Quantities:** Single tube containing 5g of Warticon Cream. The pack also contains a mirror to facilitate accurate application. **Basic NHS Price:** Warticon Cream 5g £17.40. **Product Licence Number:** PL 3863/0010. **Date of Preparation:** March 1995. **References:** 1. Kinghorn, G. et al. International Journal of STD & AIDS 1993; 4: 194-199. 2. Strand, A. et al. 1995 In press

 Perstorp Pharma

Further information is available from: Perstorp Pharma Ltd, Intec 2, Wade Road, Basingstoke, Hants RG24 8NE. Tel: 01256 477868. Fax: 01256 215

 and Warticon are registered trademarks