


Fuzzy about Logic? You should read...



LOGIC IN MEDICINE

2ND EDITION

Edited by Calbert I Philips

Logic in Medicine brings together a team of eminent doctors, scientists and philosophers to explain the process behind clinical diagnoses and management decisions.

This new revised edition features:

- An overview of the general philosophy of logic in medicine, in particular, diagnostic logic
- Clarification of difficult subjects such as Bayes's Theorem of pattern recognition
- New chapters on current major issues including: Clinical Decision Making, Statistics, Medical Law, Sociology and Medical Ethics

ISBN 0 7279 0854 5 232 pages September 1995
UK £14.95 Overseas £17.00 (BMA Members £13.95; £16.00)

Order your copy now

from BMJ Publishing Group, PO Box 295,
London WC1H 9TE or phone our credit card hotline
on 0171 383 6185/6245 (Fax: 0171 383 6662)

BMJ
Publishing
Group

MEMBERSHIP OF THE FACULTY OF FAMILY PLANNING AND REPRODUCTIVE HEALTH CARE

The MFFP examination consists of:—

- ◆ **Part 1** Multiple choice question paper (MCQ)

Dissertation

- ◆ **Part 2** (Entry to part two depends on having held the DFFP for at least two years).

There are three components:

- Modified Essay Question paper (MEQ)
- Critical Reading Question paper (CRQ)
- Objective Structured Clinical examination (OSCE)


- ◆ The Faculty Board may allow those whose degrees do not qualify them for full Registration with the GMC to enter Part 2.
- ◆ Closing dates for application for entry to Part 1 are **1st January** and **1st July**.
- ◆ The qualification is subject to re-certification every five years.

Regulations available on application to:—

The Examinations Secretary.

Faculty of Family Planning and Reproductive Health Care of
the Royal College of Obstetricians and Gynaecologists,
27 Sussex Place, Regent's Park, London NW1 4RG, UK
Tel. 0171 724 2441

Applying science to health care



ISBN 0 7279 1029 9 200 pages April 1996
UK £22.95; Overseas £25.00
(BMA members £20.95; £23.00)

Evidence based medicine is the buzz word in health care today but the concept that the design and function of health services should also be based on scientific evidence is less familiar and more radical. Grown out of a ground breaking conference, *The Scientific Basis of Health Services* examines how the activities of health services can be rooted in research.

Chapters include:

- Economic evaluation and clinical practice
- The role of the consumer in health research
- Evidence based management in health care
- Shifting the balance between primary and secondary care

With contributions from leading international figures, this unique book points the way to the future and is essential reading for all involved in the design and management of health services.

Available from: BMJ Publishing Group, PO Box 295, London WC1H 9TE,
medical booksellers or the BMJ bookshop in BMA House.
OR: Phone on our credit card hotline: 0171 383 6185/6245 or fax: 0171 383 6662
PRINT CLEARLY Please send me.....copy/ies of Scientific Basis of Health Services

Name

Address

Postcode.....Membership No:

☐ Cheque enclosed (made payable to British Medical Journal) £.....

☐ Debit my AMERICAN EXPRESS VISA MASTERCARD

Card No Expires

Signature

BMJ
Publishing
Group

Order your copy today

"Something for the next five years sir?"

SmithKline Beecham
VACCINES

SmithKline Beecham Pharmaceuticals
Welwyn Garden City, Hertfordshire AL7 1EY.

Presentation: Each 1 ml of 'Engerix B' hepatitis B vaccine (rby), P1 10592/0015, contains 20 micrograms of hepatitis B surface antigen protein, together with thiomersal 1:20,000. Pack of 1 (1 ml) prefilled syringe containing 20 micrograms, £12.74; pack of 10 (1 ml) prefilled syringes each containing 20 micrograms, £127.40; pack of 1 (1 ml) vial containing 20 micrograms, £12.55; pack of 3 (1 ml) vials each containing 20 micrograms, £37.65; pack of 10 (1 ml) vials, £125.50; pack of 1 paediatric (0.5 ml) vial containing 10 micrograms, £9.41.

Uses: Active immunisation against infections caused by hepatitis B virus.

Dosage and administration: For intramuscular use only. Shake well and inspect before use. Three doses should be given, the second one month and the third six months after the initial dose. For more rapid immunisation the third dose can be

given two months after the initial dose with a booster at 12 months.

Adults and children over 12 years: 20 micrograms (1 ml) given intramuscularly. *Neonates and children 12 years and under:* 10 micrograms (0.5 ml) given intramuscularly.

Administer in the deltoid region, though the antero-lateral thigh is the preferred site for infants. 'Engerix B' should not be administered in the buttock since this may result a lower immune response. In neonates of HBsAg positive mothers, give hepatitis B immunoglobulin at the same time as vaccine at different sites within a few hours of birth.

Contra-indications: Hypersensitivity to any component of the vaccine. Severe febrile infections.

'Engerix B' isn't just something for the weekend. It provides up to five years' protection against hepatitis B; which means you don't have to rely on your patients using a condom every time they have sex.

So who's at risk? People who are sexually active, either with multiple partners, or who travel abroad and have casual, unprotected sex.

The fact is hepatitis B can be contracted in the same way as AIDS, but it's 100 times more infectious. Worst still, it has been found in body fluids such as sweat, saliva, even tears.

It's quite reassuring to know then that, world-wide, 'Engerix B' has protected more people against the hepatitis B virus than any other vaccine.

You can order 'Engerix B' in pre-filled syringes, by calling SmithKline Beecham on 0181-913 4290. So, even if you can't prevent your patients from picking up every sexually transmitted disease, you can give them five years' protection against hepatitis B.

Precautions: Response may be impaired in renal dialysis patients or those who are immunocompromised. Adrenaline 1:1000 should be available in case of anaphylaxis. Use in pregnancy: see Data Sheet.

Adverse reactions: Mild transient local soreness, erythema and induration at the injection site. Local swelling at the injection site and angioedema have been reported rarely. Occasionally low grade fever, malaise, fatigue, arthralgia, arthritis, myalgia, headache, dizziness, syncope, nausea, vomiting, diarrhoea, abdominal pain, lymphadenopathy, abnormal liver function tests, rashes rarely with urticaria. Thrombocytopenic purpura, severe skin disorders such as erythema multiforme have exceptionally occurred. Very rarely one week or more after injection, transient arthralgia, pruritus or urticaria.

Neurological manifestations in temporal association with the vaccine, very rarely include paraesthesia and extremely rarely include paralysis, neuropathy and neuritis (including Guillain-Barré syndrome, optic neuritis and multiple sclerosis). No causal relationship established. Early onset allergic-type reactions reported rarely.

Legal category POM.
3.796.

'Engerix B' and the SB logo are trade marks.
© 1996 SmithKline Beecham Pharmaceuticals.

ENGRIXB
Hepatitis B Vaccine (rby)

0696 EB:AD/6/541



How do you...

Persuade a government to ban tobacco advertising?




Campaign against dumping of toxic waste?

Energise the fight against AIDS?

In *The Fight for Public Health*, Simon Chapman and Deborah Lupton show how lobbying through media advocacy can be a powerful tool in mobilising public support and getting public health legislation passed. Illustrated with photographs and crammed with compelling examples from around the world, this lively and entertaining guide:

- Describes what advocacy is
- Shows how public opinion has influenced public health policy in the past
- Includes an A-Z of strategies for gaining media attention.



"This splendid book moves the art of media advocacy a giant step forward. Chapman and Lupton's counsel leapfrogs continents; it will be as welcome and applicable in London and Washington as in Sydney."

Michael Pertschuk, Director, The Advocacy Institute, Washington, DC

ISBN 0 7279 0849 9
200 Pages October 1994
UK £19.95; Overseas £22.00
(BMA members £18.95; £21.00)

BMJ
Publishing Group

ORDER FORM

Available from: BMJ Publishing Group, P.O. Box 295, London WC1H 9TE (Tel: 071 383 6185/6245), medical booksellers or the BMJ bookshop in BMA House

Please send me _____ copy/ies of
THE FIGHT FOR PUBLIC HEALTH

Name _____
(Print Clearly)

Address _____

Postcode _____

BMA Membership No. _____

Cheque enclosed (made payable to British Medical Journal) £ _____
Debit my AMERICAN EXPRESS VISA MASTERCARD

Card No. _____ Exp. _____

Signature _____


☐ Please send me a BMJ PUBLISHING GROUP CATALOGUE

In the USA contact American College of Physicians, tel: 800 523 1546; in Australia contact AMA Services (WA) PTY Ltd, Tel: 389 8288

INDISPENSABLE **NEW** GUIDES FOR ALL HEALTH PROFESSIONALS

The Pocket Guide to Critical Appraisal

Iain K Crombie



This practical handbook is designed to accompany researchers on every library visit, taking them step by step through the process of appraisal.

- Introduces the rationale behind critical appraisal
- Tackles the five main methods of medical research in detail: surveys, clinical trials, cohort studies, case control studies, review papers
- Provides a comprehensive one page check list for each method to aid the reader in assessing the quality and relevance of a paper
- Identifies the common pitfalls in publishing research


Written in a lively, jargon-free style, this accessible book arms the reader with all the knowledge needed to make sense of the overwhelming mass of medical literature.

ISBN 0 7279 1099 X 72 pages 1996

UK £8.95; Overseas £11.00 (BMA members £8.45; £10.50)

Elementary Economic Evaluation in Health Care

Tom Jefferson, Vittorio Demicheli, and Miranda Mugford



Doctors, nurses and managers are increasingly responsible for delivering optimum care in the most efficient way, and must have an understanding of economic evaluation and how to practice it. *Elementary Economic Evaluation in Health Care* provides a basic introduction to the subject. Written in lay language and with case histories to illustrate the different processes, it explains the principles and how to use them. Chapters include:

- The basis of economic evaluation
- Cost of illness studies
- Cost minimisation analysis
- Cost effectiveness analysis

Including helpful checklists, flow diagrams and appendices to aid understanding, this is a comprehensive guide to getting started in economic evaluation.

ISBN 0 7279 1074 4 136 pages 1996

UK £14.95; Overseas £17.00 (BMA members £13.95; £16.00)

**ORDER
FORM**

BMJ Publishing Group, PO Box 295, London, England WC1H 9TE
Tel: 0171 383 6185/6245 Fax: 0171 383 6662

QTY

TITLE

AMOUNT

THE POCKET GUIDE TO CRITICAL APPRAISAL

ELEMENTARY ECONOMIC EVALUATION IN HEALTH CARE

TOTAL £

Membership No. where applicable

Please send me a BMJ Publishing Group catalogue

NAME (print name)

ADDRESS

Cheque enclosed (made payable to British Medical Journal) £

Debit my AMERICAN EXPRESS VISA MASTERCARD

Card No.

Expires

Signature

BMJ
Publishing
Group

WARTICON CREAM



La Crème de la Crème

Warticon Cream is the first ever podophyllotoxin cream for the self-treatment of genital warts.

The cream combines the well proven efficacy¹ and good tolerability² of podophyllotoxin in an easy to use formulation.

Men and women will find that Warticon Cream, with its discreet tube and simple application, adds convenience and dignity to the self-treatment of genital warts.

Warticon[®]
cream

Podophyllotoxin 0.15% w/w

The cream of self-treatment for genital warts

Abbreviated Prescribing Information


Warticon Cream Presentation: White homogeneous cream containing 0.15% w/w podophyllotoxin. **Uses:** For the topical treatment of condyloma acuminata affecting the penis, and the female external genitalia. **Dosage and Administration:** The affected area should be thoroughly washed with soap and water, and dried prior to application. Using a fingertip, Warticon Cream is applied twice daily for 3 days using only enough cream to just cover each wart. The hands should be thoroughly washed after each application. Residual warts should be treated with further courses of twice daily applications for 3 days at weekly intervals, if necessary, for a total of 4 weeks of treatment. Where lesions are greater in area than 4cm² it is recommended that treatment takes place under the direct supervision of medical staff. **Contraindications, Warnings etc:** Open wounds, hypersensitivity to podophyllotoxin. Avoid contact with the eyes. In the event of the preparation entering the eye, the eye should be thoroughly bathed in water. Prolonged contact with healthy skin should be avoided, as the cream contains an active pharmaceutical substance that could be harmful to healthy skin. **Side Effects:** Local irritation

may occur on second or third day of application associated with the start of wart necrosis. In the majority of cases the reactions are mild (see Data Sheet). **Use in Pregnancy:** Do not use during pregnancy or lactation. **Overdosage:** There have been no reported overdoses with Warticon Cream. No specific antidote is known. In the event of accidental ingestion give emetic or stomach washout. Treatment should be symptomatic and in severe oral overdose ensure the airway is clear and give fluids, check and correct electrolyte balance, monitor blood gases and liver function. Blood count should be monitored for at least five days. **Pharmaceutical Precautions:** Product should be stored at room temperature. **Legal Category:** POM. **Package Quantities:** Single tube containing 5g of Warticon Cream. The pack also contains a mirror to facilitate accurate application. **Basic NHS Price:** Warticon Cream 5g £17.40. **Product Licence Number:** PL 3863/0010. **Date of Preparation:** March 1995. **References:** 1. Kinghorn, G. et al. International Journal of STD & AIDS 1993; 4: 194-199. 2. Strand, A. et al. Genitourinary Medicine 1995, Vol.71, No.6, P387-390.



Perstorp Pharma

Further information is available from: Perstorp Pharma, Intec 2, Wade Road, Basingstoke, Hants RG24 8NE. Tel: 01256 477868. Fax: 01256 21508.

 and Warticon are registered trademarks