

Original
article

Assessing the impact of national anti-HIV sexual health campaigns: trends in the transmission of HIV and other sexually transmitted infections in England

Angus Nicoll, Gwenda Hughes, Mary Donnelly, Shona Livingstone, Daniela De Angelis, Kevin Fenton, Barry Evans, O Noël Gill, Mike Catchpole

Objective: To assess the impact of the sexual component of AIDS and HIV campaigns on transmission of HIV and other sexually transmitted infections (STIs).

Design: Comparison of time series data.

Setting: England, 1971–1999.

Outcome measures: HIV transmission and diagnoses among men who have sex with men (MSMs), rates of attendances and specific STI diagnoses (per 100 000 total population) at genitourinary medicine (GUM) clinics.

Results: Awareness of AIDS and campaigns in 1983–4 among homosexual men coincided with substantial declines in transmission of HIV and diagnoses of syphilis among MSMs. During general population campaigns in 1986–7 new GUM clinic attendances requiring treatment fell by 117/10⁵ in men and 42/10⁵ in women. Rates for gonorrhoea fell by 81/10⁵ and 43/10⁵ and genital herpes by 6/10⁵ and 4/10⁵, respectively. Previous rises in genital wart rates were interrupted, while rates of attendances not requiring treatment (the “worried well”) increased by 47/10⁵ and 58/10⁵ for men and women, respectively. Since 1987 diagnoses of HIV among MSMs have not declined, averaging 1300–1400 annually. Following a period of unchanging rates there have been substantial increases in GUM attendances requiring treatment, notably for gonorrhoea, syphilis, and viral STIs since 1995.

Conclusions: Self help initiatives and awareness among homosexual men in 1983–4 contributed significantly to a fall in HIV transmission among MSMs, and the general campaigns of 1986–7 were associated with similar effects on all STI transmission. Both effects seem to have occurred through changing sexual behaviour, and probably contributed to the UK’s low national HIV prevalence. Bacterial STI incidence has increased significantly since 1995 and there is no evidence that recent prevention initiatives have reduced HIV transmission among MSMs, hence sexual health initiatives need to be comprehensively reinvigorated in England.

(*Sex Transm Inf* 2001;77:242–247)

Keywords: HIV; gonorrhoea; syphilis; interventions; health promotion; homosexual males; heterosexuals; England

HIV and STI Division, Communicable Disease Surveillance Centre

A Nicoll
G Hughes
M Donnelly
K Fenton
B Evans
O N Gill
M Catchpole

Statistics Unit, Public
Health Laboratory
Service, 61 Colindale
Avenue, London
NW9 5EQ, UK
S Livingstone
D De Angelis

MRC Biostatistics
Unit, University of
Cambridge, Institute
of Public Health,
University Forvie Site,
Robinson Way, Hills
Road, Cambridge
CB3 9NF, UK
D De Angelis

Department of STDs,
Mortimer Market
Centre, Royal Free and
University College
Medical School, Off
Capper Street, London
WC1E 6AU, UK
K Fenton

Correspondence to:
Angus Nicoll
anicoll@phls.org.uk

Accepted for publication
10 April 2001

website
extra

Further figures and tables
are on the STI website

www.sextransinf.com

Introduction

The primary purpose of programmes intended to control sexually transmitted infections (STIs) is the interruption or reduction of transmission. Activities that contribute to achieving this goal include the reduction of risky sexual behaviours, case finding, screening, and partner notification.¹ Programmes targeting human immunodeficiency virus (HIV) are no different in this respect from those aimed at other STIs.

AIDS cases began to be diagnosed in the United Kingdom in late 1981. Indigenous HIV transmission then accelerated abruptly in the early 1980s, primarily through sex between men.² Public health responses began in 1983–4 with information and awareness initiatives run by, and directed at, members of the gay (male homosexual) community. They were followed, in 1986, by campaigns for the whole population (table 1).³ These interventions were more timely and vigorous than in other European countries⁴ and some have suggested that this is the reason that HIV prevalence in the United Kingdom is lower than in most of continental Western Europe.^{5–6} However, the campaigns,

and ongoing programmes, were also criticised as being unfocused, wasteful, or unnecessary.⁷ This analysis aims to evaluate the possible impact on sexual health outcomes of the anti-AIDS campaigns by examining national time series data for temporal associations between the interventions and key outcome measures: HIV transmission and diagnoses among men having sex with men (transmission through drug injecting is outside the scope of this analysis, and most of the heterosexually transmitted HIV infections diagnosed in the United Kingdom were acquired abroad⁸), attendances, and diagnoses of specific STIs at genitourinary medicine clinics. This is the first national evaluation for the United Kingdom using HIV statistics and STI surveillance data.

Methods

The period 1971–99 was chosen. A period extending back to the start of the 1970s was investigated because changes in STI epidemiology were taking place in the 1970s and it was important to distinguish these longer term trends from those that might be attributed to AIDS awareness and campaigns in the 1980s

Key messages

- AIDS awareness and campaigns to reduce HIV transmission in England in the early and mid-1980s had major impacts in reducing indigenous homosexual transmission of HIV and heterosexual transmission of other STIs, in the general population.
- The success of the early campaigns probably also explains why overall prevalence of HIV in the United Kingdom is lower than most other Western European countries.
- There is however evidence of substantial ongoing transmission of HIV among homosexual and bisexual men in the 1990s.
- Attendances at genitourinary medicine clinics with sexually transmitted diseases have increased substantially since 1995.
- These data strongly support the need for vigorous new sexual health and HIV/AIDS strategies.

and 1990s. The analyses were confined to England as detailed STI data were only available for that area. Details of AIDS awareness and campaigns from 1983 were gathered from historical sources: published articles, the HIV-AIDS media library at the London School of Hygiene and Tropical Medicine, Health Education Authority files, press records and discussions with historical authorities, and those active in the field.

Estimates of HIV transmission among men having sex with men (MSMs) were derived using back calculation² based on data to the end of 1995 (since 1996 such estimates have been profoundly affected by more effective therapy), and reports of new HIV diagnoses corrected for reporting delays and averaged to overcome seasonal reporting fluctuations. Routine data on annual attendances at genitourinary medicine (GUM) clinics in England were examined for trends.⁹ Categories examined were total attendances, new attendances requiring and not requiring treatment, and four specific STI diagnoses whose reporting format changed little over the period; infectious syphilis (KC60 code A1-A3), uncomplicated gonorrhoea (B1-B2), all genital herpes (C11) and all genital warts (C10). (Chlamydia and non-specific urethritis were excluded; chlamydia testing only became available in the late 1980s when the chlamydia diagnosis started taking cases from “non-specific urethritis.” These two categories were therefore not useful for time trend analysis over the study period.) Male infectious syphilis was especially associated with MSMs in the 1970s and 1980s¹⁰ (Routine annual GUM clinic returns did not start to identify subtotals of selected infections as being contracted through sex between men until after 1987) and male homosexual transmission of syphilis was monitored by annual ratios of male to female infectious syphilis cases.

Otherwise data were analysed separately for males and females and total population rates per 100 000 were calculated using mid-year population estimates for England from the Office for National Statistics. The annual rates of the STI categories were analysed with STATA 6.0 for Windows using normal regression models to test for trends. The pattern of change across the whole period was described, estimating linear trends. Probabilities expressed in the

results are the likelihood of differences or changes being by chance alone and refer to differences between groups (males *v* females), differences from no change in rates (for rises or falls in rates), or differences in rates over time (for when trends in rates seem to have steepened or flattened). The impact of the sexual component of the campaigns was examined by observing the trends in rates over narrower periods before, during, and after the most intense interventions of 1983 to 1984 (homosexual males) and 1986 to 1987 (general population).

Results**AIDS AND HIV SEXUAL HEALTH CAMPAIGNS**

Attempts to spread information on AIDS, and how to avoid it, began informally in the male homosexual community in early 1983, led by various non-governmental organisations (table 1).^{3, 4} The gay press gave extensive coverage to AIDS issues thereafter resulting in growing awareness in the gay community between 1983 and 1984 (table 1).³ A government led programme of education for the general population began with a major campaign in March 1986. There were larger campaigns in the winter of 1986-7 (including a leaflet sent to every household in January 1987) and the spring of 1987. These generated intense media coverage. In “AIDS week” in February 1987 the topic dominated all broadcasting schedules.^{3, 4} After 1987 general and MSM programmes continued at lower intensity (table 1). Notable features were a character with HIV appearing in a television serial (*EastEnders*) (1990), personal testimony from prominent people (1991), and a new UK HIV/AIDS strategy in 1995.¹¹ Two periods of intense national publicity could therefore be identified; a mix of awareness and then self help campaigning among MSMs from early 1983 to the end of 1987 and a more identifiable set of campaigns for the general population, from March 1986 to the end of 1987.^{3, 4} Both periods lacked distinct ends as they evolved into ongoing anti-HIV sexual health promotion. (The periods are indicated in the figures by vertical lines.)

Trends in HIV transmission and new attendances at NHS GUM clinics during and after awareness and AIDS campaigns

HIV TRANSMISSION THROUGH SEX BETWEEN MEN
Back calculation estimates indicate that incidence of HIV among MSMs was most intense around 1983 with about 6000 transmissions occurring by 1984. A rapid decrease in transmission that year was followed by plateauing (fig 1). Estimated numbers of new transmissions and diagnoses changed little after 1987, averaging 1300 to 1400 per annum (fig 1).

NEW ATTENDANCES AT NHS GUM CLINICS

Over the whole period (1971-99) the annual number of attendances, seen at GUM clinics increased threefold from 299 905 to 1 052 719. (Tables of annual data totals are available on the STI website.) The upward trend was steeper in females than males ($p < 0.001$) and from 1994 female attendance rates overtook those of males

Table 1 AIDS and HIV awareness and related sexual health promotion campaigns, UK 1983–99

Date	Event	Target group	Description
1983–5	Initial informal awareness	Gay community	Gay press coverage, informal campaigns organised by various non-government and community based organisations (NGOs and CBOs), eg Terrence Higgins Trust, Gay Switchboard, etc
1983–present	NGO and CBO campaigns, includes various leaflets about prevention, testing, treatment, housing legal issues.	Gay community (initially)	Initially, informal information sharing in the gay community. Later organised programmes of education for male gay community, other risk groups, and the general public.
March 1986	Government AIDS press campaign	General public	Press and posters, general information, massive media coverage
January 1987	Leaflet drop	General public	General household leaflet drop, massive media coverage
February 1987	“AIDS—don’t die of ignorance” campaign	General public	TV, national press, posters, massive media coverage
Spring 1987	AIDS Week	General public	All media channels, massive media coverage
	“AIDS—you’re as safe as you want to be” campaign	General public/youth	National press and youth press, extensive media coverage
Summer 1988	Holiday campaign	Young people	TV and posters
Summer 1988	AIDS the facts ads campaign	General public	TV and national press
Winter 1988–9 and annually thereafter	Safer sex campaigns	Gay community	Gay press and other gay outlets
Spring 1990	Personal testimony of experts campaign—“talking heads”	General public	TV and national press
Spring 1990	<i>EastEnders</i> introduces HIV into story line	General public	Inclusion of HIV infected person in a popular TV series
Spring 1991	“Mrs Dawson” campaign	General public	Cinema and TV
November 1991	Personal testimony	General public	Cinema and TV
Spring 1992	Red ribbon campaign	General public	General media outlets
Summer 1992	“Mrs Dawson” campaign		TV and cinema
	Personal testimony campaign	Ethnic minority groups	TV
1992	HIV/AIDS the facts	General public	Radio, local and minority ethnic press
1992–4	Travel facts, condom use	Gay community and young people	Radio, posters, TV
Spring 1994 and annually thereafter	Experimenting/uncertain campaign	Young women	Selected general press
Spring 1995	Postponement campaign	All groups	Gay press
December 1995	New HIV-AIDS strategy		Young women’s magazines Policy document

(fig w1 on the *STI* website). Within these general trends there were different patterns for component categories.

NEW ATTENDANCES REQUIRING AND NOT REQUIRING TREATMENT

Annual numbers of attendances requiring treatment more than doubled from 147 423 to

361 020 (fig 2). Between 1971 and 1986 male rates (per 100 000 total male population) increased with an average annual rise of $19.1/10^5$. This upward trend was significant ($p < 0.001$). For females the rate also increased, on average by $16.2/10^5$ annually ($p < 0.001$). During 1987, the rates fell (fig 2) by $117/10^5$ for males and by $42.1/10^5$ for females. Rates of attendances not requiring treatment (“the worried well”) unlike those of attendance requiring treatment rose rather than fell, male by $47.4/10^5$ and $57.8/10^5$ in men and women respectively. Rates in both sexes declined again in 1988. In contrast, between 1987 and 1995, rates of male attendances requiring treatment declined by an average of $7.6/10^5$ annually. Female rates re-attained and exceeded the 1986 level by 1988, but then remained essentially unchanged between 1988 and 1994 with no evidence of an upward or downward trend ($p = 0.81$). Between 1995 and 1999 male rates increased again at an average annual increase of $50.5/10^5$. The rising trend was significant ($p < 0.001$), and rates regained and exceeded 1986 levels (fig 2). Female rates also rose steeply and significantly at an average of $43.0/10^5$ per annum ($p < 0.001$).

Trends in specific diagnoses during and after awareness and AIDS campaigns

INFECTIOUS SYPHILIS

The annual number of new diagnoses decreased from 1606 to 401 (fig 3A). Trends differed for males and females. Between 1971 and 1978 male rates rose by an average of $0.7/10^5$ annually. Rates declined between 1978 and 1983, by an average of $0.6/10^5$ annually. A




Figure 1 Back calculation estimates of HIV incidence in homosexual and bisexual men: England and Wales 1979–96 (median and 95% confidence intervals) and new diagnoses of HIV probably acquired through sex between men.




Figure 2 Annual rates of new attendances requiring treatment at genitourinary medicine clinics in England; males and females, 1971-99.




Figure 3 (A) Annual rates of new diagnoses of infectious syphilis; males and females, 1971-99. (B) Annual ratio of male to female new diagnoses of infectious syphilis at genitourinary medicine clinics in England 1971 to 1999.

steeper decline began in 1983 with the rate falling by a total of $6.3/10^5$ by 1988. The rate of decline was significantly steeper than in the preceding period ($p < 0.001$) (fig 3A). The fall was statistically significant. Rates remained low between 1988 and 1998. There was a rise in 1999 especially among gay men.⁹ Female rates were lower but also declined after 1982 (fig 3A). The male to female ratio increased slowly

until the early 1980s peaking at 8:1 in 1983 because of a fall in female cases that year (fig 3B). There was then a more rapid decline to 2.7:1 in 1987. The declines in MSM HIV transmission, male infectious syphilis, and the male to female syphilis ratio coincided with the AIDS awareness and campaigns among homosexual men in 1983-4 (figs 1 and 3).

GONORRHOEA

Before 1975 trends differed between the sexes but were similar after that. Male rates declined from 1975 to 1985, with an average annual decrease of $4.1/10^5$ (fig 4) ($p < 0.001$). From 1985 to 1988 rates declined significantly more steeply compared with 1982 to 1985 ($p = 0.001$) and 1988 to 1991 ($p < 0.001$) at an average of $28.3/10^5$ annually, falling by $80.5/10^5$ (fig 4). The rates changed little between 1988 and 1991 but then fell to $28/10^5$ in 1995 after which they rose again, increasing to $43.7/10^5$ in 1999. Rates in women followed a similar pattern (fig 4) rising between 1971 and 1975, declining slowly at $2.07/10^5$ annually between 1975 and 1985, and showing a steeper decline between 1985 and 1999 with the rate of decline in that period being significantly steeper, compared with 1982 to 1985 ($p = 0.001$) and 1988 to 1991 ($p = 0.002$). There was a fall of $43.2/10^5$ between 1985 and 1988. After falling from $27.9/10^5$ in 1988 to $12.9/10^5$ in 1994, female rates started increasing again in 1995, rising to $19.6/10^5$ in 1999.

GENITAL HERPES (FIG W2 ON STI WEBSITE)

Over the whole period, the annual new diagnoses increased more than sevenfold from 3671 to 29 221 (fig w2 on STI website). Rates quadrupled for males (a total increase of $37.5/10^5$) and increased 17-fold ($64.3/10^5$) for females. Rates increased steadily between 1971 and 1981 but steepened between 1981 and 1984 (fig w2). Rates declined between 1986 and 1988, falling by $5.8/10^5$ for males and $3.5/10^5$ for females. After 1988 rates resumed their upward courses; and after 1991, female rates increased more steeply, exceeding male rates. In men the rate changed little after 1992 but for women the rate continued to increase. Rates stopped increasing between 1996 and 1999 in females (fig w2).

GENITAL WARTS

Over the whole period, the annual numbers of new diagnoses increased eightfold from 13 730 to 113 528 (fig w3 on STI website). Rates followed similar trends for males and females (fig w3). From 1971 to 1982 they rose slowly in both sexes, then the rate of increase steepened between 1982 and 1987. For males the average annual rise between 1982 and 1987 was $21.1/10^5$ but the male rates did not change significantly ($p = 0.252$) between 1987 and 1991. For females there was an average annual increase of $14.9/10^5$ between 1982 and 1987, followed by a significantly shallower ($p = 0.001$) rise of $3.7/10^5$ annually between 1987 and 1991. After a small decline in 1995 rates rose again from 1995 to 1999 increasing annually by $11.6/10^5$ and $8.0/10^5$ in males and females




Figure 4 Annual rates of attendances at genitourinary clinics in England with new diagnoses of gonorrhoea; males and females 1971 to 1999.

respectively. The upward trends after 1995 were significant for males ($p = 0.02$) and for females ($p = 0.014$).

Discussion

The data presented are consistent with the theory that sexual health components of the UK's AIDS campaigns of 1986–7 influenced sexual behaviour in the general population. The temporal associations of the campaigns and substantial rises in new attendances at GUM clinics not requiring treatment (also reflected by increases in HIV testing¹²), and falls in all treated cases, gonorrhoea, and genital herpes are very unlikely to be coincidental. The delayed slowing of previously rising rates of genital warts in 1987 is consistent with that disease's longer incubation period. The bacterial infections, syphilis, and gonorrhoea had begun declining in the 1970s (figs 3A and 4) and the reasons for this are unknown. However, the decline in the period 1996–7 was significantly steeper and also affected herpes and warts. There were also significant increases in condom sales and use at that time.¹² Whatever they achieved in preventing HIV transmission, the general population campaigns were associated with reduced transmission of other STIs, seemingly through behaviour change and heightened awareness of risky sexual practices.

The fall in HIV transmission among homosexual and bisexual men (MSMs) in 1983–4 (fig 1) coincided with the AIDS awareness and informal AIDS campaigns conducted by the gay community. However, could that decline be because most men at highest behavioural risk had been infected with HIV by that date? The 6000 men estimated to have been infected before 1984 represented only 6% of MSMs attending GUM clinics in England, a fraction of the MSM population at higher behavioural risk.¹⁴ Therefore an "exhaustion of the susceptibles" is unlikely. The simultaneous fall in the incidence of male infectious syphilis (fig 3), and contemporary reports of changes in gay male behaviour around 1983–4,^{3 15 16} make it more plausible that the growing gay male awareness and AIDS campaigns caused falls in transmission of both HIV and syphilis among MSMs through changing behaviour. The

decline in ulcerative syphilis may also have accelerated the reduction in HIV transmission.¹⁷

Whether the sexual health campaigns of 1983–7, and initiatives against drug injecting,⁸ protected the United Kingdom from the impact of HIV experienced by many other Western European countries (where the population prevalence of HIV is up to six times higher than in the United Kingdom⁶) is more arguable. However, early interventions against HIV emerging in a country are considered to be most effective.¹⁸ It is striking that among Western European countries only Sweden, Norway, and Finland (countries which also mounted vigorous public health campaigns against HIV early relative to their local epidemics⁴) have HIV prevalences below that of the United Kingdom.⁶ This is despite a particular overlay in the United Kingdom of HIV from African countries.¹⁹ Therefore it is probable that an early vigorous approach to AIDS blunted the penetration of HIV in the United Kingdom.

The improving trends in gay sexual health indices in the mid-1980s were not sustained. High levels of risk taking behaviour were recorded in gay and bisexual men in the early 1990s.^{20 21} The evidence of recent, ongoing HIV transmission among MSMs, and increasing STI diagnoses in the general population is especially disquieting.⁹ There is little evidence from these data of any positive impact of the 1995 strategy.^{9 11} "Risk profiles," and other HIV reporting data for homosexual and bisexual men show results consistent with ongoing HIV transmissions and continuing or even rising levels of risk taking.^{22 23} Specifically since 1995 gonorrhoea diagnoses have increased significantly,^{9 24} several outbreaks of syphilis have occurred^{25 26} and behavioural monitoring suggests increasing behavioural risk among MSMs.²⁷ There has been a rise in rectal gonorrhoea so that the rise in gonorrhoea in gay men is unlikely to be the result of adoption or safe sex practices.¹⁹

The recent general increases in STI diagnoses are likely to be genuine. There were changes in reporting practices and codes in 1995 but these were minor compared with those in 1988, which occurred without appreciable effect on trends.⁹ Responsibility for the collection of clinic returns was transferred to CDSC in 1995.⁹ However, the upward trends have been sustained for over 4 years^{9 24} and are unlikely to be explicable by minor alterations in coding or collection practices, which would mostly affect a single year. Nevertheless, it should be emphasised that the rises in attendances requiring treatment do not necessarily indicate increased incidence of all STIs. Increasing ability to diagnose chlamydia, greater healthcare seeking behaviour, or improved clinic services explain some of the rises.⁹ However, rates of gonorrhoea, short term trends in which are considered a sensitive indicator of sexual behaviour, have increased by over 30% since 1994.^{9 24} They cannot be explained by new diagnostic molecular techniques which are not yet in widespread use.

This rise, in combination with the trends in STIs and behaviours among MSMs,^{23 25 26} makes it likely that the impact of the HIV/AIDS campaigns of the 1980s has worn off. New generations becoming sexually active since then are likely to be less aware of the dangers of HIV and it seems likely that there has recently been an increase in risky sexual behaviour.⁸

In conclusion, the results of this analysis show that almost certainly the AIDS awareness and campaigns among gay men had a major impact, reducing HIV transmission in England in the early 1980s, and that the later general population campaigns reduced the transmission of other STIs among heterosexuals, probably also reducing the penetration of HIV into the population. The more recent sustained HIV transmission among gay men and increasing transmission of gonorrhoea and syphilis among all gay men and heterosexuals indicate that the forthcoming sexual health strategy for England, in particular its HIV/AIDS and STI components, will have to be comprehensive and vigorous.^{28 29}

This analysis is entirely reliant on primary STI data supplied by genitourinary medicine clinics and HIV data were from voluntary confidential reporting by clinicians, microbiologists, and others to CDSC. The authors would also like to thank Michael Bland for producing visual presentations of these data and Angela McHenry and Pauline Deeks for management of the HIV and STI data. Virginia Berridge, Janet Mortimer, and Donald Acheson provided helpful advice and comments on earlier drafts of this paper.

Contributors: AN designed the analysis and wrote the paper; MD gathered the primary information on the HIV/AIDS campaigns and made early drafts of the paper while working as specialist registrar on placement at CDSC. All statistical analyses were undertaken by SL, apart from back calculation of HIV incidence, which was undertaken by DDeA with ONG. STI data were collated, compiled, and analysed under the directions of GH and MC; BE and ONG undertook this role for HIV data; GH, MC, BE, KF, and ONG helped with interpretation of the data and commented on the text. GH and KF contributed significantly to the writing of the paper. Angus Nicoll acts as guarantor of the paper.

Conflicting interests: There were no known conflicting interests of the authors and the costs of the study were met from core PHLS funding.

- 1 Catchpole MA. The role of epidemiology and surveillance systems in the control of sexually transmitted diseases. *Genitourin Med* 1996;72:321-9.
- 2 Report of a Working Group (Chairman: Professor NE Day) convened by the Director of the Public Health Laboratory Service on behalf of the Chief Medical Officers. The incidence and prevalence of AIDS and the prevalence of other severe HIV disease in England and Wales for 1995 to 1999; projections using data to the end of 1994. *CDR Rev* 1996;6:R1-24.
- 3 Berridge V. *AIDS in the UK, the making of policy*. 1981-94. Oxford: OUP, 1996.
- 4 Wellings K, Field B. *Stopping AIDS. AIDS/HIV public education and the mass media in Europe*. London: Longman, 1996.
- 5 Department of Health. *HIV/AIDS strategy conference report (27 October 1998)*. London: DoH, 1998.
- 6 Joint United Nations Programme on AIDS (UNAIDS) and the World Health Organization (WHO). *Report on the global HIV/AIDS epidemic June 1998*. Geneva: UNAIDS and WHO, 1998.
- 7 Craven BM, Stewart GT, Taghavi D. Amateurs confronting specialists: expenditure on AIDS in England. *J Public Policy* 1994;13:305-25.

- 8 Berridge V. *AIDS and British drug policy: continuity or change. AIDS and contemporary history*. Cambridge: CUP, 1993.
- 9 PHLS, DHSS and PS and the Scottish ISD(D)5 Collaborative Group. *Trends in sexually transmitted infections in the United Kingdom 1990-1999*. London: Public Health Laboratory Service, 2000.
- 10 Waugh MA. Studies on the recent epidemiology of early syphilis in West London. *Br J Vener Dis* 1972;48:534-41.
- 11 Departments of Health. *HIV and AIDS health promotion: an evolving strategy*. London: Department of Health, 1995.
- 12 Communicable Disease Surveillance Centre, Scottish Centre for Infection and Environmental Health, Institute of Child Health (London), and Oxford Haemophilia Centre. *AIDS and HIV Infection in the United Kingdom: monthly report. Sentinel surveillance of diagnostic HIV tests in England. CDR* 1997;17:307-8.
- 13 Goodrich J, Wellings K, McVey D. Using condom data to assess the impact of HIV/AIDS preventive interventions. *Health Educ Res* 1998;13:267-74.
- 14 Giesecke J, Johnson A, Hawkins A, et al. An estimate of the prevalence of human immunodeficiency virus infection in England and Wales by using a direct method. *J Roy Stat Soc A* 1994;157:89-103.
- 15 Carne CA, Weller IVD, Johnson AM, et al. Prevalence of antibodies to human immunodeficiency virus, gonorrhoea rates, and changed sexual behaviour in homosexual men in London. *Lancet* 1987;i:656-8.
- 16 Weller IVD, Hindley DJ, Adler MW, et al. Gonorrhoea in homosexual men and media coverage of the acquired immune deficiency syndrome in London 1982-3. *BMJ* 1984;289:1041.
- 17 Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Inf* 1999;75:3-17.
- 18 Robinson NJ, Mulder DW, Auvert B, et al. Proportions of HIV infections attributable to other sexually transmitted diseases in a rural Ugandan population: simulation model estimates. *Int J Epidemiol* 1997;26:180-9.
- 19 Communicable Disease Surveillance Centre, Scottish Centre for Infection and Environmental Health, Institute of Child Health (London), and Oxford Haemophilia Centre. *AIDS and HIV Infection in the United Kingdom: monthly report. HIV infection acquired through sex between men and women. Commun Dis Rep* 2000;9:157-60.
- 20 Hart GJ, Fitzpatrick RM, Boulton M, et al. Risk behaviour, anti-HIV and anti-hepatitis B core prevalence in clinic and non-clinic samples of gay men in England, 1991-1992. *AIDS* 1993;7:863-9.
- 21 Davies PM, Weatherburn P, Hunt AJ, et al. The sexual behaviour of young gay men in England and Wales. *AIDS Care* 1992;4:259-72.
- 22 Unlinked Anonymous Surveys Steering Group. 'Risk Profiles' in prevalence of HIV in the United Kingdom 1998. Report of the Unlinked Anonymous Seroprevalence Monitoring Programme in England and Wales. Department of Health, Public Health Laboratory Service, Institute of Child Health (London), Scottish Centre for Infection and Environmental Health, December 1999: 25-8.
- 23 Communicable Disease Surveillance Centre, Scottish Centre for Infection and Environmental Health, Institute of Child Health (London), and Oxford Haemophilia Centre. *AIDS and HIV-1 Infection in the United Kingdom: monthly report. HIV and sexually transmitted infections acquired through sex between men. Commun Dis Rep* 2000; 10:237-40.
- 24 Fenton KA, Rogers PA, Simms I, et al. Increasing N gonorrhoea reports—not only in London. *Lancet* 2000;355: 1907.
- 25 Sexually transmitted diseases quarterly report: syphilis national and international epidemiology. *Commun Dis Rep* 1999;9:38-9.
- 26 Communicable Disease Surveillance Centre. Increased transmission of syphilis in men who have sex with men reported from Brighton and Hove. *CDR* 2000;10:177.
- 27 Dodds JP, Nardone A, Mercey DE, et al. Increasing high risk sexual behaviour amongst homosexual men in London 1996-1998: a repeated cross sectional questionnaire study. *BMJ* 2000;320:1510-1.
- 28 First ever government strategy on sexual health launched. Press Release, 23 March 1999. London: Department of Health (No 1999/0166).
- 29 Department of Health and the HIV/AIDS Strategy Steering Group. *Developing an HIV/AIDS strategy: progress report*. London: NHS Executive, April 2000.