

# Subject index

- aborted lesions**, Aborted genital herpes simplex virus lesions: findings from a randomised controlled trial with valaciclovir, **435**
- abortion**, Prevalence of and risk indicators for STIs among women seeking induced abortions in two urban family planning clinics in Shandong province, People's Republic of China, **e3**
- access to services**, A survey of STI policies and programmes in Europe: preliminary results, **380**
- adenovirus**, Characteristics of adenovirus associated urethritis, **445**
- adolescent reproductive health**, Adolescent reproductive health interventions, **315**
- adolescent sexual health**, Adolescence: a time of risk taking, **314**  
Adolescent sexual health in Sweden, **352**  
"A single petticoat", **318**  
Herpes simplex virus type 1 infection: a sexually transmitted infection of adolescence?, **346**  
National guidelines for the management of suspected sexually transmitted infections in children and young people, **324**
- adolescents**, *Chlamydia trachomatis* infection among sexually active young women in Italy, **e4**  
Detection of *Chlamydia trachomatis* in an Australian high school student population, **194**  
HIV tests in young adolescents attending a GUM clinic, **386**  
National guideline for the management of suspected sexually transmitted infections in children and young people, **324**  
News from the frontline: sexually transmitted infections in teenagers attending a genitourinary clinic in south east London, **349**  
Post-treatment sexual and prevention behaviours of adolescents with sexually transmitted infections, **365**  
Predictors of infection with *Trichomonas vaginalis*: a prospective study of low income African-American adolescent females, **360**  
Provider willingness to screen all sexually active adolescents for chlamydia, **369**
- adult entertainment industry**, HIV negative certification and sexual health issues facing performers in the adult entertainment industry in the UK, **311**
- Africa**, 15 Month follow up of African children following vaginal cleansing with benzalkonium chloride of their HIV infected mothers during late pregnancy and delivery, **267**  
Donovanosis, **452**  
HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- age**, *Chlamydia trachomatis* in non-gonococcal urethritis patients and their heterosexual partners: routine testing by polymerase chain reaction, **286**
- AIDS**, A pilot study of a rapid assessment method to identify places for AIDS prevention in Cape Town, South Africa, **i106**  
Behavioural surveillance: the value of national coordination, **398**  
HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**  
Sociodemographic context of the AIDS epidemic in a rural area in Tanzania with a focus on people's mobility and marriage, **i97**
- anogenital warts**, Adult and paediatric contact immunotherapy with squaric acid dibutylester (SADBE) for recurrent, multiple, resistant, mucocutaneous anogenital warts, **309**
- antenatal screening**, Offering routine antenatal testing for HIV and hepatitis B in the rural setting of Cornwall, **133**
- antimicrobial resistance**, Gonorrhoea in London: usefulness of first line therapies, **106**
- army**, Surveillance of HIV in the army of the Republic of Cyprus (SHARC); rationale, design, and implementation of an inexpensive system, **50**
- artificial insemination**, Four layer discontinuous gradient for HIV, **388**
- attitudes**, Knowledge about the prevention of sexually transmitted diseases: a longitudinal study of young women from 16–23 years of age, **339**
- Australia**, Screening for STIs in individuals with HIV infection, **387**  
The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice, **278**
- bacterial vaginosis**, Validation of a simplified grading of Gram stained vaginal smears for use in genitourinary medicine clinics, **413**
- Bangladesh**, Sexually transmitted infections in Bangladeshis resident in the UK: a case-control study, **120**  
STD in Bangladesh's trucking industry: prevalence and risk factors, **31**
- Benin**, Role of core and bridging groups in the transmission dynamics of HIV and STIs in Cotonou, Benin, West Africa, **i69**
- bilateral groove sign**, Bilateral groove sign with penoscrotal elephantiasis, **458**
- black Africans**, Pathways to HIV testing and care by black African and white patients in London, **37**
- Bodo saltans**, TV or not TV?, **185**
- Book reviews**, ABC of AIDS, 5th ed. M W Adler. BMJ Books, 2001, **311**  
Human papillomaviruses. Clinical and Scientific Advances. JC Sterling, SK Tyring. Arnold Publishers, 2001, **312**  
The Changing Face of HIV and AIDS. R A Weiss, M W Adler, S L Rowland-Jones. Oxford University Press, 2001, **390**  
Understanding thrush, cystitis and women's genital symptoms. C Bradbeer. Family Doctor Publications in association with the BMA, 2001, **312**
- Brazil**, Sexual and reproductive health among female adolescents: preliminary results, **386**
- Britain**, Peer led HIV prevention among homosexual men in Britain, **158**  
The accuracy of reported sensitive sexual behaviour in Britain: exploring the extent of change 1990–2000, **26**
- Canada**, Geographical variations in the epidemiology of bacterial sexually transmitted infections in Manitoba, Canada, **i139**
- Candida albicans**, Vulvovaginitis due to fluconazole resistant *Candida albicans* following self treatment with non-prescribed triazoles, **308**
- Cape Town**, A pilot study of a rapid assessment method to identify places for AIDS prevention in Cape Town, South Africa, **i106**
- case management**, Syndromic management of sexually transmitted diseases at primary care level, **e2**
- Castleman's disease**, Multicentric plasma cell variant Castleman's disease mimicking intrapulmonary malignancy, **304**
- cervical cancer**, Prevalence and risk factors of sexually transmitted infections and cervical neoplasia in women's health clinics in Nicaragua, **204**
- cervix**, Tuberculosis of the cervix: case presentation and a review of the literature, **62**
- chancroid**, Erythema nodosum induced by chancroid, **388**
- chemists and druggists**, The practice of STI treatment among chemists and druggists in Pokhara, Nepal, **223**
- children**, Managing the very young patient: a conflict between the requirements of the Children Act and the VD regulations?, **332**  
National guideline for the management of suspected sexually transmitted infections in children and young people, **324**
- Children Act**, Managing the very young patient: a conflict between the requirements of the Children Act and the VD regulations?, **332**
- China**, Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Chinese**, Hepatitis B virus infection among Chinese residents in the United Kingdom, **166**  
Reasons for not using condoms among the Hong Kong Chinese population: implications for HIV and STD prevention, **180**
- chlamydia**, Chlamydia screening and sexual health, **396**  
Factors affecting co-infection with genital chlamydia and genital gonorrhoea in an urban genitourinary medicine clinic, **387**  
Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system, **40**  
Gonorrhoea and chlamydia core groups and sexual networks in Manitoba, **i145**  
Investigation into the acceptability and effectiveness of a new contact slip in the management of *Chlamydia trachomatis* at a London genitourinary medicine clinic, **422**  
Management of screened chlamydia positive women, **155**  
Outreach health adviser in a community clinic screening programme improves management of genital chlamydia infection, **101**  
Provider willingness to screen all sexually active adolescents for chlamydia, **369**  
The value of mass screening for chlamydia control in high prevalence communities, **i64**
- chlamydia infections**, Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men, **190**
- chlamydia screening**, Provision of chlamydia testing in a nationwide service offering termination of pregnancy: with data capture to monitor prevalence of infection, **416**
- Chlamydia trachomatis**, *Chlamydia trachomatis* diagnostics, **232**  
*Chlamydia trachomatis* in an Australian high school student population, **194**  
*Chlamydia trachomatis* infection among sexually active young women in Italy, **e4**  
*Chlamydia trachomatis* infections in eastern Europe: legal aspects, epidemiology, diagnosis, and treatment, **115**

- Chlamydia trachomatis* in non-gonococcal urethritis patients and their heterosexual partners: routine testing by polymerase chain reaction, **286**
- Clinical characteristics of *Chlamydia trachomatis* infections in a general outpatient department of obstetrics and gynaecology in the Netherlands, **e6**
- Comparing cost effectiveness of screening women for *Chlamydia trachomatis* in systematic and opportunistic approaches, **73**
- Cost effectiveness analysis of a population based screening programme for asymptomatic *Chlamydia trachomatis* infection in women, **76**
- Cost effectiveness of screening for *Chlamydia trachomatis*: a review of published studies, **406**
- Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- Chlamydia trachomatis assay**, Resolution of the recent performance problem of Abbott LCx *Chlamydia trachomatis* assay Issues of repeat testing for confirmation of chlamydial infection, **225**
- chlamydia transmission**, Sexual network structure as an indicator of epidemic phase, **i152**
- ciprofloxacin**, Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**
- clarithromycin**, Neuropsychiatric reaction induced by clarithromycin, **75**
- cohort study**, Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Colorado**, Reaching homeless youths for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* screening in Denver, Colorado, **357**
- community effects**, Community effects on the risk of HIV infection in rural Tanzania, **261**
- computed tomograph scan**, Multicentric plasma cell variant Castleman's disease mimicking intrapulmonary malignancy, **304**
- condom use**, Condom access does not ensure condom use: you've got to be putting me on, **225**
- Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Post-treatment sexual and prevention behaviours of adolescents with sexually transmitted infections, **365**
- Reasons for not using condoms among the Hong Kong Chinese population: implications for HIV and STD prevention, **180**
- Short term interventions are not enough, **80**
- condoms**, Health seeking and sexual behaviour in patients with sexually transmitted infections: the importance of traditional healers in Thyolo, Malawi, **127**
- congenital syphilis**, Congenital syphilis—missed opportunities for prenatal intervention, **224**
- contraception**, Adolescent sexual health in Sweden, **352**
- News from the frontline: sexually transmitted infections in teenagers attending a genitourinary clinic in south east London, **349**
- core group**, Gonorrhoea and chlamydia core groups and sexual networks in Manitoba, **i145**
- The feasibility of integrated STI prevalence and behaviour surveys in developing countries, **187**
- cost benefit analysis**, The potential role of suppressive therapy for sex partners in the prevention of neonatal herpes: a health economic analysis, **425**
- cost effectiveness**, Comparing cost effectiveness of screening women for *Chlamydia trachomatis* in systematic and opportunistic approaches, **73**
- Cost effectiveness analysis of a population based screening programme for asymptomatic *Chlamydia trachomatis* infection in women, **76**
- Cost effectiveness of screening for *Chlamydia trachomatis*: a review of published studies, **406**
- Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- cost effectiveness analysis**, The potential role of suppressive therapy for sex partners in the prevention of neonatal herpes: a health economic analysis, **425**
- costs**, A survey of STI policies and programmes in Europe: preliminary results, **380**
- co-trimoxazole**, Gonococcal perianal abscess: re-emergence after cessation of co-trimoxazole, **389**
- counselling**, Low incidence of syphilis among factory workers in Ethiopia: effect of an intervention based on education and counselling, **123**
- criminalisation**, The criminalisation of HIV transmission, **448**
- Crohn's disease**, Perianal Crohn's disease masquerading as perianal warts, **302**
- Cyprus**, Surveillance of HIV in the army of the Republic of Cyprus (SHARC); rationale, design, and implementation of an inexpensive system, **50**
- demography**, Sociodemographic context of the AIDS epidemic in a rural area in Tanzania with a focus on people's mobility and marriage, **i97**
- dermatitis**, Human T lymphotropic virus-I (HTLV-I) infection in patients with unclassifiable dermatitis in central Kerala, south India: a preliminary study, **e7**
- diagnosis**, Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**
- diagnostic test**, An evidence based approach to testing for antibody to herpes simplex virus type 2, **430**
- Using the evidence base on genital herpes: optimising the use of diagnostic tests and information provision, **160**
- donovanosis**, Donovanosis, **452**
- duty to warn**, The criminalisation of HIV transmission, **448**
- eastern Europe**, *Chlamydia trachomatis* infections in eastern Europe: legal aspects, epidemiology, diagnosis, and treatment, **115**
- education**, Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**
- Low incidence of syphilis among factory workers in Ethiopia: effect of an intervention based on education and counselling, **123**
- efavirenz**, Nevirapine + efavirenz based salvage therapy in heavily pretreated HIV infected patients, **72**
- empathy**, Assisting medical students to conduct empathic conversations with patients from a sexual medicine clinic, **246**
- enforcing obligation**, The criminalisation of HIV transmission, **448**
- England**, A postal survey to identify and describe nurse led clinics in genitourinary medicine services across England, **98**
- Resurgence of syphilis in England, **308**
- Enterobacter sakazakii**, Vaginal infection by *Enterobacter sakazakii*, **467**
- environmental contamination**, Contamination of environmental surfaces by genital human papillomaviruses, **135**
- eosinophilia**, Immune reconstitution eosinophilia due to schistosomiasis, **76**
- epidemic phase**, Risk network structure in the early epidemic phase of HIV transmission in Colorado Springs, **i159**
- The impact of the phase of an epidemic of sexually transmitted infection on the evolution of the organism, **i20**
- epidemicity**, Determinants of STD epidemics: implications for phase appropriate intervention strategies, **i3**
- epidemic**, Sexual network structure as an indicator of epidemic phase, **i152**
- The impact of HIV epidemic phases on the effectiveness of core group interventions: insights from mathematical models, **i78**
- epidemiology**, An introduction to mathematical models in sexually transmitted disease epidemiology, **7**
- Behavioural surveillance: the value of national coordination, **398**
- Chlamydia trachomatis* infections in eastern Europe: legal aspects, epidemiology, diagnosis, and treatment, **115**
- Current characteristics of male gonorrhoea outpatients in Athens, Greece, **e5**
- Geographical variations in the epidemiology of bacterial sexually transmitted infections in Manitoba, Canada, **i139**
- Heterosexual outbreak of infectious syphilis: epidemiological and ethnographic analysis and implications for control, **i164**
- HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**
- Risk network structure in the early epidemic phase of HIV transmission in Colorado Springs, **i159**
- The evolution of sexually transmitted infections in the Ukraine, **219**
- The geographical and temporal evolution of sexually transmitted disease epidemics, **i14**
- Epstein-Barr virus**, Progressive intrathoracic lymphadenopathy: EBV associated non-Hodgkin's lymphoma, **13**
- erythema nodosum**, Erythema nodosum induced by chancroid, **388**
- Ethiopia**, Low incidence of syphilis among factory workers in Ethiopia: effect of an intervention based on education and counselling, **123**
- ethnicity**, HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- Europe**, A survey of STI policies and programmes in Europe: preliminary results, **380**
- Services for sexually transmitted infections in Europe and central Asia, **320**
- exit editorial**, The curtain call, **392**
- factory workers**, Low incidence of syphilis among factory workers in Ethiopia: effect of an intervention based on education and counselling, **123**
- female**, Predictors of infection with *Trichomonas vaginalis*: a prospective study of low income African-American adolescent females, **360**

- female adolescents**, Sexual and reproductive health among female adolescents: preliminary results, **386**
- female sex worker**, HIV infection among people of foreign origin voluntarily tested in Spain. A comparison with national subjects, **250**  
Part time female sex workers in a suburban community in Kenya: a vulnerable hidden population, **271**  
Response of a sexually transmitted infection epidemic to a treatment and prevention programme in Nairobi, Kenya, **i114**
- field testing**, Performance of the rapid plasma reagin and the rapid syphilis screening tests in the diagnosis of syphilis in field conditions in rural Africa, **282**
- France**, Seroprevalence of HSV-1 and HSV-2 infection in the general French population, **201**
- general practitioners**, Genital herpes: general practitioners' knowledge and opinions, **198**
- genital herpes**, Genital herpes: general practitioners' knowledge and opinions, **198**  
Successful use of valaciclovir in a case of recurrent urticaria associated with genital herpes, **468**  
Using the evidence base on genital herpes: optimising the use of diagnostic tests and information provision, **160**
- genital ulcer**, Genital ulcers, stigma, HIV, and STI control in sub-Saharan Africa, **143**
- genital ulcer disease**, Coexistence of urethritis with genital ulcer disease in South Africa: influence on provision of syndromic management, **274**  
Donovanosis, **452**
- genitourinary medicine clinic attendees**, Recreational drug use by GUM clinic attendees, **310**
- genitourinary medicine clinics**, A postal survey to identify and describe nurse led clinics in genitourinary medicine services across England, **98**
- geographic epidemiology**, Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system, **40**
- GIS system**, Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system, **40**
- gonorrhoea**, Characteristics of gonorrhoea in Kermanshah, Iran, **460**  
Factors affecting co-infection with genital chlamydia and genital gonorrhoea in an urban genitourinary medicine clinic, **387**  
Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system, **40**  
Gonorrhoea and chlamydia core groups and sexual networks in Manitoba, **i145**  
Gonorrhoea in London: usefulness of first line therapies, **106**  
Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men, **190**
- granuloma inguinale**, Donovanosis, **452**
- Greece**, Current characteristics of male gonorrhoea outpatients in Athens, Greece, **e5**
- guidelines**, National guideline for the management of suspected sexually transmitted infections in children and young people, **324**  
Revised UK national guidelines on sexually transmitted infections and closely related conditions 2002, **81**
- Guinea-Bissau**, *Mycoplasma genitalium* is not associated with adverse outcomes of pregnancy in Guinea-Bissau, **289**
- GUM clinic**, HIV tests in young adolescents attending a GUM clinic, **386**  
Sexual health clinics for women led by specialist nurses or senior house officers in a central London GUM service: a randomised controlled trial, **93**
- gyrA gene**, Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**
- HAART**, Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**  
*Mycobacterium xenopi* pulmonary infection in an HIV infected patient under highly active antiretroviral treatment, **60**
- health advisers**, Outreach health adviser in a community clinic screening programme improves management of genital chlamydia infection, **101**
- health education**, Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**
- health policy**, From community to clients: the professionalisation of HIV prevention among gay men and its implications for intervention selection, **i176**
- health seeking behaviour**, Health seeking and sexual behaviour in patients with sexually transmitted infections: the importance of traditional healers in Thyolo, Malawi, **127**
- hepatitis**, Hepatitis, syphilis, and HIV sentinel surveillance in Mongolia 1999–2000, **223**
- hepatitis A**, Seroprevalence of hepatitis A immunity in male genitourinary medicine clinic attendees: a case control study of heterosexual and homosexual men, **174**
- hepatitis B**, Hepatitis B virus infection among Chinese residents in the United Kingdom, **166**  
Offering routine antenatal testing for HIV and hepatitis B in the rural setting of Cornwall, **133**  
Prevalence and incidence of hepatitis B virus infection in STD clinic attendees in Pune, India, **169**  
Understanding correlates of hepatitis B virus vaccination in men who have sex with men: what have we learned?, **374**
- hepatitis C**, A new way to catch hepatitis C, **459**
- herpes genitalis**, Polymerase chain reaction for diagnosis of genital herpes in a genitourinary medicine clinic, **21**
- herpes simplex virus**, Aborted genital herpes simplex virus lesions: findings from a randomised controlled trial with valaciclovir, **435**  
Herpes simplex virus type 1 infection: a sexually transmitted infection of adolescence?, **346**  
Polymerase chain reaction for diagnosis of genital herpes in a genitourinary medicine clinic, **21**  
Seroprevalence of HSV-1 and HSV-2 infection in the general French population, **201**  
Using the evidence base on genital herpes: optimising the use of diagnostic tests and information provision, **160**
- herpes simplex virus type 2**, An evidence based approach to testing for antibody to herpes simplex virus type 2, **430**
- herpes virus infections**, The potential role of suppressive therapy for sex partners in the prevention of neonatal herpes: a health economic analysis, **425**
- heterosexual men**, Chlamydia screening and sexual health, **396**
- heterosexuality**, Seroprevalence of hepatitis A immunity in male genitourinary medicine clinic attendees: a case control study of heterosexual and homosexual men, **174**
- high performance testing**, Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- HIV**, 15 Month follow up of African children following vaginal cleansing with benzalkonium chloride of their HIV infected mothers during late pregnancy and delivery, **267**  
Anti-HIV serology in patients with sexual dysphoria in screening test before sex change surgery, **75**  
Behavioural surveillance: the value of national coordination, **398**  
Community effects on the risk of HIV infection in rural Tanzania, **261**  
Diverse realities: sexually transmitted infections and HIV in India, **i31**  
Does increased aromatase activity in adipose fibroblasts cause low sexual desire in patients with HIV lipodystrophy?, **64**  
Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**  
Four layer discontinuous gradient for HIV, **388**  
From community to clients: the professionalisation of HIV prevention among gay men and its implications for intervention selection, **i176**  
Genital ulcers, stigma, HIV, and STI control in sub-Saharan Africa, **143**  
HIV negative certification and sexual health issues facing performers in the adult entertainment industry in the UK, **311**  
HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**  
*Mycobacterium xenopi* pulmonary infection in an HIV infected patient under highly active antiretroviral treatment, **60**  
Nevirapine + efavirenz based salvage therapy in heavily pretreated HIV infected patients, **72**  
Offering routine antenatal testing for HIV and hepatitis B in the rural setting of Cornwall, **133**  
Part time female sex workers in a suburban community in Kenya: a vulnerable hidden population, **271**  
Pathways to HIV testing and care by black African and white patients in London, **37**  
Peer led HIV prevention among homosexual men in Britain, **158**  
Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**  
Prevention interventions for HIV positive individuals, **393**  
Prostatic tuberculosis in an HIV infected male, **147**  
Response of a sexually transmitted infection epidemic to a treatment and prevention programme in Nairobi, Kenya, **i114**  
Risk network structure in the early epidemic phase of HIV transmission in Colorado Springs, **i159**  
Risky sexual behaviours among injection drugs users with high HIV prevalence: implications for STD control, **i170**  
Role of core and bridging groups in the transmission dynamics of HIV and STIs in Cotonou, Benin, West Africa, **i69**  
Screening for STIs in individuals with HIV infection, **387**  
Sex in the city: sexual behaviour, societal change, and STDs in Saigon, **i47**  
STD/HIV intervention and research programme Mwanza Region, NW Tanzania, **i91**  
STI epidemics in the Indian Ocean region: can the phase be assessed?, **i121**  
The criminalisation of HIV transmission, **448**

- The impact of HIV epidemic phases on the effectiveness of core group interventions: insights from mathematical models, **i78**
- The profile and context of the epidemics of sexually transmitted infections including HIV in Zimbabwe, **i40**
- Tuberculosis of the cervix: case presentation and a review of the literature, **62**
- Uptake of HIV testing in patients with a confirmed sexually transmitted infection, **389**
- HIV clinic services**, Pathways to HIV testing and care by black African and white patients in London, **37**
- HIV disease**, Lactic acidosis and abnormal liver function in advanced HIV disease, **139**
- HIV epidemicity**, HIV epidemicity in context of STI declines: a telling discordance, **467**
- HIV infection**, Peripheral neuropathy in patients with HIV infection: consider dual pathology, **462**
- Post-exposure prophylaxis for non-occupational exposure to HIV: current clinical practice and opinions in the UK, **130**
- Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men, **190**
- HIV prevalence**, HIV infection among people of foreign origin voluntarily tested in Spain. A comparison with national subjects, **250**
- HIV prevention**, Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**
- HIV surveillance**, Surveillance of HIV in the army of the Republic of Cyprus (SHARC); rationale, design, and implementation of an inexpensive system, **50**
- The feasibility of integrated STI prevalence and behaviour surveys in developing countries, **187**
- HIV testing**, HIV infection among people of foreign origin voluntarily tested in Spain. A comparison with national subjects, **250**
- HIV tests in young adolescents attending a GUM clinic, **386**
- HIV/STD interactions**, The effect of HIV, behavioural change, and STD syndromic management on STD epidemiology in sub-Saharan Africa: simulations of Uganda, **i55**
- homeless youths**, Reaching homeless youths for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* screening in Denver, Colorado, **357**
- homosexual men**, Peer led HIV prevention among homosexual men in Britain, **158**
- homosexuality**, Seroprevalence of hepatitis A immunity in male genitourinary medicine clinic attenders: a case control study of heterosexual and homosexual men, **174**
- human papillomavirus**, Contamination of environmental surfaces by genital human papillomaviruses, **135**
- Oestrogen receptor transcripts associated with cervical human papillomavirus infection, **210**
- Preventive human papillomavirus vaccination, **4**
- Risk factors for genital HPV DNA in men resemble those found in women: a study of male attendees at a Danish STD clinic, **215**
- human T lymphotropic virus-1**, Human T lymphotropic virus-1 (HTLV-1) infection in patients with unclassifiable dermatitis in central Kerala, south India: a preliminary study, **e7**
- immigrants**, Current characteristics of male gonorrhoea out-patients in Athens, Greece, **e5**
- immune restoration syndrome**, *Mycobacterium xenopi* pulmonary infection in an HIV infected patient under highly active antiretroviral treatment, **60**
- immunisation**, Hepatitis B virus infection among Chinese residents in the United Kingdom, **166**
- immunotherapy**, Adult and paediatric contact immunotherapy with squaric acid dibutylester (SADBE) for recurrent, multiple, resistant, mucocutaneous anogenital warts, **309**
- India**, Diverse realities: sexually transmitted infections and HIV in India, **i31**
- Human T lymphotropic virus-1 (HTLV-1) infection in patients with unclassifiable dermatitis in central Kerala, south India: a preliminary study, **e7**
- Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**
- Syndromic management of sexually transmitted diseases at primary care level, Mozambique, **67**
- Indian Ocean**, STI epidemics in the Indian Ocean region: Can the phase be assessed, **i121**
- infectious disease**, The criminalisation of HIV transmission, **448**
- injection drug user**, Risky sexual behaviours among injection drugs users with high HIV prevalence: implications for STD control, **i170**
- intervention programme**, Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Short term interventions are not enough, **80**
- intervention study**, Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**
- intrathoracic lymphadenopathy**, Progressive intrathoracic lymphadenopathy: EBV associated non-Hodgkin's lymphoma, **13**
- intrauterine contraceptive**, An update on pelvic inflammatory disease, **18**
- Iran**, Characteristics of gonorrhoea in Kermanshah, Iran, **460**
- journalology**, The curtain call, **392**
- Kaposi's sarcoma**, Kaposi's sarcoma infiltrating skeletal muscle, **464**
- Kenya**, Part time female sex workers in a suburban community in Kenya: a vulnerable hidden population, **271**
- Response of a sexually transmitted infection epidemic to a treatment and prevention programme in Nairobi, Kenya, **i114**
- knowledge**, Knowledge about the prevention of sexually transmitted diseases: a longitudinal study of young women from 16–23 years of age, **339**
- laboratory techniques**, Polymerase chain reaction for diagnosis of genital herpes in a genitourinary medicine clinic, **21**
- lactic acidosis**, Acute onset lactic acidosis and pancreatitis in the third trimester of pregnancy in HIV-1 positive women taking antiretroviral medication, **58**
- Lactic acidosis and abnormal liver function in advanced HIV disease, **139**
- legalisation**, Will the legalisation of street sex work improve health?, **309**
- ligase chain reaction**, Epidemiology and natural history of ligase chain reaction detected chlamydial and gonococcal infections, **208**
- linked surveys**, The feasibility of integrated STI prevalence and behaviour surveys in developing countries, **187**
- lipodystrophy**, Does increased aromatase activity in adipose fibroblasts cause low sexual desire in patients with HIV lipodystrophy?, **64**
- liver function**, Lactic acidosis and abnormal liver function in advanced HIV disease, **139**
- London**, News from the frontline: sexually transmitted infections in teenagers attending a genitourinary clinic in south east London, **349**
- Pathways to HIV testing and care by black African and white patients in London, **37**
- lung**, Multicentric plasma cell variant Castleman's disease mimicking intrapulmonary malignancy, **304**
- lymphogranuloma venereum**, Bilateral groove sign with penoscrotal elephantiasis, **458**
- Lymphogranuloma venereum, **90**
- malignancy**, Multicentric plasma cell variant Castleman's disease mimicking intrapulmonary malignancy, **304**
- management**, Outreach health adviser in a community clinic screening programme improves management of genital chlamydia infection, **101**
- Manitoba**, Gonorrhoea and chlamydia core groups and sexual networks in Manitoba, **i145**
- marijuana**, Predictors of infection with *Trichomonas vaginalis*: a prospective study of low income African-American adolescent females, **360**
- mass screening**, Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men, **190**
- mathematical model**, An introduction to mathematical models in sexually transmitted disease epidemiology, **7**
- The impact of HIV epidemic phases on the effectiveness of core group interventions: insights from mathematical models, **i78**
- men**, Risk factors for genital HPV DNA in men resemble those found in women: a study of male attendees at a Danish STD clinic, **215**
- migrants**, HIV infection among people of foreign origin voluntarily tested in Spain. A comparison with national subjects, **250**
- military installation**, Geographic epidemiology of gonorrhoea and chlamydia on a large military installation: application of a GIS system, **40**
- molecular epidemiology**, Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**
- Mongolia**, Hepatitis, syphilis, and HIV sentinel surveillance in Mongolia 1999–2000, **223**
- Mozambique**, Syndromic management of sexually transmitted diseases at primary care level, **e2**
- MSM**, Current characteristics of male gonorrhoea outpatients in Athens, Greece, **e5**
- Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men, **190**
- Mycobacterium xenopi**, *Mycobacterium xenopi* pulmonary infection in an HIV infected patient under highly active antiretroviral treatment, **60**
- Mycoplasma genitalium**, Coexistence of urethritis with genital ulcer disease in South Africa: influence on provision of syndromic management, **274**
- Mycoplasma genitalium* is not associated with adverse outcomes of pregnancy in Guinea-Bissau, **289**
- national sexual health strategy**, The national strategy for sexual health and HIV: implications for genitourinary medicine, **83**

- natural history**, Epidemiology and natural history of ligase chain reaction detected chlamydial and gonococcal infections, **208**
- Neisseria gonorrhoeae***, Detection of *Neisseria gonorrhoeae* by PCR using *orf1* gene as target, **72**  
Inhibition of *Neisseria gonorrhoeae* by vaginal lubricants, **468**  
Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**  
Detection of *Neisseria gonorrhoeae* by PCR using *orf1* gene as target, **72**
- Nepal**, The practice of STI treatment among chemists and druggists in Pokhara, Nepal, **223**
- neuropsychiatric reaction**, Neuropsychiatric reaction induced by clarithromycin, **75**
- nevirapine**, Nevirapine + efavirenz based salvage therapy in heavily pretreated HIV infected patients, **72**
- newborn**, The potential role of suppressive therapy for sex partners in the prevention of neonatal herpes: a health economic analysis, **425**
- non-gonococcal urethritis**, *Chlamydia trachomatis* in non-gonococcal urethritis patients and their heterosexual partners: routine testing by polymerase chain reaction, **286**
- non-Hodgkin's lymphoma**, Progressive intrathoracic lymphadenopathy: EBV associated non-Hodgkin's lymphoma, **13**
- non-occupational post-exposure prophylaxis**, Post-exposure prophylaxis for non-occupational exposure to HIV: current clinical practice and opinions in the UK, **130**
- non-specific urethritis**, Characteristics of adenovirus associated urethritis, **445**
- novel mutations**, Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**
- nucleic acid amplification**, *Chlamydia trachomatis* diagnostics, **232**
- nurse led care**, A postal survey to identify and describe nurse led clinics in genitourinary medicine services across England, **98**  
Sexual health clinics for women led by specialist nurses or senior house officers in a central London GUM service: a randomised controlled trial, **93**
- nursing**, The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 1 Role, **292**  
The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 2 Education and training, **298**
- obituary**, James Oliver Doyle, MB BCH, BAO 1943, LM 1944 Dublin, MRCP 1948, MD 1948 NUI (1918–2001), **154**  
Robert Steel Morton 1917–2002, **322**  
Sydney M Laird, BSc, MD, DPH, FRCP (1911–2001), **20**
- oestrogen receptors**, Oestrogen receptor transcripts associated with cervical human papillomavirus infection, **210**
- oestrogen**, Does increased aromatase activity in adipose fibroblasts cause low sexual desire in patients with HIV lipodystrophy?, **64**
- oophoritis**, An update on pelvic inflammatory disease, **18**
- orf1* gene** Detection of *Neisseria gonorrhoeae* by PCR using *orf1* gene as target, **72**
- pancreatitis**, Acute onset lactic acidosis and pancreatitis in the third trimester of pregnancy in HIV-1 positive women taking antiretroviral medication, **58**
- parC* gene** Mutation patterns in *gyrA* and *parC* genes of ciprofloxacin resistant isolates of *Neisseria gonorrhoeae* from India, **440**
- partner notification**, Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**  
Investigation into the acceptability and effectiveness of a new contact slip in the management of *Chlamydia trachomatis* at a London genitourinary medicine clinic, **422**  
Sex difference in partner notification: results from three population based surveys in France, **45**
- peer education**, Peer led HIV prevention among homosexual men in Britain, **158**
- pelvic inflammatory disease**, Cost effectiveness of screening for *Chlamydia trachomatis*: a review of published studies, **406**
- penoscrotal elephantiasis**, Bilateral groove sign with penoscrotal elephantiasis, **458**
- perianal abscess**, Gonococcal perianal abscess: re-emergence after cessation of co-trimoxazole, **389**
- perianal warts**, Perianal Crohn's disease masquerading as perianal warts, **302**
- peripheral neuropathy**, Peripheral neuropathy in patients with HIV infection: consider dual pathology, **462**
- pH dependent colonisation**, Vaginal infection by *Enterobacter sakazakii*, **467**
- phase specific strategies**, Heterosexual outbreak of infectious syphilis: epidemiological and ethnographic analysis and implications for control, **i164**
- polymerase chain reaction**, Contamination of environmental surfaces by genital human papillomaviruses, **135**  
Detection of *Neisseria gonorrhoeae* by PCR using *orf1* gene as target, **72**  
The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice, **278**
- population**, Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**
- postal survey**, A postal survey to identify and describe nurse led clinics in genitourinary medicine services across England, **98**
- pregnancy**, 15 Month follow up of African children following vaginal cleansing with benzalkonium chloride of their HIV infected mothers during late pregnancy and delivery, **267**  
Acute onset lactic acidosis and pancreatitis in the third trimester of pregnancy in HIV-1 positive women taking antiretroviral medication, **58**  
*Mycoplasma genitalium* is not associated with adverse outcomes of pregnancy in Guinea-Bissau, **289**
- prenatal intervention**, Congenital syphilis—missed opportunities for prenatal intervention, **224**
- prevention strategies**, Phase specific strategies for the prevention, control, and elimination of sexually transmitted infections: case study in Lambeth, Southwark, and Lewisham, London, UK, **i133**
- prevention**, Canadian STI national goals and phase specific strategies, **i189**  
Phase specific strategies for the prevention, control, and elimination of sexually transmitted diseases: background country profile, England and Wales, **i125**  
Populations, pathogens, and epidemic phases: closing the gap between theory and practice in the prevention of sexually transmitted diseases, **i183**  
Understanding correlates of hepatitis B virus vaccination in men who have sex with men: what have we learned?, **374**
- prostatic tuberculosis**, Prostatic tuberculosis in an HIV infected male, **147**
- prostitution**, Characteristics of gonorrhoea in Kermanshah, Iran, **460**
- public health**, Behavioural surveillance: the value of national coordination, **398**  
From community to clients: the professionalisation of HIV prevention among gay men and its implications for intervention selection, **i176**  
HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**  
Populations, pathogens, and epidemic phases: closing the gap between theory and practice in the prevention of sexually transmitted diseases, **i183**  
Sex difference in partner notification: results from three population based surveys in France, **45**
- qualitative research methods**, The application of qualitative research methods to the study of sexually transmitted infections, **87**
- race**, HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- randomised controlled trials**, Sexual health clinics for women led by specialist nurses or senior house officers in a central London GUM service: a randomised controlled trial, **93**
- rapid plasmin reagin**, Performance of the rapid plasma reagin and the rapid syphilis screening tests in the diagnosis of syphilis in field conditions in rural Africa, **282**
- rapid syphilis test**, Performance of the rapid plasma reagin and the rapid syphilis screening tests in the diagnosis of syphilis in field conditions in rural Africa, **282**
- recreational drug use**, Recreational drug use by GUM clinic attendees, **310**
- repeat testing**, Resolution of the recent performance problem of Abbott LCx *Chlamydia trachomatis* assay Issues of repeat testing for confirmation of chlamydial infection, **225**
- reproductive health**, Sexual and reproductive health among female adolescents: preliminary results, **386**
- reproductive morbidity**, Syndromic management of sexually transmitted diseases at primary care level, Mozambique, **67**
- risk factors**, Clinical characteristics of *Chlamydia trachomatis* infections in a general outpatient department of obstetrics and gynaecology in the Netherlands, **e6**  
Risk factors for genital HPV DNA in men resemble those found in women: a study of male attendees at a Danish STD clinic, **215**
- salpingitis**, An update on pelvic inflammatory disease, **18**
- schistosomiasis**, Immune reconstitution eosinophilia due to schistosomiasis, **76**
- screening**, Anti-HIV serology in patients with sexual dysphoria in screening test before sex change surgery, **75**  
*Chlamydia* screening and sexual health, **396**  
Comparing cost effectiveness of screening women for *Chlamydia trachomatis* in systematic and opportunistic approaches, **73**  
Cost effectiveness of screening for *Chlamydia trachomatis*: a review of published studies, **406**

- Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- Management of screened chlamydia positive women, **155**
- Prevalence of and risk indicators for STIs among women seeking induced abortions in two urban family planning clinics in Shandong province, People's Republic of China, **e3**
- Provider willingness to screen all sexually active adolescents for chlamydia, **369**
- Screening for STIs in individuals with HIV infection, **387**
- The national strategy for sexual health and HIV: implications for genitourinary medicine, **83**
- The value of mass screening for chlamydia control in high prevalence communities, **i64**
- screening programme**, Cost effectiveness analysis of a population based screening programme for asymptomatic *Chlamydia trachomatis* infection in women, **76**
- self treatment**, Vulvovaginitis due to fluconazole resistant *Candida albicans* following self treatment with non-prescribed triazoles, **308**
- semen**, Four layer discontinuous gradient for HIV, **388**
- sensitivity**, An evidence based approach to testing for antibody to herpes simplex virus type 2, **430**
- seroconverters**, Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**
- seroprevalence**, Seroprevalence of HSV-1 and HSV-2 infection in the general French population, **201**
- serotesting**, Genital herpes: general practitioners' knowledge and opinions, **198**
- sex change**, Anti-HIV serology in patients with sexual dysphoria in screening test before sex change surgery, **75**
- sex worker**, Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Short term interventions are not enough, **80**
- sexual abuse**, Adolescent sexual health in Sweden, **352**
- sexual behaviour**, Behavioural surveillance: the value of national coordination, **398**
- HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- Part time female sex workers in a suburban community in Kenya: a vulnerable hidden population, **271**
- Post-treatment sexual and prevention behaviours of adolescents with sexually transmitted infections, **365**
- Prevention interventions for HIV positive individuals, **393**
- Risky sexual behaviours among injection drugs users with high HIV prevalence: implications for STD control, **i170**
- Seroprevalence of hepatitis A immunity in male genitourinary medicine clinic attenders: a case control study of heterosexual and homosexual men, **174**
- The accuracy of reported sensitive sexual behaviour in Britain: exploring the extent of change 1990–2000, **26**
- The effect of HIV, behavioural change, and STD syndromic management on STD epidemiology in sub-Saharan Africa: simulations of Uganda, **i55**
- sexual conversations**, Assisting medical students to conduct empathic conversations with patients from a sexual medicine clinic, **246**
- sexual desire**, Does increased aromatase activity in adipose fibroblasts cause low sexual desire in patients with HIV lipodystrophy?, **64**
- sexual dysphoria**, Anti-HIV serology in patients with sexual dysphoria in screening test before sex change surgery, **75**
- sexual health**, Assisting medical students to conduct empathic conversations with patients from a sexual medicine clinic, **246**
- Chlamydia screening and sexual health, **396**
- HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- Sexual and reproductive health among female adolescents: preliminary results, **386**
- Young people's views on provision of sexual health services, **342**
- sexual health issues**, HIV negative certification and sexual health issues facing performers in the adult entertainment industry in the UK, **311**
- sexual network**, Heterosexual outbreak of infectious syphilis: epidemiological and ethnographic analysis and implications for control, **i164**
- Risk network structure in the early epidemic phase of HIV transmission in Colorado Springs, **i159**
- Sexual network structure as an indicator of epidemic phase, **i152**
- sexual transmission**, Prevalence and incidence of hepatitis B virus infection in STD clinic attendees in Pune, India, **169**
- Sexually acquired hepatitis, **235**
- sexually transmitted disease**, An introduction to mathematical models in sexually transmitted disease epidemiology, **7**
- Attitudes about sexual disclosure and perceptions of stigma and shame, **334**
- Canadian STI national goals and phase specific strategies, **i189**
- Characteristics of gonorrhoea in Kermanshah, Iran, **460**
- Decreasing STD incidence and increasing condom use among Chinese sex workers following a short term intervention: a prospective cohort study, **110**
- Determinants of STD epidemics: implications for phase appropriate intervention strategies, **i3**
- Diverse realities: sexually transmitted infections and HIV in India, **i31**
- Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**
- Geographical variations in the epidemiology of bacterial sexually transmitted infections in Manitoba, Canada, **i139**
- Hepatitis, syphilis, and HIV sentinel surveillance in Mongolia 1999–2000, **223**
- Knowledge about the prevention of sexually transmitted diseases: a longitudinal study of young women from 16–23 years of age, **339**
- News from the frontline: sexually transmitted infections in teenagers attending a genitourinary clinic in south east London, **349**
- Phase specific strategies for the prevention, control, and elimination of sexually transmitted diseases: background country profile, England and Wales, **i125**
- Populations, pathogens, and epidemic phases: closing the gap between theory and practice in the prevention of sexually transmitted diseases, **i183**
- Prevalence and incidence of hepatitis B virus infection in STD clinic attendees in Pune, India, **169**
- Prevalence of and risk indicators for STIs among women seeking induced abortions in two urban family planning clinics in Shandong province, People's Republic of China, **e3**
- Reasons for not using condoms among the Hong Kong Chinese population: implications for HIV and STD prevention, **180**
- Revised UK national guidelines on sexually transmitted infections and closely related conditions 2002, **81**
- Role of core and bridging groups in the transmission dynamics of HIV and STIs in Cotonou, Benin, West Africa, **i69**
- Seroprevalence of hepatitis A immunity in male genitourinary medicine clinic attenders: a case control study of heterosexual and homosexual men, **174**
- Sex difference in partner notification: results from three population based surveys in France, **45**
- Sex in the city: sexual behaviour, societal change, and STDs in Saigon, **i47**
- Sexual network structure as an indicator of epidemic phase, **i152**
- Short term interventions are not enough, **80**
- Sociodemographic context of the AIDS epidemic in a rural area in Tanzania with a focus on people's mobility and marriage, **i97**
- STD/HIV intervention and research programme Mwanza Region, NW Tanzania, **i91**
- STD in Bangladesh's trucking industry: prevalence and risk factors, **31**
- Syndromic management of sexually transmitted diseases at primary care level, **e2**
- The geographical and temporal evolution of sexually transmitted disease epidemics, **i14**
- The profile and context of the epidemics of sexually transmitted infections including HIV in Zimbabwe, **i40**
- The value of mass screening for chlamydia control in high prevalence communities, **i64**
- Update of trichomoniasis, **378**
- sexually transmitted infection**, A pilot study of a rapid assessment method to identify places for AIDS prevention in Cape Town, South Africa, **i106**
- A survey of STI policies and programmes in Europe: preliminary results, **380**
- Adolescent sexual health in Sweden, **352**
- Behavioural surveillance: the value of national coordination, **398**
- Detection of *Chlamydia trachomatis* in an Australian high school student population, **194**
- Donovanosis, **452**
- Epidemiology and natural history of ligase chain reaction detected chlamydial and gonococcal infections, **208**
- Genital ulcers, stigma, HIV, and STI control in sub-Saharan Africa, **143**
- Health seeking and sexual behaviour in patients with sexually transmitted infections: the importance of traditional healers in Thyolo, Malawi, **127**
- HIV epidemicity in context of STI declines: a telling discordance, **467**
- Human T lymphotropic virus-I (HTLV-I) infection in patients with unclassifiable dermatitis in central Kerala, south India: a preliminary study, **e7**
- National guideline for the management of suspected sexually transmitted infections in children and young people, **324**
- Phase specific strategies for the prevention, control, and elimination of sexually transmitted infections: case study in Lambeth, Southwark, and Lewisham, London, UK, **i133**
- Prevalence and risk factors of sexually transmitted infections and cervical neoplasia in women's health clinics in Nicaragua, **204**

- Prevalence and risk indicators for STIs among women seeking induced abortions in two urban family planning clinics in Shandong province, People's Republic of China, **e3**
- Prevention interventions for HIV positive individuals, **393**
- Reaching homeless youths for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* screening in Denver, Colorado, **357**
- Response of a sexually transmitted infection epidemic to a treatment and prevention programme in Nairobi, Kenya, **i114**
- Revised UK national guidelines on sexually transmitted infections and closely related conditions 2002, **81**
- Screening for STIs in individuals with HIV infection, **387**
- Services for sexually transmitted infections in Europe and central Asia, **320**
- Sexually transmitted infections in Bangladeshi resident in the UK: a case-control study, **120**
- STI epidemics in the Indian Ocean region: can the phase be assessed?, **i121**
- The application of qualitative research methods to the study of sexually transmitted infections, **87**
- The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice, **278**
- The evolution of sexually transmitted infections in the Ukraine, **219**
- The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 1 Role, **292**
- The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 2 Education and training, **298**
- The impact of the phase of an epidemic of sexually transmitted infection on the evolution of the organism, **i20**
- The practice of STI treatment among chemists and druggists in Pokhara, Nepal, **223**
- Understanding racial-ethnic and societal differentials in STI, **2**
- Uptake of HIV testing in patients with a confirmed sexually transmitted infection, **389**
- Sexually Transmitted Infections journal**, The year ahead, **6**
- shame**, Attitudes about sexual disclosure and perceptions of stigma and shame, **334**
- simulation models**, The effect of HIV, behavioural change, and STD syndromic management on STD epidemiology in sub-Saharan Africa: simulations of Uganda, **i55**
- skeletal muscle**, Kaposi's sarcoma infiltrating skeletal muscle, **464**
- social epidemiology**, Understanding racial-ethnic and societal differentials in STI, **2**
- social networks**, Understanding racial-ethnic and societal differentials in STI, **2**
- South Africa**, Evaluation of a video based health education strategy to improve sexually transmitted disease partner notification in South Africa, **53**
- Spain**, Factors influencing HIV progression in a seroconverter cohort in Madrid from 1985 to 1999, **255**
- HIV infection among people of foreign origin voluntarily tested in Spain. A comparison with national subjects, **250**
- Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**
- specificity**, An evidence based approach to testing for antibody to herpes simplex virus type 2, **430**
- squaric acid dibutylester**, Adult and paediatric contact immunotherapy with squaric acid dibutylester (SADBE) for recurrent, multiple, resistant, mucocutaneous anogenital warts, **309**
- STI programmes**, A survey of STI policies and programmes in Europe: preliminary results, **380**
- Services for sexually transmitted infections in Europe and central Asia, **320**
- stigma**, Attitudes about sexual disclosure and perceptions of stigma and shame, **334**
- Genital ulcers, stigma, HIV, and STI control in sub-Saharan Africa, **143**
- STI/HIV services**, The national strategy for sexual health and HIV: implications for genitourinary medicine, **83**
- street sex work**, Will the legalisation of street sex work improve health?, **309**
- sub-Saharan Africa**, Genital ulcers, stigma, HIV, and STI control in sub-Saharan Africa, **143**
- subtypes**, Prevalence of different HIV-1 subtypes in an urban clinic in Madrid, **e1**
- surveillance**, Behavioural surveillance: the value of national coordination, **398**
- HIV testing and high risk sexual behaviour among London's migrant African communities: a participatory research study, **241**
- Sweden**, Adolescent sexual health in Sweden, **352**
- Knowledge about the prevention of sexually transmitted diseases: a longitudinal study of young women from 16–23 years of age, **339**
- syndromic management**, Coexistence of urethritis with genital ulcer disease in South Africa: influence on provision of syndromic management, **274**
- syphilis**, Heterosexual outbreak of infectious syphilis: epidemiological and ethnographic analysis and implications for control, **i164**
- Low incidence of syphilis among factory workers in Ethiopia: effect of an intervention based on education and counselling, **123**
- Performance of the rapid plasma reagin and the rapid syphilis screening tests in the diagnosis of syphilis in field conditions in rural Africa, **282**
- Resurgence of syphilis in England, **308**
- STD/HIV intervention and research programme Mwanza Region, NW Tanzania, **i91**
- Tanzania**, Community effects on the risk of HIV infection in rural Tanzania, **261**
- Sociodemographic context of the AIDS epidemic in a rural area in Tanzania with a focus on people's mobility and marriage, **i97**
- STD/HIV intervention and research programme Mwanza Region, NW Tanzania, **i91**
- teenage abortion**, Adolescent sexual health in Sweden, **352**
- teenage pregnancy**, Adolescent reproductive health interventions, **315**
- termination of pregnancy**, Provision of chlamydia testing in a nationwide service offering termination of pregnancy: with data capture to monitor prevalence of infection, **416**
- the Netherlands**, Clinical characteristics of *Chlamydia trachomatis* infections in a general outpatient department of obstetrics and gynaecology in the Netherlands, **e6**
- traditional healers**, Health seeking and sexual behaviour in patients with sexually transmitted infections: the importance of traditional healers in Thyolo, Malawi, **127**
- triazole resistance**, Vulvovaginitis due to fluconazole resistant *Candida albicans* following self treatment with non-prescribed triazoles, **308**
- Trichomonas vaginalis**, TV or not TV?, **185**
- trichomoniasis**, Predictors of infection with *Trichomonas vaginalis*: a prospective study of low income African-American adolescent females, **360**
- Update of trichomoniasis, **378**
- tropical medicine**, Donovanosis, **452**
- Lymphogranuloma venereum, **90**
- Sexually acquired hepatitis, **235**
- truck drivers**, STD in Bangladesh's trucking industry: prevalence and risk factors, **31**
- tuberculosis**, Tuberculosis of the cervix: case presentation and a review of the literature, **62**
- Uganda**, The effect of HIV, behavioural change, and STD syndromic management on STD epidemiology in sub-Saharan Africa: simulations of Uganda, **i55**
- Ukraine**, The evolution of sexually transmitted infections in the Ukraine, **219**
- United Kingdom**, HIV negative certification and sexual health issues facing performers in the adult entertainment industry in the UK, **311**
- Revised UK national guidelines on sexually transmitted infections and closely related conditions 2002, **81**
- Sexually transmitted infections in Bangladeshi resident in the UK: a case-control study, **120**
- The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 1 Role, **292**
- The historical role and education of nurses for the care and management of sexually transmitted infections in the United Kingdom: 2 Education and training, **298**
- urban genitourinary medicine clinic**, Factors affecting co-infection with genital chlamydia and genital gonorrhoea in an urban genitourinary medicine clinic, **387**
- urban slum**, Syndromic management of sexually transmitted diseases at primary care level, Mozambique, **67**
- urethritis**, Characteristics of adenovirus associated urethritis, **445**
- Coexistence of urethritis with genital ulcer disease in South Africa: influence on provision of syndromic management, **274**
- urine specimens**, Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- urticaria**, Successful use of valciclovir in a case of recurrent urticaria associated with genital herpes, **468**
- vaccination**, Preventive human papillomavirus vaccination, **4**
- Understanding correlates of hepatitis B virus vaccination in men who have sex with men: what have we learned?, **374**
- vaginal infection**, Vaginal infection by *Enterobacter sakazakii*, **467**
- vaginal lubricants**, Inhibition of *Neisseria gonorrhoeae* by vaginal lubricants, **468**
- vaginal microbicide**, 15 Month follow up of African children following vaginal cleansing with benzalkonium chloride of their HIV infected mothers during late pregnancy and delivery, **267**
- vaginal smears**, Validation of a simplified grading of Gram stained vaginal smears for use in genitourinary medicine clinics, **413**

- vaginal swabs**, The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice, **278**
- vaginitis emphysematosa**, Vaginitis emphysematosa, **155**
- valaciclovir**, Aborted genital herpes simplex virus lesions: findings from a randomised controlled trial with valaciclovir, **435**  
Successful use of valaciclovir in a case of recurrent urticaria associated with genital herpes, **468**
- VD regulations**, Managing the very young patient: a conflict between the requirements of the Children Act and the VD regulations?, **332**
- Vietnam**, Sex in the city: sexual behaviour, societal change, and STDs in Saigon, **i47**
- viral hepatitis**, Sexually acquired hepatitis, **235**
- women**, Comparing cost effectiveness of screening women for *Chlamydia trachomatis* in systematic and opportunistic approaches, **73**  
Cost effectiveness analysis of a population based screening programme for asymptomatic *Chlamydia trachomatis* infection in women, **76**
- Knowledge about the prevention of sexually transmitted diseases: a longitudinal study of young women from 16–23 years of age, **339**
- Major improvements in cost effectiveness of screening women for *Chlamydia trachomatis* using pooled urine specimens and high performance testing, **74**
- Management of screened chlamydia positive women, **155**  
*Chlamydia trachomatis* infection among sexually active young women in Italy, **e4**
- women's health**, Prevalence and risk factors of sexually transmitted infections and cervical neoplasia in women's health clinics in Nicaragua, **204**
- young people**, Young people's views on provision of sexual health services, **342**
- Zimbabwe**, The profile and context of the epidemics of sexually transmitted infections including HIV in Zimbabwe, **i40**