

Subject index

- access**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
Treating sexually transmitted infections in primary care: a missed opportunity?, **134**
- accident and emergency department**, Urine based screening for asymptomatic/undiagnosed genital chlamydial infection in young people visiting the accident and emergency department is feasible, acceptable, and can be epidemiologically helpful, **229**
- adolescence**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- adolescent health**, Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- adolescents**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- adult respiratory distress syndrome**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- Africa**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
Sexually transmitted infections in Africa: single dose treatment is now affordable, **432**
- age**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- AIDS**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
Television coverage of AIDS and transmission of the AIDS virus, **170**
- AIDS/STD**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- amphotericin B**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
- antenatal syphilis screening**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- antibiotics**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
- antibodies**, Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, **351**
- antiretroviral therapy**, Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- appointments**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
- attitude**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- Australia**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, **332**
- azithromycin**, Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, **453**
- bacterial vaginosis**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- behaviour**, Determinants of disclosure of genital herpes to partners, **42**
- Belgium**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- bisexuality**, Sexual behaviour of lesbians and bisexual women, **147**
- black township**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- Botswana**, Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
- Brazil**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- Britain**, Male circumcision in Britain: findings from a national probability sample survey, **499**
- Cambodia**, Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Candida albicans**, Vulvovaginal carriage of yeasts other than *Candida albicans*, **249**
- Candida glabrata**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
Vulvovaginal carriage of yeasts other than *Candida albicans*, **249**
- Castleman's disease**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
- cutaneous larva migrans**, Cutaneous larva migrans of the penis, **500**
- ceftriaxone**, Treatment of neurosyphilis with ceftriaxone, **415**
- cervical cancer**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, **191**
- cervical screening**, Awareness of human papillomavirus among women attending a well woman clinic, **320**
- cervix**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- cervix uterus**, Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- chancere**, Perforating chancre: any cause-effect relation with HIV infection?, **262**
- chancroid**, Chancroid: clinical manifestations, diagnosis, and management, **68**
Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, **294**
- chaperone**, Chaperoning in genitourinary medicine clinics, **422**
- chaperones**, Chaperoning in genitourinary medicine: supporting patients and protecting doctors, **498**
- chlamydia**, Chlamydia testing before termination of pregnancy, **352**
Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
Recent pilot studies of chlamydia screening, **352**
Repeat chlamydia screening by mail, San Francisco, **28**
Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
Validity and utility of screening tests for STIs, **356**
- chlamydia screening**, Urine based screening for asymptomatic/undiagnosed genital chlamydial infection in young people visiting the accident and emergency department is feasible, acceptable, and can be epidemiologically helpful, **229**

- Chlamydia trachomatis**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
- Are all genital *Chlamydia trachomatis* infections pathogenic? A study in men, **349**
- Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
- Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
- Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- Spending money to save money, **4**
- chlamydial infection**, Chlamydia screening in the United Kingdom, **3**
- First, do not harm: also an issue in NAA assay diagnostics for chlamydial infection, **76**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- circumcision**, Circumcision practice patterns in South Korea: community based survey, **65**
- HIV and circumcision: new factors to consider, **495**
- Issues associated with the introduction of circumcision into a non-circumcising society, **427**
- Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
- clinical practice variations**, Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- cognitions**, Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- cohort study**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- Colombia**, Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- computerised questionnaire**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- condom negotiation skills**, Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- condoms**, Don't overlook condoms for HIV prevention, **268**
- congenital syphilis**, Congenital syphilis in the Russian Federation: magnitude, determinants, and consequences, **106**
- continuing medical education**, Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- cost effectiveness**, A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
- cost effectiveness analysis**, Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Spending money to save money, **4**
- Costa Rica**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- cost-effectiveness analysis**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- counselling**, Genital herpes serotyping: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- couple transmission**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- cranial tuberculomas**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- Cryptosporidium**, Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- cytokines**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- cytolethal distending toxin**, Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, **294**
- DALYs**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- dark ground microscopy**, Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- delay**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
- demographic risk factors**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- diagnosis**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- diagnostics**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- doctors**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- drug resistance**, New drugs for treating drug resistant HIV-1, **176**
- dual neoplastic diseases**, Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- education**, Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- education initiatives**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- EIA amplicon detection**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
- England**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- environmental contamination**, Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
- epidemiology**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- Estonia**, Seroprevalence study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- ethnicity**, Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- Europe**, A survey of STI policies and programmes in Europe, **171**
- family planning**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- female bar workers**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- female circumcision**, Female genital mutilation in the Sudan: survey of the attitude of Khartoum university students towards this practice, **220**
- female genital mutilation**, Female genital mutilation in the Sudan: survey of the attitude of Khartoum university students towards this practice, **220**
- female sex workers**, Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- females**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**

- flucytosine**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
- follow up**, Patients lost to follow up: experience of an HIV clinic, **349**
- foreskin**, Circumcision practice patterns in South Korea: community based survey, **65**
- general practice**, Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- genital chlamydia infection**, Who is being tested for genital chlamydia in primary care?, **234**
- genital diseases**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**
- genital herpes**, Determinants of disclosure of genital herpes to partners, **42**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Genital herpes serotyping: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- Is HSV serology useful for the management of first episode genital herpes?, **276**
- genital infection**, Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
- genital ulcer disease**, Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- genital warts**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- genital warts, circumcision, treatment**, Circumcision in genital warts—let us not forget!, **265**
- genitourinary diseases**, Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- genitourinary infection**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
- genitourinary infections**, Self obtained samples for genitourinary infections, **171**
- genitourinary medicine**, Chaperoning in genitourinary medicine: supporting patients and protecting doctors, **498**
- genitourinary medicine clinics**, Chaperoning in genitourinary medicine clinics, **74**
- Raising awareness of UK GUM clinic activities, **264**
- genitourinary medicine education**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- genotypes**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- Germany**, Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, **351**
- glucocorticoids**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- gold mining**, Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- gonorrhoea**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
- Gonorrhoea reinfection in heterosexual STD clinic attendees: longitudinal analysis of risks for first reinfection, **124**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- Pharyngeal gonorrhoea: the forgotten reservoir, **345**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- The fall and rise of gonorrhoea incidence in Israel: an international phenomenon?, **116**
- Greater Manchester**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- GUM clinics**, Maintaining patient access to GUM clinics: is it compatible with appointments?, **11**
- GUM services**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- gynaecological examination**, The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- HAART**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- Polymorphic immune restoration syndrome after effective HAART, **171**
- Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- Haemophilus ducreyi**, Chancroid: clinical manifestations, diagnosis, and management, **68**
- healthcare settings**, Chlamydia screening in the United Kingdom, **3**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- hepatitis B**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- hepatitis C**, Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- hepatitis C testing**, Hepatitis C testing in HIV infected patients, **76**
- hepatitis C virus**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**
- Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- herpes simplex virus**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Epidemiology of recurrent genital herpes simplex virus types 1 and 2, **456**
- Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- herpes simplex virus serology**, Is HSV serology useful for the management of first episode genital herpes?, **276**
- herpes simplex virus seroprevalence**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- HIV**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals, **137**
- Don't overlook condoms for HIV prevention, **268**
- Enhanced risk of HIV sexual transmission during structured treatment interruption, **74**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Hepatitis C testing in HIV infected patients, **76**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**

- HIV and circumcision: new factors to consider, **495**
- HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Lower motor neuron syndrome and HIV infection, **351**
- Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- Male circumcision: an acceptable strategy for HIV prevention in Botswana, **214**
- New drugs for treating drug resistant HIV-1, **176**
- Perforating chancres: any cause-effect relation with HIV infection?, **262**
- Polymyositis masquerading as mitochondrial toxicity, **417**
- Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- Superior mesenteric artery syndrome in an HIV positive patient, **262**
- The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, **453**
- Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- View 1: Assisted conception in couples with HIV infection, **185**
- Commissioning HIV services, **84**
- View 2: Assisted conception in couples with HIV infection, **189**
- HIV clinic**, Patients lost to follow up: experience of an HIV clinic, **349**
- HIV infection**, Emerging of dual AIDS associated neoplastic diseases in the era of highly active antiretroviral therapy, **345**
- Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
- Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- What's new for injection drug users with HIV infection?, **80**
- HIV prevention**,
- HIV testing attitudes**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- HIV transmission risks**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- HIV treatment adherence**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- HIV/AIDS epidemic**, The HIV/AIDS epidemic in Ukraine: stable or still exploding?, **263**
- HIV/AIDS**, Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- homosexual men**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- homosexuality**, Sexual behaviour of lesbians and bisexual women, **147**
- hospital clinic**, Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- human immunodeficiency virus**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- human papillomavirus**, Awareness of human papillomavirus among women attending a well woman clinic, **320**
- Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
- Interactive continuing medical education (CME) and its influence on the working practices of genitourinary clinicians, **350**
- Prevalence and determinants of *Chlamydia trachomatis* infections in women from Bogota, Colombia, **474**
- human parvovirus B19**, Primary human parvovirus B19 infection in an HIV infected patient on highly active antiretroviral therapy, **336**
- hygiene**, Circumcision practice patterns in South Korea: community based survey, **65**
- immune reconstitution**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
- income inequality**, Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- India**, Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
- indigenous people**, Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, **332**
- infertility**, View 1: Assisted conception in couples with HIV infection, **185**
- View 2: Assisted conception in couples with HIV infection, **189**
- injecting drug use**, Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- injection drug users**, What's new for injection drug users with HIV infection?, **80**
- internet**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- intimate examination**, Chaperoning in genitourinary medicine clinics, **422**
- Israel**, The fall and rise of gonorrhoea incidence in Israel: an international phenomenon?, **116**
- Haemophilus ducreyi**, Expression of the cytolethal distending toxin in a geographically diverse collection of *Haemophilus ducreyi* clinical isolates, **294**
- Sexually Transmitted Infections**, STI case management at a South African teaching hospital, **498**
- Taking up the reins, **2**
- Jacobson JM, editor Immunotherapy for Infectious Diseases**, Treatment of *Candida glabrata* using topical amphotericin B and flucytosine, **265**
- journalology**, Taking up the reins, **2**
- Kaposi's sarcoma**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
- knowledge**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- lactic acidosis**, Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
- lesbians**, Sexual behaviour of lesbians and bisexual women, **147**
- leucocyte esterase dipstick**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- ligase chain reaction**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- lipodystrophy**, A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals, **137**

- lymphocytes**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- lymphogranuloma venereum**, Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin, **453**
- male circumcision**, Male circumcision in Britain: findings from a national probability sample survey, **499**
- male clients**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- maternal syphilis**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- men**, Are all genital *Chlamydia trachomatis* infections pathogenic? A study in men, **349**
- Mexico**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- microscopy**, Poor sensitivity and consistency of microscopy in the diagnosis of low grade non-gonococcal urethritis, **487**
- middle aged men**, Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- middle aged women**, Clearance of HPV infection in middle aged men and women after 9 years' follow up, **348**
- mitochondrial toxicity**, Lactic acidosis in HIV infected patients: a systematic review of published cases, **340**
Polymyositis masquerading as mitochondrial toxicity, **417**
- mixed infection**, Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
- molecular typing**, Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
- monosymptomatic hypochondriacal psychosis**, Monosymptomatic hypochondriacal psychosis, **425**
- Morocco**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- mother to child transmission**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- motor neuron**, Lower motor neuron syndrome and HIV infection, **351**
- Mozambique**, Geographical focusing: an intervention to address increased risk for sexually transmitted diseases during repatriation and resettlement in post-war Mozambique, **77**
- MSM**, Evaluation of ligase chain reaction for the non-cultural detection of rectal and pharyngeal gonorrhoea in men who have sex with men, **484**
Risk factors leading to *Cryptosporidium* infection in men who have sex with men, **412**
- MSSVD**, The MSSVD, the National Sexual Health and HIV strategy for England and genitourinary medicine education, **166**
- multidrug resistance**, Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- muscle biopsy**, Polymyositis masquerading as mitochondrial toxicity, **417**
- Mycoplasma genitalium***, Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
- Mycoplasma* infections**, Tetracycline treatment does not eradicate *Mycoplasma genitalium*, **318**
- myopathy**, Polymyositis masquerading as mitochondrial toxicity, **417**
- Neisseria gonorrhoeae***, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
Detection of mixed infection of *Neisseria gonorrhoeae*, **56**
Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
- neurology**, Lower motor neuron syndrome and HIV infection, **351**
- neurosyphilis**, Treatment of neurosyphilis with ceftriaxone, **415**
- non-gonococcal urethritis**, Poor sensitivity and consistency of microscopy in the diagnosis of low grade non-gonococcal urethritis, **487**
- non-paying partners**, Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- nucleic acid amplification assay**, First, do not harm: also an issue in NAA assay diagnostics for chlamydial infection, **76**
- nurses**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- opportunistic testing**, Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- outreach programmes**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- outreach service**, Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- paediatric AIDS**, A pilot study to evaluate the safety and feasibility of the administration of AZT/3TC fixed dose combination to HIV infected pregnant women and their infants in Rio de Janeiro, Brazil, **448**
- Papanicolaou smear**, Was the Papanicolaou smear responsible for the decline of *Trichomonas vaginalis*?, **263**
- papillomavirus**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, **191**
- partner notification**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
Partner notification in primary care, **264**
- pelvic inflammatory disease**, Associations between *Mycoplasma genitalium*, *Chlamydia trachomatis*, and pelvic inflammatory disease, **154**
Diagnosis of pelvic inflammatory disease: time for a rethink, **491**
Spending money to save money, **4**
- penis**, Cutaneous larva migrans of the penis, **500**
- perianal**, Perianal verrucous epidermal naevus mimicking perianal warts, **424**
- perianal verrucous epidermal naevus**, Perianal verrucous epidermal naevus mimicking perianal warts, **424**
- perinatal mortality**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- perinatal transmission**, Selective transmission of multidrug resistant HIV to a newborn related to poor maternal adherence, **419**
- Peru**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- pharmacists**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- phimosi**, Circumcision practice patterns in South Korea: community based survey, **65**
- pneumonitis**, Lymphocytic interstitial pneumonitis in HIV infected adults, **88**
- podophyllin**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- podophyllotoxin**, Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts, **270**
- point of care diagnosis**, Validity and utility of screening tests for STIs, **356**
- polymerase chain reaction**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
Contamination of environmental surfaces by genital human papillomaviruses (HPV): a follow up study, **426**
Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
Increasing relative prevalence of HSV-2 infection among men with genital ulcers from a mining community in South Africa, **202**
Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
- polymorphic immune restoration syndrome**, Polymorphic immune restoration syndrome after effective HAART, **171**
- polymyositis**, Polymyositis masquerading as mitochondrial toxicity, **417**

- pontine myelinolysis**, Central pontine myelinolysis complicating treatment of multicentric Castleman's disease and Kaposi's sarcoma in a patient with AIDS, **179**
- population surveys**, LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- post-risk behaviours**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- postmenopausal woman**, Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- PPNG**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
- pregnancy**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- pretest counselling**, Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- primary care**, Partner notification in primary care, **264**
Surveillance of sexually transmitted infections in primary care, **174**
Treating sexually transmitted infections in primary care: a missed opportunity?, **134**
Who is being tested for genital chlamydia in primary care?, **234**
- priority setting**, Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- private practice**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- psychology**, Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- psychosocial impact**, Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- public health**, Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
Validity and utility of screening tests for STIs, **356**
- public understanding**, Awareness of human papillomavirus among women attending a well woman clinic, **320**
- qualitative assessment**, Psychosocial impact of serological diagnosis of herpes simplex virus type 2: a qualitative assessment, **280**
- randomised controlled trial**, Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- recurrent infection**, Repeat chlamydia screening by mail, San Francisco, **28**
- reinfection**, Gonorrhoea reinfection in heterosexual STD clinic attendees: longitudinal analysis of risks for first reinfection, **124**
- reproductive technology**, View I: Assisted conception in couples with HIV infection, **185**
View 2: Assisted conception in couples with HIV infection, **189**
- reproductive tract infections**, Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
The reliability of a structured examination protocol and self administered vaginal swabs: a pilot study of gynaecological outpatients in Goa, India, **251**
- risk behaviour**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- risk behaviours**, Avoiding risky sex partners: perception of partners' risks v partners' self reported risks, **197**
- risk factors**, Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- risk taking**, National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- Russian Federation**, Congenital syphilis in the Russian Federation: magnitude, determinants, and consequences, **106**
- sampling**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- screening**, A population based dynamic approach for estimating the cost effectiveness of screening for *Chlamydia trachomatis*, **426**
Chlamydia screening in the United Kingdom, **3**
- Chlamydial infection: an accurate model for opportunistic screening in general practice, **313**
- Estimating the costs and benefits of screening monogamous, heterosexual couples for unrecognised infection with herpes simplex virus type 2, **45**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- LCR testing for gonorrhoea and chlamydia in population surveys and other screenings of low prevalence populations: coping with decreased positive predictive value, **94**
- Managing partners of people diagnosed with *Chlamydia trachomatis*: a comparison of two partner testing methods, **358**
- Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
- Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- Recent pilot studies of chlamydia screening, **352**
- Repeat chlamydia screening by mail, San Francisco, **28**
- Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- Spending money to save money, **4**
- The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- Validity and utility of screening tests for STIs, **356**
- Who is being tested for genital chlamydia in primary care?, **234**
- self collected urine samples**, Improving response rates for self collected urine samples, **346**
- self obtained samples**, Self obtained samples for genitourinary infections, **171**
- sensitivity**, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- seroepidemiological study**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- serology**, Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- seroprevalence**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area, **497**
- serotesting**, Genital herpes serotesting: a study of the epidemiology and patients' knowledge and attitude among STD clinic attenders in Coventry, UK, **35**
- sex**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
Sexual behaviour of lesbians and bisexual women, **147**
Who is being tested for genital chlamydia in primary care?, **234**
- sex behaviour**, Sexual behaviour of lesbians and bisexual women, **147**
- sex differences**, Sex differences in the experience of testing positive for genital chlamydia infection: a qualitative study with implications for public health and for a national screening programme, **372**
- sex on premises venues**, A novel research approach in sex on premises venues (SOPV): objective measure of sexual behaviour and low level intrusion to patrons, **78**
- sex resort**, Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- sex workers**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
Social and behavioural factors associated with condom use among direct sex workers in Siem Reap, Cambodia, **163**
- sexual abuse**, Psychosocial predictors of HIV/STI risk behaviours in a sample of homosexual men, **142**
- sexual assault**, Follow up of patients who have been recently sexually assaulted, **349**

- sexual behaviour**, A novel research approach in sex on premises venues (SOPV): objective measure of sexual behaviour and low level intrusion to patrons, **78**
- Associations between internet sex seeking and STI associated risk behaviours among men who have sex with men, **466**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- Reducing the risk of gonorrhoea in black Caribbean men: can we identify risk factors?, **119**
- sexual health**, Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- Sexual health—health of the nation, **85**
- sexual health services**, Staff training in integrated sexual health services, **354**
- sexual risk**, HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy, **59**
- sexual risk factors**, A population based study of herpes simplex virus 2 seroprevalence in rural Costa Rica, **460**
- sexual transmission**, A national survey of genitourinary medicine clinic attenders provides little evidence of sexual transmission of hepatitis C virus infection, **301**
- Absence of hepatitis C virus transmission in a prospective cohort of heterosexual serodiscordant couples, **160**
- sexually transmitted disease**, Chancroid: clinical manifestations, diagnosis, and management, **68**
- sexually transmitted diseases**, Avoiding risky sex partners: perception of partners' risks v partners' self reported risks, **197**
- Determinants of disclosure of genital herpes to partners, **42**
- Geographical focusing: an intervention to address increased risk for sexually transmitted diseases during repatriation and resettlement in post-war Mozambique, **77**
- Tetracycline treatment does not eradicate *Mycoplasma genitalium*, **318**
- Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
- Variation in clinical practice in genitourinary medicine clinics in the United Kingdom, **240**
- Which factors affect access to STD care? A comparison of a hospital based clinic and an outreach service, **347**
- sexually transmitted infection**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- Sexually Transmitted Infections**, Taking up the reins, **2**
- sexually transmitted infections**, High postpartum rates of sexually transmitted infections among teens: pregnancy as a window of opportunity for prevention, **469**
- Is use of antiretroviral therapy among homosexual men associated with increased risk of transmission of HIV infection?, **7**
- National survey of doctors' actions following the diagnosis of a bacterial STD, **254**
- Predictors of the prevalence of bacterial STI among young disadvantaged Indigenous people in north Queensland, Australia, **332**
- Rural sex work in Cambodia: work characteristics, risk behaviours, HIV, and syphilis, **e2**
- Sexual health—health of the nation, **85**
- Sexually transmitted infections in Africa: single dose treatment is now affordable, **432**
- STI services in the United Kingdom: a way forward, **439**
- STI services in the United Kingdom: how shall we cope?, **435**
- Surveillance of sexually transmitted infections in primary care, **174**
- Treating sexually transmitted infections in primary care: a missed opportunity?, **134**
- Use of a leaflet to replace verbal pretest discussion for HIV: effects and acceptability, **243**
- Validity and utility of screening tests for STIs, **356**
- Sexually Transmitted Infections Foundation**, Impact of the Sexually Transmitted Infections Foundation course on the knowledge of family planning nurses and doctors, **346**
- shared care**, Shared care in the management of genital *Chlamydia trachomatis* infection in primary care, **369**
- South Africa**, HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa, **442**
- Impact of on-site testing for maternal syphilis on treatment delays, treatment rates, and perinatal mortality in rural South Africa: a randomised controlled trial, **208**
- STI case management at a South African teaching hospital, **498**
- specificity**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- Sri Lanka**, Seroepidemiological study of herpes simplex virus types 1 and 2 in Brazil, Estonia, India, Morocco, and Sri Lanka, **286**
- staff training**, Staff training in integrated sexual health services, **354**
- STD**, STD trends and patterns of treatment for STD by physicians in private practice in Peru, **403**
- STD clinic**, Prevalence of and risk factors for hepatitis C virus infection among STD clinic clientele in Miami, Florida, **e1**
- STD clinics**, Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India, **425**
- STD intervention trials**, Comparison of STD prevalences in the Mwanza, Rakai, and Masaka trial populations: the role of selection bias and diagnostic errors, **98**
- STD prevention**, Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
- STI policies**, A survey of STI policies and programmes in Europe, **171**
- STI prevalence**, Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- STIs**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- Chaperoning in genitourinary medicine clinics, **422**
- Sexually transmitted infections in male clients of female sex workers in Benin: risk factors and reassessment of the leucocyte esterase dipstick for screening of urethral infections, **388**
- streptococcal infections**, Group B streptococcus carriage and vulvovaginal symptoms: causal or casual? A case-control study in a GUM clinic population, **246**
- superior mesenteric artery syndrome**, Superior mesenteric artery syndrome in an HIV positive patient, **262**
- survey**, A survey of STI policies and programmes in Europe, **171**
- survey method**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- susceptibility**, Antimicrobial susceptibility of *Neisseria gonorrhoeae* strains isolated in Guangzhou, China, 1996–2001, **399**
- syndromic diagnosis**, Prevalence of sexually transmitted infections and performance of STI syndromes against aetiological diagnosis, in female sex workers of red light area in Surat, India, **111**
- syphilis**, An outbreak no longer: factors contributing to the return of syphilis in Greater Manchester, **291**
- High frequency of antibodies to syphilis and HIV in hepatitis C virus positive blood donors may reflect its sexual transmission in this region, **170**
- Is increased surveillance for asymptomatic syphilis in an HIV outpatient department worthwhile?, **257**
- Perforating chancre: any cause-effect relation with HIV infection?, **262**
- Social capital, poverty, and income inequality as predictors of gonorrhoea, syphilis, chlamydia and AIDS case rates in the United States, **62**
- Syphilis prevalence has rapidly decreased in South Korea, **323**
- Syphilis specific antibodies in newborn infants in Lower Saxony, Germany 1993–2001, **351**
- Use of PCR in the diagnosis of early syphilis in the United Kingdom, **479**
- Tanzania**, Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya Region, Tanzania, **382**
- Is antenatal syphilis screening still cost effective in sub-Saharan Africa, **375**
- Variations of HIV and STI prevalences within communities neighbouring new goldmines in Tanzania: importance for intervention design, **307**
- telephone survey**, Effects of two telephone survey methods on the level of reported risk behaviours, **325**
- television**, Television coverage of AIDS and transmission of the AIDS virus, **170**
- termination of pregnancy**, Chlamydia testing before termination of pregnancy, **352**
- test of cure**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**

- tinidazole**, Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
- treatment interruption**, Enhanced risk of HIV sexual transmission during structured treatment interruption, **74**
- Trichomonas vaginalis***, Comparison of culture and different PCR assays for detection of *Trichomonas vaginalis* in self collected vaginal swab specimens, **393**
Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic, **151**
Was the Papanicolaou smear responsible for the decline of *Trichomonas vaginalis*?, **263**
- trichomoniasis**, Diagnosis and treatment of presumed STIs at Mexican pharmacies: survey results from a random sample of Mexico City pharmacy attendants, **224**
- tuberculosis**, Adult respiratory distress syndrome as a severe immune reconstitution disease following the commencement of highly active antiretroviral therapy, **337**
Coexistent cranial tuberculomas and tuberculosis of the cervix in a postmenopausal woman, **496**
- tuberculous meningitis**, Cytokine profiles in HIV seropositive patients with tuberculous meningitis, **75**
- type specific serology**, The acceptability of the introduction of a type specific herpes antibody screening test into a genitourinary medicine clinic in the United Kingdom, **129**
- Ukraine**, The HIV/AIDS epidemic in Ukraine: stable or still exploding?, **263**
- United Kingdom**, Sexual health—health of the nation, **85**
STI services in the United Kingdom: a way forward, **439**
STI services in the United Kingdom: how shall we cope?, **435**
- urine**, Correlation between culture testing of swabs and ligase chain reaction of first void urine from patients recently treated for *Chlamydia trachomatis*, **237**
- urine testing**, Chlamydia screening in the United Kingdom, **3**
Opportunistic screening for genital chlamydial infection. I: Acceptability of urine testing in primary and secondary healthcare settings, **16**
Opportunistic screening for genital chlamydial infection. II: Prevalence among healthcare attenders, outcome, and evaluation of positive cases, **22**
- vaccination**, Hepatitis B vaccination for sex workers: do outreach programmes perform better?, **157**
- vaginitis**, Late recurrence of resistant *Trichomonas vaginalis* vaginitis: relapse or re-infection?, **260**
- varicella zoster virus**, Detection of varicella zoster virus in genital specimens using a multiplex polymerase chain reaction, **298**
- VDRL test**, Syphilis prevalence has rapidly decreased in South Korea, **323**
- virus-like particles**, Prospective study on cervical neoplasia: presence of HPV DNA in cytological smears precedes the development of cervical neoplastic lesions, **191**
- vulvovaginal yeasts**, Vulvovaginal carriage of yeasts other than *Candida albicans*, **249**
- women**, Adolescence and other risk factors for *Chlamydia trachomatis* genitourinary infection in women in Melbourne, Australia, **31**
Adolescent women underestimate their susceptibility to sexually transmitted infections, **408**
Sensitivity requirements for the point of care diagnosis of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in women, **363**
Seroprevalence of reproductive tract infections in women in northern India—a relatively low prevalence area