IN PRACTICE

Methods employed by genitourinary medicine clinics in the United Kingdom to diagnose bacterial vaginosis

F E A Keane, R Maw, C Pritchard, C A Ison

Sex Transm Infect 2005;81:155–157. doi: 10.1136/sti.2004.009902

See end of article for
authors' affiliations

Correspondence to:
Dr Frances Keane,
Department of
Genitourinary Medicine,
Royal Cornwall Hospital,
Treliske, Truro TR1 3LJ,
UK; keanefr@rcht.
cornwall.nhs.uk

Accepted for publication
3 June 2004

Objective: To determine the methods used by genitourinary medicine (GUM) clinics in the United Kingdom for the diagnosis of bacterial vaginosis (BV).

Methods: A questionnaire survey of UK GUM clinics was conducted.

Results: 148/221 (67%) clinics returned a questionnaire. 96/148 (64.9%) clinics reported using Amsel's criteria to diagnose BV but only 29 (30.5%) of these used all four of the composite criteria. 139/148 (93.9%) clinics used the appearance of a Gram stained vaginal smear as an aid in BV diagnosis, although a variety of scoring methods was employed. In the majority of clinics, 92/148 (62.2%), one staff discipline provided the microscopy service, in 50 (33.8%) clinics two staff disciplines provided microscopy services. The bulk of microscopy services within UK GUM clinics is provided by nurses.

Conclusions: Most UK GUM clinics utilise the appearance of a Gram stained vaginal smear for the diagnosis of BV although there is little consensus at present about the type of scoring method employed. Adaptation of a uniform scoring method would have enormous benefits, including consistency and reproducibility of results and the development of quality assurance schemes for BV diagnosis on a national basis. There are important issues to be addressed regarding the initial training and ongoing support for nurses providing microscopy services within UK GUM clinics.

Bacterial vaginosis (BV) is characterised by an alteration in the vaginal ecology where the normal *Lactobacillus* dominated flora is replaced by a mixed bacterial flora that includes *Gardnerella vaginalis*, *Mycoplasma hominis*, and anaerobes such as *Mobiluncus* spp and *Prevotella* spp. Although the condition was first described (at that time as *Haemophilus vaginalis* vaginitis) in 1955 by Gardener and Dukes¹ it was not until 1983 that Amsel *et al*² proposed standardised diagnostic criteria. This method, which has remained the gold standard for the diagnosis of BV until recent times, proposed that BV was diagnosed on the presence of three out of four composite criteria—that is, a characteristic thin, homogeneous vaginal discharge, vaginal pH above 4.5, a positive “whiff” test, and the presence of clue cells on dark ground microscopy of a saline wet mount preparation. In the same year, Spiegel *et al*,³ of whom Amsel was one, published their work on the use of the appearance of a Gram stained vaginal smear for the diagnosis of BV. They assigned a simple scoring system based on the presence of *Lactobacillus* spp, *Gardnerella* spp, and other morphotypes. A direct comparison of the two methods by Eschenbach *et al*⁴ found varying sensitivities (43% to 97%) and specificities (53% to 99%) of each of the four criteria when compared to Spiegel's Gram stain criteria. Since then, several other scoring systems, including Nugent's,⁵ a modified Spiegel's (Hay-Ison),⁶ and a modified Ison-Hay⁷ method have been proposed and validated.

Genitourinary medicine (GUM) clinics in the United Kingdom routinely perform in-house diagnosis of BV. Until now, their usage of the available diagnostic tools has not been studied. A recent pilot external quality assurance scheme for the diagnosis of BV on the appearance of Gram stained vaginal smears revealed little consistency of scoring system preference between the five participating centres (F Keane, personal communication). The following questionnaire survey was designed to investigate current diagnostic practices for BV within UK GUM clinics.

METHODS

A questionnaire (designed by FK) was distributed by the British Co-operative Clinical Group to all lead clinicians in UK GUM clinics. They were surveyed about the use of Amsel's criteria, examination of a Gram or acridine orange stained vaginal smear, or other method to diagnose BV in their routine practice. Clinics using the appearance of Gram stained vaginal smears were asked to disclose their scoring system. Information was also gathered on the staff disciplines providing routine microscopy services within each clinic and the use of microbiology laboratories to provide a diagnostic service for BV to GUM clinics. Data were analysed using the SPSS statistical package.

RESULTS

Replies were received from clinicians responsible for 148/221 (67%) clinics.

Diagnostic methods for BV

A total of 96/148 (64.9%) clinics reported using Amsel's composite criteria to diagnose BV. One of these did not detail how many of the criteria it routinely employed. Of the remaining 95 clinics, 10 (10.5%) reported using only one of the criteria, 27 (28.4%) two, 29 (30.5%) three, and only 29 (30.5%) reported using all four of the criteria necessary to make a diagnosis of BV by this method. The use of the individual criteria is detailed in table 1. Of the 25 (16.9%) replying no to the use of Amsel's criteria and the 27 (18.2%) who left the reply blank, a substantial proportion reported the use of at least some of the composite criteria (table 1); 139/148 (93.9%) clinics reported using examination of a Gram stained vaginal smear for the diagnosis of BV; 44 (31.7%) of these did not report any scoring method. Of the

Abbreviations: BV, bacterial vaginosis; GUM, genitourinary medicine; MLSOs, medical laboratory scientific officers

Table 1 Use of Amsel's criteria

Component of Amsel's criteria	Yes (n = 96) (64.9%)	No (n = 25) (16.9%)	Blank reply (n = 27) (18.2%)
Discharge	91 (95.8)	11 (44.0)	26 (96.3)
pH	58 (61.1)	2 (8.0)	13 (48.1)
"Whiff test"	37 (38.9)	4 (16.0)	5 (18.5)
Clue cells on "wet" preparation	81 (85.3)	9 (36.0)	24 (88.9)

remaining 95 clinics, 94 employed one scoring system alone, but one clinic reported five different methods. Of the 99 scoring methods employed—24 were Nugent's scoring system, eight Spiegel's, 33 Hay/Ison method, and 34 "some other" scoring method.

Only five (3.3%) clinics used the appearance of an acridine stained vaginal smear for the diagnosis of BV. Ten clinics reported "other" diagnostic methods," consisting of various combinations of the above and the use of anaerobic cultures in two cases. Samples were sent to microbiology laboratories by 36/148 (24%) clinics to aid in the diagnosis of BV. Of these, 14 sent air dried and nine Gram stained vaginal smears respectively; 31 sent charcoal swabs for culture of *Gardnerella vaginalis*.

Staff disciplines performing microscopy

The microscopy service was provided by one staff discipline in 92/148 (62.2%) clinics; in 49 (53.3%) of these clinics by nurses, 27 (29.3%) by medical laboratory scientific officers (MLSOs), 15 (16.3%) by doctors, and in one by some "other" unspecified staff member. In 50 (33.8%) clinics, microscopy services were provided by two different staff disciplines: in 34 (68%) of these clinics by doctors and nurses, in six (12%) by doctors and MLSOs, in six (12%) by nurses and MLSOs, and in four (by doctors and "other"). In five clinics there was a

multidisciplinary provision of microscopy services and one clinic had no "in-house" laboratory service.

DISCUSSION

This is the first survey to examine routine "in-house" diagnostic methods for BV within UK GUM clinics. The data regarding the use of Amsel's criteria are difficult to interpret: less than a third of those using this method actually employed all of the composite criteria to do so, thus, by definition, invalidating the method. This would suggest that the use of Amsel's criteria is not the method of choice for diagnosing BV in routine clinical practice. Interestingly, the overwhelming majority of survey respondents were using the appearance of a Gram stained vaginal smear to aid their diagnosis of BV. However, although there was an unwritten consensus that the Gram stain was the most useful method for the diagnosis of BV, there was poor agreement about the most suitable scoring system. An important caveat to mention is that although the appearance of the vaginal Gram stain should be used to interpret a patient's vaginal flora; the result needs to be viewed in a clinical context to establish whether a patient requires treatment. It is well accepted that *G vaginalis* can be isolated from women both with and without BV^{4,8} and thus sending cultures for *G vaginalis* is unhelpful in trying to establish a diagnosis of BV.

The adoption of a single scoring system for the diagnosis of BV on Gram stained vaginal smears by all GUM clinics in the United Kingdom would be extremely advantageous. Firstly, from the point of consistency; there would be increased confidence that a Gram stain vaginal smear taken in any UK clinic would be interpreted in a similar manner. Secondly, adoption of the one system would aid enormously in the development/implementation of both internal and external quality assurance schemes for the diagnosis of BV in UK GUM clinics. Furthermore, use of a single scoring system would strengthen the body of BV research in the United Kingdom as studies would, by definition, become more directly comparable than they are at present. Most BV researchers now favour the Gram stain method over Amsel's criteria; slides may be stored and both Gram stained and read in batches. The method offers convenience and an opportunity for confirmation of findings that is lost when Amsel's method is employed.

Ison and Hay have validated their modified simple grading system⁷ against both Amsel's criteria² and Nugent's method³ in an international comparison⁹ and have found excellent correlations. The Ison-Hay method can be easily incorporated into routine clinic practice, unlike Nugent's scoring system, which requires the counting of different types of bacteria, making it too time consuming for use in a busy clinical setting. The Ison-Hay method is recommended by the Bacterial Special Interest Group (BSIG) as the method of choice for the diagnosis of BV in GUM clinics. It has also been recommended for screening for BV in the guidelines for screening for sexually transmitted infections (Clinical Effectiveness Group/BSIG of BASHH, in preparation). Uniform adoption of the Ison-Hay method across UK GUM

Key messages

- This survey, which examines current diagnostic practice for BV in UK GUM clinics, reveals the following:
 - the use of Amsel's criteria is largely unsatisfactory as most clinics do not employ all of the criteria—an essential prerequisite of this method
 - most UK clinics use the appearance of a Gram stained vaginal smear to aid in the diagnosis of BV, but there is little consensus about the scoring methods they employ
 - the bulk of microscopy services within UK GUM clinics are provided by nurses
- The authors recommend the universal adoption of the Ison-Hay scoring system to
 - improve diagnostic consistency across clinics
 - assist in the development /implementation of both internal and external quality assurance schemes for the diagnosis of BV
 - strengthen the body of BV research in the UK as studies
- There are important issues to be addressed regarding the initial training and on-going support for nurses providing microscopy services within UK GUM clinics.

clinics could achieve the desired consistency of BV diagnosis in areas of both clinical and research practice and assist in the development of robust quality assurance schemes for BV diagnosis in GUM clinic laboratories.

The survey also revealed that GUM nurses provide the bulk of microscopy services with UK GUM clinics. This raises important issues regarding initial training, on-going support and clinical governance for the delivery of microscopy services within GUM services.

ACKNOWLEDGEMENTS

The authors would like to thank the Bacterial Special Interest Group of the British Association for Sexual Health and HIV and the British Co-operative Clinical Group for their support of this project and are indebted to all the clinics that completed the survey questionnaires.

CONTRIBUTORS

FK designed the questionnaire, performed the initial statistical analysis, and wrote the manuscript; RM contributed to the questionnaire design, distributed the questionnaire, collated the responses, and commented on the manuscript; CP carried out more detailed statistical analysis; CI contributed to the questionnaire design and the discussion section of the manuscript.

Authors' affiliations

F E A Keane, Department of Genito-urinary Medicine, Royal Cornwall Hospital, Truro, UK

R Maw, Department of Genito-urinary Medicine, Royal Victoria Hospital, Belfast, UK

C Pritchard, Research and Development Support Unit, Royal Cornwall Hospital, Truro, UK

C A Ison, Department of Infectious Diseases and Microbiology, Imperial College London, St Mary's Campus, London, UK

Conflict of interest: None.

REFERENCES

- 1 **Gardner HL**, Dukes CD. Haemophilus vaginalis vaginitis. A newly defined specific infection previously classified "non-specific" vaginitis. *Am J Obstet Gynecol* 1955;**69**:962-76.
- 2 **Amsel R**, Totten PA, Spiegel CA, *et al*. Nonspecific vaginitis: diagnostic criteria and microbial and epidemiological associations. *Am J Med* 1983;**74**:14-22.
- 3 **Spiegel CA**, Amsel R, Holmes KK. Diagnosis of bacterial vaginosis by direct Gram stain of vaginal fluid. *J Clin Microbiol* 1983;**18**:170-7.
- 4 **Eschenbach DA**, Hillier S, Critchlow C, *et al*. Diagnosis and clinical manifestations of bacterial vaginosis. *Am J Obstet Gynecol* 1988;**158**:819-28.
- 5 **Nugent RP**, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardised method of Gram-stain interpretation. *J Clin Microbiol* 1991;**29**:297-301.
- 6 **Hay PE**, Lamont RF, Taylor-Robinson D, *et al*. Abnormal bacterial colonisation of the genital tract and subsequent preterm delivery and late miscarriage. *BMJ* 1994;**308**:295-8.
- 7 **Ison CA**, Hay PE. Validation of a simplified grading of Gram-stained vaginal smears for use in genitourinary medicine clinics. *Sex Transm Infect* 2002;**78**:413-5.
- 8 **Totten PA**, Amsel R, Hale J, *et al*. Selective differential human blood bilayer media for isolation of Gardnerella (Haemophilus) vaginalis. *J Clin Microbiol* 1982;**15**:141-7.
- 9 **Forsum U**, Jakobsson T, Larsson PG, *et al*. An international study of the interobserver variation between interpretations of vaginal smear criteria of bacterial vaginosis. *APMIS* 2002;**110**:811-8.