

# **P1-S2.61 HIGH HIV RATES AMONG MEN WHO HAVE SEX WITH MEN IN JAMAICA DESPITE INCREASED PREVENTION EFFORTS**

doi:10.1136/sextrans-2011-050108.118

<sup>1</sup>J P Figueroa, <sup>2</sup>S Weir, <sup>3</sup>C J Cooper, <sup>4</sup>L Byfield, <sup>4</sup>S Eastman, <sup>2</sup>M Hobbs, <sup>4</sup>J Duncan.  
<sup>1</sup>University of the West Indies, Kingston, Jamaica; <sup>2</sup>University of North Carolina, Chapel Hill, USA; <sup>3</sup>Epidemiology and Research Training Unit, Kingston, Jamaica; <sup>4</sup>Ministry of Health, Jamaica

**Background** Adult HIV prevalence in Jamaica is 1.7% while rates among men who have sex with men (MSM) were 32% in a 2007 survey. Following this survey prevention interventions among MSM were increased. The objective of this study was to estimate HIV rates among MSM in 2011 and compare findings with the 2007 survey.

**Methods** An experienced research nurse attended prevention workshops for MSM as well as their social activities and invited MSM to participate in a cross sectional survey. MSM were interviewed and tested for HIV and other STIs. Sensitive behavioural data was interviewer administered and self-completed. Confidential results were given to participants with treatment as indicated. Ethical approval and informed consent was obtained. The 2011 survey is ongoing so results are preliminary.

**Results** 33% of 157 MSM were HIV positive in 2011 compared with 32% of 201 MSM in 2007. In 2011, 8 of 157 (5.1%) MSM were positive for syphilis and 3.8% were indeterminate. 12 of 139 MSM (8.9%) were positive for Chlamydia and 2 (1.4%) had gonorrhoea. In 2007 HIV positive MSM were more likely to ever have a STI (37.5% vs 19.0%,  $p=0.004$ ) and to be receptive (73.4% vs 59.1%,  $p=0.005$ ). 33% of MSM reported sex with a woman in the past 4 weeks and 65% reported ever having sex with a woman. Nearly 60% of HIV positive MSM had not disclosed their status to their partner. MSM who were of low socio-economic status, ever homeless and victims of physical violence were significantly more likely to be HIV positive.

**Conclusions** HIV prevalence among MSM remains unacceptably high. MSM are more socially vulnerable than the general population. Many MSM reported having sex with a woman and not disclosing their HIV status to their partners. MSM are likely to act as a bridge for HIV into the general population. Prevention efforts among MSM need to be scaled up urgently and measures taken to reduce their social vulnerability including stigma and discrimination.

Abstract P1-S2.61 Figure 1

# **P1-S2.62 FACTORS ASSOCIATED WITH HIGH-RISK PENETRATIVE SEX IN A COHORT OF TREATMENT EXPERIENCED HIV-POSITIVE MEN WHO HAVE SEX WITH MEN (MSM) IN BRITISH COLUMBIA**

doi:10.1136/sextrans-2011-050108.119

<sup>1</sup>W Michelow, <sup>2</sup>G Talany, <sup>1</sup>N O'Brien, <sup>1</sup>H Wang, <sup>1</sup>J Forrest, <sup>1</sup>A Palmer, <sup>1</sup>J S G Montaner, <sup>2</sup>R Hogg. <sup>1</sup>BC Centre for Excellence in HIV/AIDS, Canada; <sup>2</sup>Simon Fraser University, Vancouver, Canada

**Background** In Canada, men who have sex with men (MSM) are disproportionately infected with HIV. Despite much evidence demonstrating the secondary preventive value of HAART, we have not fully capitalised on potential synergies between treatment and

prevention. This study investigates high-risk sexual behaviour among treatment-experienced MSM in British Columbia, Canada.

**Methods** We analysed data from the Longitudinal Investigations into Supportive and Ancillary Health Services (LISA) project on MSM (gay, bisexual, or reported sex with men in the 6 months prior to interview). High-risk sexual behaviour was penetrative anal or vaginal sex in the 6 months prior to interview with less than 100% condom use. Multivariate logistic regression was used to identify factors associated with high-risk penetrative sex.

**Results** Of 346 HIV-positive MSM, median age was 46 years, 268 (78%) were currently on HAART, with 185 (54%) having  $\geq 350$  CD4 cells/ $\mu$ L. Ninety-eight (28%) participants endorsed seeking out HIV-positive partners for sex, and 147 (43%) reported being less likely to use a condom with an HIV-positive partner. In the regression model, HIV-positive MSM with a CD4 count  $\geq 350$  cells/ $\mu$ L were more likely to report unprotected penetrative sex (Adjusted OR [AOR] =2.7; 95% CI 1.5 to 5.2), more likely to not use condoms with their HIV-positive partners (AOR=12.0; 95% CI:6.4 to 22.3), and less likely to report sex with anonymous partners in the past 6 months (AOR=0.28; 95% CI 0.1 to 0.8). No significant associations were found between high-risk penetrative sex and current HAART use, viral suppression, or treatment adherence.

**Conclusion** HIV-positive MSM who report engaging in high-risk penetrative sex are more likely to be healthier (higher CD4 counts), to have unprotected sex with other HIV-positive partners (serosorting) and with known rather than anonymous sex partners. Despite healthier MSM reporting more high-risk sex, results suggest this risk-taking is discriminate and would not necessarily lead to more onward HIV transmission.

# **P1-S2.63 ASSOCIATION BETWEEN AGE AND STI AMONG MEN WHO HAVE SEX WITH MEN (MSM) IN JAMAICA**

doi:10.1136/sextrans-2011-050108.120

<sup>1</sup>S Weir, <sup>2</sup>P Figueroa, <sup>3</sup>C Jones-Cooper, <sup>4</sup>T Hylton-Kong, <sup>1</sup>M Hobbs, <sup>1</sup>K Rich, <sup>3</sup>L Byfield. <sup>1</sup>University of North Carolina, Chapel Hill, USA; <sup>2</sup>University of the West Indies, Kingston, Jamaica; <sup>3</sup>Ministry of Health, Kingston, Jamaica; <sup>4</sup>Epidemiology & Research Training Unit, Kingston, Jamaica

**Background** In 2007, 201 men who have sex with men were recruited at social venues and through peer referral in five urban areas of Jamaica for an exploratory study of the prevalence of sexually transmitted infection in the population and identification of risk factors.

**Objective** To describe the association between age and current infection with HIV, syphilis, and chlamydia and explore reasons for the differences.

**Methods** MSM were invited to participate in a survey and testing. Urine and blood samples were collected at the time of the interview and later tested for evidence of HIV, syphilis, gonorrhoea and chlamydia. Men were given options about where to obtain results.

**Results** Of the three infections, HIV had the strongest positive association with age; syphilis a weaker positive association; and chlamydia a weak negative association. Risk factors varied somewhat by age and infection see Abstract P1-S2.63 Figure 1. For example, not using a condom in the past 4 weeks was associated with chlamydia infection among MSM age 15–19 but not for older MSM. Having a main sexual partner was associated with HIV infection for MSM age 15–19 but not for older MSM.

**Conclusion** Strategies for prevention of transmission need to be aware of differences in risk by age and by infection.