

from 46 public STD clinics in the 4 provinces. Specimens that tested positive for *C. trachomatis* by polymerase chain reaction were genotyped for the infecting *C. trachomatis* strain by amplifying and sequencing the genovar-specific *ompA* gene, which encodes the chlamydial major outer membrane protein. Nine *C. trachomatis* genovars were identified from 129 specimens; they consisted of the F (25.6%, n = 33), E (17.1%, n = 22), J (16.3%, n = 21), D (15.5%, n = 20), G (11.6%, n = 15), K (5.4%, n = 7), H (3.9%, n = 5), I (1.6%, n = 2), and B (0.8%, n = 1) genovars. Nine genovars were found in specimens from Guangxi province, 6 were found in specimens from each of Guangdong and Hainan provinces, but only 5 were found in specimens from Jiangsu province. Significant differences were observed in *C. trachomatis* genovar distributions between different provinces: G/Ga was absent among male STD clinic patients (MSPs) from the eastern province of Jiangsu (Fisher exact test = 0.036), while being prevalent in the 3 southern provinces.

P3.261 GLOBAL MULTILOCUS SEQUENCE TYPE (MLST) ANALYSIS OF CHLAMYDIA TRACHOMATIS STRAINS FROM 16 COUNTRIES

doi:10.1136/sextrans-2013-051184.0717

¹J Isaksson, ²R Bom, ¹I Saleh, ¹B Herrmann, ²S Bruisten. ¹Department of Clinical Microbiology, Uppsala, Sweden; ²Public Health Laboratory, Public Health Service of Amsterdam, Amsterdam, The Netherlands

Background The *Chlamydia trachomatis* MLST database was established in 2007 and is based on five target regions (non-housekeeping genes) and the conventional *ompA* gene. It enables worldwide epidemiological analyses.

Methods Samples were included from 11 studies focusing on specific risk groups and with different study objectives. Geographical distribution of MLST profiles was carried out and eBURST analysis identified genetic founders.

Results A total of 414 MLST profiles were recognised from 2087 entries. Polymorphism of target regions was reflected in varying numbers of alleles; *hctB* 89, CT058 51, CT144 30, CT172 41, *pbpB* 35. With addition of 49 *ompA* gene variants 459 profiles exist.

There were 16 countries from which samples originated, the majority coming from the Netherlands 39%, Sweden 16% and Norway 12%, but also samples from countries in North and South America, Africa and other European countries. Overall 605 (29%) of the database samples originated from men having sex with men (MSM) while the remaining samples were assumed to be from heterosexuals.

Singletons or small clusters emerged from 377 of the MLST profiles that were found 1–9 times while 31 profiles were found 10–43 times each. There were 8 profiles that predominated and were each found between 83 and 140 times and constituted large clusters that comprised 868 samples (41.6%). Four of the predominating profiles were strongly associated with MSM, with 96–100% of the samples coming from MSM. In the other four large clusters heterosexuals comprised > 90% of the samples. eBURST analysis identified 3 of the 8 predominating profiles as founders, and another 3 as subgroup founders. The genetic diversity was much lower in the MSM clusters compared to clusters in heterosexuals.

Conclusions Worldwide a few *C. trachomatis* MLST profiles predominate. Different MLST profiles predominate among MSM and heterosexuals.

P3.262 COMPARISON OF URINE SAMPLES AND PENILE SWABS IN THE DETECTION OF HUMAN PAPILLOMA VIRUS IN MEN USING THE SPF₁₀ LINE PROBE ASSAY

doi:10.1136/sextrans-2013-051184.0718

¹F M P A Koene, ¹P F G Wolffs, ¹A A T P Brink, ²N H T M Dukers-Muijers, ³W G V Quint, ¹C A Bruggeman, ^{1,2}C J P A Hoebe. ¹Department of Medical Microbiology, School of Public Health and Primary Care (CAPHRI), Maastricht University Medical Center, Maastricht, The Netherlands; ²Department of Sexual Health, Infectious Diseases and Environment, South Limburg Public Health Service, Geleen, The Netherlands; ³DDL Diagnostic Laboratory, Rijswijk, The Netherlands

Background Penile swab sampling is the gold standard when testing for human papilloma virus (HPV) in men. Urine could provide a more convenient sampling material. Therefore we compared the detection and typing of HPV in urine samples and penile swabs using a highly sensitive method.

Methods First void urine and self-obtained penile swabs were collected from 120 men visiting a Sexually Transmitted Infections (STI) clinic in South Limburg, The Netherlands. When exclusion criteria were met, the results of 111 men (mean age 29.4 years) were analysed. Broad-spectrum HPV DNA amplification and genotyping were performed using SPF₁₀-DEIA-LiPA₂₅ system (Labo Bio-Medical Products, Rijswijk, The Netherlands). There are 12 high risk HPV types (hrHPV), 5 possible hrHPV types and 8 low risk HPV types (lrHPV) detectable by SPF₁₀-LiPA₂₅.

Results HPV DNA was found in 75 (67.6%) men. In 73 men HPV DNA was detected in the penile swab or both samples and in two men HPV DNA was detected only in the urine sample. Sixty-six paired samples were concordant in being HPV positive (n = 30) or negative (n = 36). Eleven of the 30 matching samples were concordant in their genotypes, fourteen pairs were comparable (≥ 1 genotype identical) and five pairs were discordant in genotypes. Three of these discordant and one comparable urine sample contained (possible) high risk types which were not found in the swab. Furthermore, the two positive urines with a negative swab contained (possible) high risk types as well.

Conclusion Urine samples are not comparable to penile swabs in the detection of HPV in men. However, in the urine sample of six men high risk types were found that could not be detected in the penile swab alone. This could be an indication of an additional value of the urine sample aside the penile swab in the detection of HPV in men.

P3.263 DO MEN HARBOUR THE SAME STRAIN OF GONOCOCCUS AT MULTIPLE ANATOMICAL SITES?

doi:10.1136/sextrans-2013-051184.0719

K Eastick. Scottish Bacterial STI Reference Laboratory, Royal Infirmary of Edinburgh, Edinburgh, UK

Background All *N. gonorrhoeae* clinical isolates in Scotland are routinely typed by *Neisseria gonorrhoeae* multi-antigen sequence typing (NG-MAST). The frequency with which individuals harboured the same *N. gonorrhoeae* strain at > 1 anatomical site concurrently was examined.

Methods Episodes of gonorrhoea between 2004 and January 2013 comprising typed isolates from two or more anatomical sites of the same male individual submitted within a one-month period were analysed for similarity.

Results 410 episodes of gonorrhoea were identified with > 1 site cultured. The table shows differences in sequence type (ST) within episodes. Forty episodes with three cultures are included in all three pairwise combinations. Overall 91.2% of episodes had identical STs at all sites.

Where STs differed at only one allele, the sequences were compared using CLUSTALW. Thirteen of 14 alleles compared were at the *porB* locus. Nine pairs were 99.8% similar, representing a difference of 1 nucleotide. The remaining pairs showed 88.4%, 98.2%, 97.8%, 96.9% and 75.1% (*tbpB*) similarity.