Highlights from this issue

doi:10.1136/sextrans-2013-051498

Jackie A Cassell, *Editor*

Welcome to our first issue of 2014. UK readers will have followed with interest the recent publication of the 3rd National Survey of Sexual Attitudes and Lifestyles (Natsal—see natsal.ac.uk), which coincided with World AIDS day 2013. Population surveys however have their limitations—particularly in their ability to describe in useful detail minority sexual practices. These need to be explored through customised surveys using targeted recruitment, or through data collection systems of sufficiently large scale to capture useful numbers of these groups.

Two populations in which Natsal has limited power are men who have sex with men (MSM) and sex workers of both genders. Improving data quality in GUMCAD, the electronic records based surveillance system for genitourinary medicine in England, is enabling improving power and accuracy in the study of vulnerable minority populations. McGrath-Lone and colleagues¹ report on the sexual health of male sex workers in England—0.08% of all men attending clinics were coded as sex workers, and this group experienced high rates of sexually transmitted infections (STI). Data on behavioural risk factors are currently limited, but likely to improve with planned changes in surveillance. The STI and HIV transmission potential of sex work relates to levels of condom use, which varies enormously across the world. While rates of unprotected sex tend to be higher in developing countries, Eccles *et al*² report advertisements for unprotected sex by female indoor sex workers in the north of England. This was associated with lower hourly rates and the offer of anal intercourse. These findings have important implications for health promotion to this group.

Sex parties are also hard to study using population-based surveys, so Grov *et al*³ used banner advertising on a website to recruit MSM for an online survey of sex party participation. MSM attending such parties reported high levels of many of risk characteristics, suggesting the need for the targeting of HIV and STI promotion activities for this group. Clinics and surveillance systems tend to assign infections to the highest risk activity or contact group, and in so doing can overlook the potential contribution of other practices to STI and HIV transmission. Recognising this challenge, Nash *et al*⁴ undertook a cross-sectional

survey of electronic health records of MSM in Melbourne exploring the contribution of practices other than anal sex to transmission of STI. They conclude that while urethral chlamydia was rarely transmitted through other routes, oral sex and other practices in which condoms are either not used or not useful make an important contribution to transmission of primary syphilis and urethral gonorrhoea. This is particularly important given the high rates of asymptomatic infection reported in this group, as reported by Duareva-Vizule *et al*⁵ and under-testing of affected anatomical sites.⁶

What should be provided within a sexual health service? With many shifts in commissioning and the structure of sexual health services, there is a growing need to use audit in assessing the value of services. In this month's educational article, Sonnex⁷ explores the use of colposcopy in the assessment of common gynaecological complaints in the genitourinary clinic setting. Burton *et al*⁸ report on the use of text reminders to encourage re-attendance—by contrast with other studies, they find little effect in their setting, which already had a high re-attendance rate.

Prevention needs to be epidemiologically driven. Sometimes this can be relatively straightforward, as we see in a reassuring report from the Netherlands suggesting a good degree of hepatitis B control has been achieved amongst MSM by vaccination levels of 50%.⁹ However in this month's Programme Science article¹⁰ Mishra *et al* emphasise the importance of distinguishing between the sources of HIV transmission, and the distribution of new diagnoses. They suggest that interventions are likely to be more effective when directed by transmission source than by the distribution of new diagnoses.

Other papers include a report on successful quality assurance for point of care testing in the Brazilian Amazon,¹¹ suboptimal adherence to antibiotic treatment in urethritis,¹² the distribution of pathogens in paraurethral gland infection,¹³ *Gardnerella vaginalis* in lactobacillus predominant vaginal flora,¹⁴ susceptibilities of ureaplasmas and mycoplasmas,¹⁵ preferences for microbicides,¹⁶ antenatal care in Ecuador¹⁷ and HPV detection in adolescent women.

Competing interests None.

Provenance and peer review Commissioned; not peer reviewed.

REFERENCES

- McGrath-Lone L, Marsh K, Hughes G, *et al*. The sexual health of male sex workers in England: analysis of cross-sectional data from genitourinary medicine clinics. *Sex Transm Infect* 2014;90:38–40.
- Eccles C, Clarke J. Levels of advertised unprotected vaginal and oral sex by independent indoor female sex workers in West Yorkshire, UK. *Sex Transm Infect* 2014;90:36–37.
- Grov C, Rendina HJ, Breslow AS, *et al*. Characteristics of men who have sex with men (MSM) who attend sex parties: results from a national online sample in the USA. *Sex Transm Infect* 2014;90:26–32.
- Nash JL, Hocking JS, Read TRH, *et al*. Contribution of sexual practices (other than anal sex) to bacterial sexually transmitted infection transmission in men who have sex with men: a cross-sectional analysis using electronic health records. *Sex Transm Infect* 2014;90:55–57.
- Duareva-Vizule S, Haar K, Sailer A, *et al*. Prevalence of pharyngeal and rectal Chlamydia trachomatis and Neisseria gonorrhoeae infections among men who have sex with men in Germany. *Sex Transm Infect* 2014;90:46–51.
- van Liere GAFS, Hoebe CJPA, Dukers-Muijers NHTM. Evaluation of the anatomical site distribution of chlamydia and gonorrhoea in men who have sex with men and in high-risk women by routine testing: cross-sectional study revealing missed opportunities for treatment strategies. *Sex Transm Infect* 2014;90:58–60.
- Sonnex C. Providing a genitourinary medicine colposcopy service. *Sex Transm Infect* 2014;90:8–10.
- Burton J, Brook G, McSorley J, *et al*. The utility of short message service (SMS) texts to remind patients at higher risk of STIs and HIV to reattend for testing: a controlled before and after study. *Sex Transm Infect* 2014;90:11–13.
- Gamedara N, Weerakoon AP, Zou H, *et al*. Cross-sectional study of hepatitis B immunity in MSM between 2002 and 2012. *Sex Transm Infect* 2014;90:41–45.
- Mishra S, Pickles M, Blanchard JF, *et al*. Distinguishing sources of HIV transmission from the distribution of newly acquired HIV infections: why is it important for HIV prevention planning? *Sex Transm Infect* 2014;90:19–25.
- Benzaken AS, Bazzo ML, Galban E, *et al*. External quality assurance with dried tube specimens (DTS) for point-of-care syphilis and HIV tests: experience in an indigenous populations screening programme in the Brazilian Amazon. *Sex Transm Infect* 2014;90:14–18.
- Khosropour CM, Manhart LE, Colomba DV, *et al*. Suboptimal adherence to doxycycline and treatment outcomes among men with non-gonococcal urethritis: a prospective cohort study. *Sex Transm Infect* 2014;90:3–7.
- Fan W, Zhang Q, Jiang T. Pathogen profile in men with inflammation of paraurethral glands. *Sex Transm Infect* 2014;90:52–54.
- Schwabke JR, Flynn MS, Rivers CA. Prevalence of Gardnerella vaginalis among women with lactobacillus-predominant vaginal flora. *Sex Transm Infect* 2014;90:61–3.
- Vargović M, Pasini M, Papić N, *et al*. Antimicrobial susceptibility of Ureaplasma urealyticum and Mycoplasma hominis. *Sex Transm Infect* 2014;90:69.
- Galea JT, Kinsler JJ, Imrie J, *et al*. Rectal douching and implications for rectal microbicides among populations vulnerable to HIV in South America: a qualitative study. *Sex Transm Infect* 2014;90:33–5.
- Sánchez-Gómez A, Grijalva MJ, Silva-Aycaguer LC, *et al*. HIV and syphilis infection in pregnant women in Ecuador: prevalence and characteristics of antenatal care. *Sex Transm Infect* 2014;90:70–5.