CrossMark

Highlights from this issue

Jackie A Cassell, *Editor in Chief*

doi:10.1136/sextrans-2015-052457

Throughout human history, military conflict and the presence of armies have been associated with higher rates of STI. This relates to many drivers of increased transmission – displacement and migration of populations, growth of sex work and sexual violence to name but a few. However times are changing, at least in the standing military populations of developed countries. In this month's issue, Harbertson *et al* report on the sexual behaviour and STI experience of recently deployed shipboard US military personnel.¹ The picture is complex, and is discussed in an accompanying editorial by Joel Gaydos and colleagues² which reflects on the implications for providers and policymakers. Their recommendations focus on the need for all military medical practitioners to be able to take travel and sexual histories. An interesting model of military sexual health care is presented by Desai *et al*,³ who describe the re-organisation of sexual health care for UK military overseas. The service for British forces in Germany was reconfigured into a nurse-led model – the authors describe the new hub and spoke service and compare its performance to the previous model of care.

Chemsex – sex under the influence of psychoactive substances – continues to be a hot topic, as you will see if you follow us on Twitter @sti_bmj. This month we publish in print a study by Bourne⁴ which describes the risk propensity of chemsex episodes compared with other sexual encounters among British men who have sex with men (MSM). This report has had extensive coverage in the press and was discussed in the *BMJ* – you can listen to the authors' podcast on our website at sti.bmj.com. The transmission risks that cluster with chemsex are demonstrated in Gilbert *et al*'s study⁵ which describes the epidemiological and behavioural features of an outbreak of *Shigella flexneri* 3a in England and Wales. A high proportion of those affected were HIV positive, and sex parties organised through social media were a common experience. In this month's BASHH column, Clutterbuck reflects on the role of the BASHH MSM special interest group in responding to the emerging and changing health needs of our MSM community, emphasizing particularly the importance of integrated and contraceptive focussed services to have capacity to respond to the needs of MSM.

African communities, at home or in diaspora, are a strong theme this month. In a qualitative study⁶ De Jesus *et al* describe the differing attitudes and experiences to HIV

testing of African-American and East African women in the US, emphasizing the implications for targeted testing and screening. Adebajo and colleagues explore differing approaches to the offering of HIV testing to MSM and drug using men in Nigeria, and report improved results with peer led counselling and testing.⁷ The role of heterosexual anal sex in sexual risk is often overlooked, so it is good to see Mtenga *et al*'s qualitative study of practices and beliefs about anal sex in rural Tanzania, which raises some important issues for sex education and health promotion.⁸ Finally Heaton *et al* describe the impact of the US PEPFAR (President's Emergency Plan for AIDS Relief) on HIV prevention and treatment in Africa.⁹

Innovative approaches to partner notification and the use of near patient testing are closely intertwined, and bound up with the increasingly wide range of settings where STIs are diagnosed. Estcourt *et al*¹⁰ describe a pilot randomised controlled trial of accelerated partner therapy for people with chlamydia diagnosed in primary care. This was a challenging study, and the authors reflect on the need for service level randomisation if robust evaluations of such interventions are to be achieved. In terms of our readiness to offer quick answers to index patients and partners, Brook's state-of-the-art review of currently available rapid tests for chlamydia will provide a helpful overview for the many clinicians who are seeking to configure efficient and safe patient pathways.¹¹

The HIV cascade is increasingly seen as the key measure of health system response to HIV – this month we report a study from the Netherlands which reports generally good linkage to care, but still some significant delays in newly diagnosed individuals.¹² The serology of STIs is explored in two studies on *Trichomonas vaginalis*¹³ and *Mycoplasma genitalium*.¹⁴

Finally, we have the answer to two big questions: were the Olympic and Paralympic games associated with increased clinic attendances?¹⁵ And is the incidence of STI greater in big cities?¹⁶

Competing interests None.

Provenance and peer review Commissioned; Not peer reviewed.

REFERENCES

- 1 Harbertson J, Scott PT, Moore J, *et al*. Sexually transmitted infections and sexual behaviour of deploying shipboard US military personnel: a cross-sectional analysis. *Sex Transm Inf* 2015;91:581–8.

- 2 Gaydos JC, McKee KT Jr, Faix DJ. Sexually transmitted infections in the military: new challenges for an old problem. *Sex Transm Inf* 2015;91:536–7.
- 3 Desai M, Littler JE, Samuel M, *et al*. How to set up a remotely supported hub and spoke sexual health service for a military population. *Sex Transm Inf* 2015;91:545–7.
- 4 Bourne A, Reid D, Hickson F, *et al*. Illicit drug use in sexual settings ('chemsex') and HIV/STI transmission risk behaviour among gay men in South London: findings from a qualitative study. *Sex Transm Inf* 2015;91:564–8.
- 5 Gilbert VL, Simms I, Jenkins C, *et al*. Sex, drugs and smart phone applications: findings from semistructured interviews with men who have sex with men diagnosed with *Shigella flexneri* 3a in England and Wales. *Sex Transm Inf* 2015;91:598–602.
- 6 De Jesus M, Carrete C, Maine C, *et al*. Attitudes, perceptions and behaviours towards HIV testing among African-American and East African immigrant women in Washington, DC: implications for targeted HIV testing promotion and communication strategies. *Sex Transm Inf* 2015;91:569–75.
- 7 Adebajo S, Eluwa G, Njab J, *et al*. Evaluating the effect of HIV prevention strategies on uptake of HIV counselling and testing among male most-at-risk-populations in Nigeria; a cross-sectional analysis. *Sex Transm Inf* 2015;91:555–60.
- 8 Mtenga S, Shamba D, Wamoyi J, *et al*. How long-distance truck drivers and villagers in rural southeastern Tanzania think about heterosexual anal sex: a qualitative study. *Sex Transm Inf* 2015;91:576–80.
- 9 Heaton LM, Bouey PD, Fu J, *et al*. Estimating the impact of the US President's Emergency Plan for AIDS Relief on HIV treatment and prevention programmes in Africa. *Sex Transm Inf* 2015;91:615–20.
- 10 Estcourt CS, Sutcliffe LJ, Copas A, *et al*. Developing and testing accelerated partner therapy for partner notification for people with genital Chlamydia trachomatis diagnosed in primary care: a pilot randomised controlled trial. *Sex Transm Inf* 2015;91:548–54.
- 11 Brook G. The performance of non-NAAT point-of-care (POC) tests, rapid NAAT tests for chlamydia, gonorrhoea infections. An assessment of currently available assays. *Sex Transm Inf* 2015;91:539–44.
- 12 van Veen MG, Trienekens SCM, Heijman T, *et al*. Delayed linkage to care in one-third of HIV-positive individuals in the Netherlands. *Sex Transm Inf* 2015;91:603–9.
- 13 Ton Nu PA, Rappelli P, Dessi D, *et al*. Kinetics of circulating antibody response to *Trichomonas vaginalis*: clinical and diagnostic implications. *Sex Transm Inf* 2015;91:561–3.
- 14 Idahl A, Jurstrand M, Olofsson JJ, *et al*. Mycoplasma genitalium serum antibodies in infertile couples and fertile women. *Sex Transm Inf* 2015;91:589–91.
- 15 Hall V, Charlett A, Hughes G, *et al*. Olympics and Paralympics 2012 mass gathering in London: time-series analysis shows no increase in attendances at sexual health clinics. *Sex Transm Inf* 2015;91:592–7.
- 16 Patterson-Lomba O, Goldstein E, Gómez-Liévano A, *et al*. Per capita incidence of sexually transmitted infections increases systematically with urban population size: a cross-sectional study. *Sex Transm Inf* 2015;91:610–4.