Methods We analysed follow up (FU) data from women with a CT infection who visited the STI clinic of Amsterdam, the Netherlands, from September 2015 through June 2016. After giving informed consent, participants underwent baseline and three FU speculum examinations to obtain cervical swabs for both CT culture and NAAT testing. Speculum examinations were scheduled at 7, 21 and 49 days after treatment (single dose 1000 mg azithromycin). Collected samples were analysed using a RNA and DNA-based NAAT. CT cell culture was performed on all samples at baseline, and in FU samples that were NAAT-positive. Clearance was defined as conversion to negative NAAT results at any FU visit.

Results We included 78 women with NAAT proven CT infection prior to receiving treatment of whom 58 (74%) were also culture positive. At the first visit after treatment (median 7 days; IQR 7–8) 44 (47%) women were NAAT positive, of whom three tested also positive by culture. CT infection was cleared in 73 women (94%), of whom 61 (78%) at their second FU visit (median 21 days; IQR 21–25). Of the five women who did not clear their infection, three were also culture positive indicating a viable infection. All five reported unprotected sexual contact after inclusion prior to their last FU visit, indicating potentially new infections.

Conclusion We observed prolonged and intermittent positive results over time for both NAAT tests. For three participants (4%) viable CT infections were detected 49 days after treatment. All three cases reported new sexual contacts. In conclusion, persisting infections or treatment failure were rare.

Support: Hologic provided Aptima test materials and kits in-kind. Roche provided Cobas test materials and kits in-kind. Copan provided Universal Transport Medium in-kind

P3.19 RISK FACTORS FOR HIV INFECTION AMONG FEMALE COMMERCIAL SEX WORKERS IN BANGUI, CENTRAL AFRICAN REPUBLIC

¹Belec, ²Jean De Dieu Longo. ¹Paris V University, Paris, France; ²Centre National de Référence Des Maladies Sexuellement Transmissibles Et de La Thérapie Antirétrovir, Bangui – Central African Republic

10.1136/sextrans-2017-053264.256

Introduction The categorization of female commercial sex work according socio-anthropologic criteria constitutes a prerequisite to assess differential risks of HIV exposure.

Methods A cross-sectional questionnaire survey was conducted to describe the spectrum of commercial sex work in Bangui, the capital city of the Central African Republic, among 345 sexually active women having more than 2 sexual partners other than their regular partner during the last 3 months and reporting to have received money or “gifts” in return of their sexual relationships.

Results HIV infection in study female sex worker (FSW) population was strongly associated with anal sex practice with last clients (OR, 4.3), irregular condom use in last 3 months (OR, 24.9), and alcohol consumption before sex (adjusted OR, 2.8). Networks of commercial sex work comprised six different FSW categories, including two groups of “official” professional FSW primarily classified according to their site of work [i) “kata” (18.6%) representing women working in poor neighbourhoods of Bangui; ii) “pupulenge” (13.9%) working in

hotels and night clubs to seek White men] and four groups of “clandestine” nonprofessional FSW classified according to their reported main activity [i) “market and street vendors” (20.8%); ii) “schoolgirls or students” (19.1%) involved in occasional transactional sex (during holidays); iii) “housewives or unemployed women” (15.7%); “civil servants” (11.9%) working as soldiers or in public sector]. HIV varied according to FSW categories. HIV prevalence was 6-fold higher among “kata” than “pupulenge” (39.1% versus 6.3%). “Students”, “civil servants” and “housewives” were the less HIV-infected (6.1%, 9.8%, 13.0%, respectively), whereas “sellers” constituted the category of highest HIV prevalence (31.9%).

Conclusion Our observations highlight the high level of vulnerability of both poor professional “kata” and nonprofessional “street vendors” FSW categories which should be particularly taken in account when designing prevention programs for STIs/HIV control purposes.

P3.20 PREVALENCE OF *CHLAMYDIA TRACHOMATIS*, *NEISSERIA GONORRHOEAE* AND *TRICHOMONAS VAGINALIS* IN FEMALE SEX WORKERS IN MOROCCO

¹B Bellaji, ¹A Hançali, ¹S Jennane, ¹H Oukouchoud, ¹C Ouanaim, ²A Bennani, ²A Latifi, ¹R Charof. ¹National Institute of Hygiene, Rabat, Morocco; ²National Aids Program, Ministry of Health, Rabat, Morocco

10.1136/sextrans-2017-053264.257

Introduction The incidence of reported sexually transmitted infections (STIs) in Morocco is over than 350.000 per year. The management of cases is based on syndromic approach especially in women. While female sex workers (FSWs) are assumed to be at increased risk of STIs, there are limited comparative data with other population groups available. Two studies have been conducted between 2013–2014 in different cities in Morocco in order, to explore the prevalence of *Chlamydia trachomatis* (CT), *Neisseria gonorrhoeae* (GC), and *Trichomonas vaginalis* (TV) among FSWs and to compare them with prevalences found in women consultant in family planning units (FPU).

Methods A sample of 519 FSWs and 537 asymptomatic women consultant in FPU was recruited in basic health services and NGOs. All consenting FSWs and women consultant in FPU underwent pretest counselling and provided socio-demographic and behavioural data using a structured questionnaire. The women were also asked to provide vaginal and cervical specimens to detect the respective STIs. GC identification was performed by culture and PCR, CT was detected by PCR and TV was detected by culture.

Results The prevalence of CT, GC, and TV were 20,7%, 9,35%, and 13,3%, respectively, in the FSWs, compared with 3%, 0,4%, and 5,6% respectively in the women consultant in FPU. These results show a high prevalence of CT, GC and TV in Moroccan FSWs than women consultant in FPU.

Conclusion Most STIs prevalences are lower in comparison with prevalences found for FSWs in other countries. However, the National Aids Program conclude that continued close monitoring of the prevalence of CT, GC and TV infection in FSWs is important for preventing the dissemination of these microorganisms, and that further investigation of CT as a sexually transmitted pathogen in women is needed.