pregnancy and infant outcomes and may also increase the risk of HIV mother-to-child-transmission. Additionally, these infections may be vertically transmitted to neonates, resulting in colonisation, conjunctivitis, and pneumonia. There is a lack of data on the neonatal incidence of CT, NG and TV due to lack of systematic screening of mothers and neonates and reliance on STI syndromic management. We conducted an analysis to determine rates of vertical transmission of CT, NG and TV from HIV-infected women to their neonates.

Methods We tested HIV-infected pregnant women for CT, NG and TV on vulvo-vaginal swab specimens during their first clinic visit post-delivery. Neonates of all women who tested positive for an STI were also evaluated for these STIs using nasopharyngeal swab specimens collected the same day. Testing was done using the Xpert CT/NG and Xpert TV [Cepheid, Sunnyvale, CA].

Results Of the 54 women tested for CT, NG and TV postnatally, 27 (50%) had at least one STI. Twenty three (85.2%) of the 27 neonates born to women with STIs were evaluated for STI colonisation. Overall, 18 (78.3%) neonates tested positive for at least one STI; 77.3% (17/22) neonates born to mothers with CT infection, 50% (2/4) with TV, and none with NG tested positive for those respective infections. STI colonisation rates also differed by birth weight with 100% of low birth weight, 77.8% (14/18) with normal birth weight, and 50% (1/2) of high birth weight neonates testing positive for at least one of those infections. Rates of neonatal STI colonisation (83.3%) were also high among the six infants delivered by C-section.

Conclusion Our analysis revealed high rates STIs among post-partum HIV-infected women, coupled with high rates of neonatal STI colonisation, particularly among low birth weight neonates.

P3.142 EPIDEMIOLOGICAL AND CLINICAL PROFILE OF HIV PATIENTS OF A PUBLIC HEALTH UNIT FROM BELO HORIZONTE, MINAS GERAIS

MI Maciel, SVF Carvalho, LM Gomes, JM Freire, CG Coelho. *Municipal Health Department, Belo Horizonte, Brazil, Belo Horizonte – MG, Brazil*

10.1136/sextrans-2017-053264.377

Introduction More than 8 00 000 individuals are HIV-infected in Brazil and a small increase of new infections (4%) was observed in past years. Nowadays, more than half (64%) of people living with HIV in Brazil are receiving antiretroviral treatment (ARVT). The epidemiological and clinical profiles of HIV patients in medium-sized Brazilian cities are unknown, and it is necessary to study them to allow comparisons. The present study aims to identify the profile of HIV-infected individuals in a public health unit, from 2013 to 2016.

Methods A descriptive study was carried out in a public health care unit, which has specialised care for HIV-infected and sexually transmitted infections (STIs). The population of interest in this study were patients (319 patients) infected with HIV and under ARVT. Individuals whose registry was in duplicate (n=1), whose ARVT started before 2013 (n=8), and those who abandoned ARVT (n=26) were excluded. Demographic data (age, sex and self-identified sexual identity) and information on the patients' clinical situation (HIV diagnosis year, ARVT start date, presence of undetectable viral load (VL) and time to undetectable VL) were obtained. Descriptive analysis

of quantitative variables was performed using the Stata software, version 12.0.

Results Among the 285 patients included in the study, most of them were male (89.44%), aged 20–29 years (47.35%) and self-identified their sexual identity as homosexual (69.01%). Most of the diagnoses were performed in the year 2015 (39.08%) and the majority of the patients had an undetectable VL (78.95%). The mean time for VL to become and remain undetectable after starting ARVT was 7.49 (+6.74) months.

Conclusion The present study reflects the epidemiology of HIV infection in Brazil, with concentration of the disease in young adults and men who have sex with men. The virological response to ARVT is directly related to adherence to regular treatment and should be the target investment of health services specialised to people with HIV/AIDS.

P3.143 SEROLOGICAL PREVALENCE OF ANTIBODIES TO HUMAN IMMUNODEFICIENCY VIRUS AMONG NEPALESE POPULATION

¹Manoj Kumar Sah, ²Prabhakar Verg Shah, ³Rupa Adhikari, ⁴Shyam Kumar Mishra, ⁴Keshab Parajuli. ¹Kathmandu University, Kantipur Dental College Teaching Hospital and Research Centre, Kathmandu, Nepal; ²B and B Hospital, Kathmandu, Nepal; ³Namaste Nepal Medical Centre, Kathmandu, Nepal; ⁴Tribhuvan University Teaching Hospital, Kathmandu, Nepal

10.1136/sextrans-2017-053264.378

Introduction HIV is one of the major public health problems in recent time. Because of limited work opportunities in Nepal, seasonal labour migration especially disadvantaged groups to overseas become common. Nepalese government is promoting overseas employment opportunities for its citizens because of poor absorption of growing labour force in the country. This study was aimed to determine the serological prevalence of antibodies Human Immunodeficiency Virus (HIV) among the Nepalese population intended to depart overseas.

Methods This study was conducted at Namaste Nepal Medical Centre and Polyclinic, Kathmandu, Nepal. Twenty Nine Thousand Four Hundred Forty One healthy populations were included for this study over a period from January 2013 to December 2014. The venous blood was collected aseptically and subjected to screening for anti HIV 1/2 (HIV Tri Dot, Jay Mitra, India) and confirmed by third generation HIV 1/2 ELISA (SD, Bio Lines, Korea) prior to overseas migration for labour employment. Data were analysed by using SPSS version 17.0 software.

Results The serological prevalence of antibodies to HIV 1/2 were found to be 0.04% (11/29,441). Among the HIV infected population 0.03% (9/29,441) was male while only 0.01% (2/29,441) was female. Majority of the HIV infected people with HIV 1/2 was found to be 20–30 years old and belonging from disadvantaged groups like Dalit, Janajati and Kshetri.

Conclusion The HIV seroprevalence showed decreasing trend, due to the cumulative effect of increasing awareness of HIV and better HIV testing and counselling (HTC) in Nepal. This study reflects that it would be alarming situation because young age population becomes more infected. Therefore, a continuous surveillance, advocacy and social mobilisation need to be considered for further improvements of transmission of HIV infection.

P3.144 EVALUATING THE PREVALENCE AND KNOWLEDGE OF SEXUALLY TRANSMITTED INFECTION AMONG HOMELESS PERSONS IN CENTRAL BRAZIL

Marcos André de Matos, Karlla Antonieta Amorim Caetano, Sheila Araujo Teles, Raquel Silva Pinheiro, Paulie Marcelly Ribeiro dos Santos Carvalho, Luana Rocha da Cunha Rosa, Márcia Maria de Souza, Ana Luiza Neto Junqueira, Brenda Kelly Gonçalves Nunes, Andressa Cunha Paula, Thaynara Ferreira de Amorim. Universidade Federal De Goiás, Goiânia – GO, Brazil

10.1136/sextrans-2017-053264.379

Introduction The objective of this study was to estimate the prevalence of Sexually Transmitted Infections and to explore the general understanding of transmission and prevention of these infections among the homeless in Goiânia, Central Brazil.

Methods This is a cross-sectional study. Between August 2014 and June 2015, residents of a public homeless shelter were recruited in Goiânia, Goiás, in Central Brazil. Survey items included demographics and knowledge of STIs. Blood samples were submitted to rapid tests for HIV, syphilis, and hepatitis B (HBV), and a positive result on any one or more of these tests indicated the presence of an STI. This study was approved by the Research Ethics Committee of the Federal University of Goiás.

Results The sample included 209 homeless persons. The mean age of participants was 38 years. The prevalence of STI was 27.8% (58/209), for HIV 3.3% (7/209), syphilis 24.9% (52/209) and hepatitis B 1.0% (2/209). When asked about transmission of HIV, syphilis, and hepatitis B in public restrooms, 4.8%, 8.6% and 10.5% reported that this can occur, respectively. On syringe sharing as an IST transmission route, 78%, 11.5% and 23.9% believe that HIV, syphilis and hepatitis B can be acquired through this parenteral media, respectively. The male condom was investigated as a method of prevention for these sexual infections, and 73.7%, 38.3% and 14.4% affirmed the efficacy of condom use for protection against HIV, syphilis and hepatitis B, respectively. It was found that age (PR: 1.01; 95% CI: 1.00–1.03) and confirming HIV transmission through syringe sharing (PR: 0.47; 95% CI: 0.27–0.81) were statistically significant associated with STI positivity ($p \leq 0.05$).

Conclusion The dissemination of knowledge about STIs is an important tool for breaking the chain of transmission of sexual infections. In Brazil, access of the homeless population to health and sexuality education services can be improved using mobile health teams, known as Street Outreach Offices.

P3.145 GENITAL CHLAMYDIA TRACHOMATIS INFECTIONS AMONG PATIENTS ATTENDING THE UNIVERSIDAD DE BUENOS AIRES HOSPITAL DURING 2010–2015

Gallo Vaulet Maria Lucia, L Amaya, M Mestre, V Gualtieri, ML Gallo Vaulet, AC Entrocassi. Universidad de Buenos Aires, Facultad De Farmacia Y Bioquímica, Depto. Bioquímica Clínica, Buenos Aires, Argentina

10.1136/sextrans-2017-053264.380

Introduction: *Chlamydia trachomatis* is the most prevalent bacterial sexually transmissible infection (STI) worldwide. Since the majority of the cases remain asymptomatic, lack of treatment can lead to serious sequels, principally in women. Previous studies at the University Hospital (period 2001–2009) reported a *C. trachomatis* prevalence of 2.1%. The aim of

this study was to analyse the prevalence of *C. trachomatis* infections in patients attending the University Hospital in the 2010–2015 period.

Methods All genital samples (cervical swabs, male urethral swabs and first void urine) taken from patients attending the University Hospital laboratory between 1/1/2010 and 31/12/2015 with indication of *C. trachomatis* detection were included in this study. *C. trachomatis* was detected by ompA nested PCR. Genotyping of *C. trachomatis* was done by PCR-RFLP.

Results A total of 2664 samples (1998 feminine and 666 masculine) were included in this study. Global prevalence of *C. trachomatis* was 2.25% (60/2664), 5.55% in males (37/666) and 1.15% (23/1998) in females. Among *C. trachomatis* positive patients, 56.8% of male (21/37) and 26.1% of female (6/23) patients were symptomatic. *C. trachomatis* prevalence was 13% at the STI clinic, 3.4% at the OBGYN clinic and 0.5% at the fertility program. Among male infected patients, 63.6% were under 30 years, and 47.8% of infected women were under 25 years. The most frequently detected genotype was E (35%) followed by D (21%) and F (13%).

Conclusion Global prevalence of *C. trachomatis* did not change respect the 2001–2009 period. Comparing to 2001–2009 period, an increase in the frequency of infection in males was observed (5.55% vs 3.29% $p=0.07$). Difference was observed in prevalence of infection in the OBGYN clinic vs Fertility Program (3.4% vs 0.5% $p<0.05$). Genotype distribution is similar to previously reported in this population and does not differ from international distribution. According to our results vulnerable populations for *C. trachomatis* infection are young and male symptomatic patients.

P3.146 FIRST BRAZILIAN NATIONAL ANTIMICROBIAL SUSCEPTIBILITY SURVEILLANCE FOR NEISSERIA GONORRHOEAE

¹Maria Luiza Bazzo, ¹Lisléia Golfetto, ²Miriam Franchini, ²Pâmela Cristina Gaspar, ³Ana Flávia Pires, ⁴Ligia MariaBedeschi Costa, ⁵Mauro Cunha Ramos, ⁶Loeci Natalina Timm, ⁷William Antunes Ferreira, ⁸Purificação Pereira da Silva Maria Da, ⁹Carvalho da Silva Roberto José, ¹⁰Fonseca Andrade Lidiane da, ¹¹Fátima Mendes Pereira Lúcia de, ¹Felipe de Rocco, ¹Jéssica Martins, ¹Hanaldyia Machado, ¹Marcos André Schörner, ¹Thais Mattos Santos, ⁴Faria de Carvalho Simone Veloso, ⁴Luciane Guimarães Dias, ⁶Letícia Eidt, ⁶de Oliveira Amhold Guilherme Henrique, ⁶Souza Coelho Muniz Chayane Ariel, ⁷Waldemara de Souza Vasconcelos, ⁷Jairo de Souza Gomes, ⁷Pinto Da Silva Maria De Fátima, ⁸Rosan Barboza de Matos, Cláudio Campos do, ⁹Porto, ¹²Leonor Henriette de Lannoy, ²Adele Schwartz Benzaken. ¹Federal University of Santa Catarina, Florianópolis – SC, Brazil; ²Department of Surveillance, Prevention and Control of STI, HIV/AIDS and Viral Hepatitis, Brasília – DF, Brazil; ³Postgraduate Program In Collective Health Of University of Brasília, Brasília – DF, Brazil; ⁴Municipal Secretary of Health Belo Horizonte, Belo Horizonte – MG, Brazil; ⁵Brazilian Society of STD, Porto Alegre – RS, Brazil; ⁶State Department of Health of Rio Grande do Sul, Porto Alegre – RS, Brazil; ⁷Alfredo da Mata Foundation, Manaus – AM, Brazil; ⁸State Centre Specialised in Diagnosis, Assistance And Research, Salvador – BA, Brazil; ⁹STD Reference and Training Centre of São Paulo, São Paulo – SP, Brazil; ¹⁰Testing and Counselling Centre of Distrito Federal, Brasília – DF, Brazil; ¹¹Central Public Health Laboratory of Distrito Federal, Brasília – DF, Brazil; ¹²Day Hospital of Brasília, Brasília – DF, Brazil

10.1136/sextrans-2017-053264.381

Introduction The threat of multidrug resistant *Neisseria gonorrhoeae* (NG) is a concern worldwide, especially in settings with emerging resistance to the extended spectrum cephalosporins. Since 2009 WHO recommendation to the countries to perform Antimicrobial Surveillance has been reinforced. Brazilian's sexually transmitted infection guideline recommends