P3.144 EVALUATING THE PREVALENCE AND KNOWLEDGE OF SEXUALLY TRANSMITTED INFECTION AMONG HOMELESS PERSONS IN CENTRAL BRAZIL

Marcos André de Matos, Karlla Antonieta Amorim Caetano, Sheila Araujo Teles, Raquel Silva Pinheiro, Paulie Marcelly Ribeiro dos Santos Carvalho, Luana Rocha da Cunha Rosa, Márcia Maria de Souza, Ana Luiza Neto Junqueira, Brenda Kelly Gonçalves Nunes, Andressa Cunha Paula, Thaynara Ferreira de Amorim. Universidade Federal De Goiás, Goiânia – GO, Brazil

10.1136/sextrans-2017-053264.379

Introduction The objective of this study was to estimate the prevalence of Sexually Transmitted Infections and to explore the general understanding of transmission and prevention of these infections among the homeless in Goiânia, Central Brazil.

Methods This is a cross-sectional study. Between August 2014 and June 2015, residents of a public homeless shelter were recruited in Goiânia, Goiás, in Central Brazil. Survey items included demographics and knowledge of STIs. Blood samples were submitted to rapid tests for HIV, syphilis, and hepatitis B (HBV), and a positive result on any one or more of these tests indicated the presence of an STI. This study was approved by the Research Ethics Committee of the Federal University of Goiás.

Results The sample included 209 homeless persons. The mean age of participants was 38 years. The prevalence of STI was 27.8% (58/209), for HIV 3.3% (7/209), syphilis 24.9% (52/209) and hepatitis B 1.0% (2/209). When asked about transmission of HIV, syphilis, and hepatitis B in public restrooms, 4.8%, 8.6% and 10.5% reported that this can occur, respectively. On syringe sharing as an IST transmission route, 78%, 11.5% and 23.9% believe that HIV, syphilis and hepatitis B can be acquired through this parenteral media, respectively. The male condom was investigated as a method of prevention for these sexual infections, and 73.7%, 38.3% and 14.4% affirmed the efficacy of condom use for protection against HIV, syphilis and hepatitis B, respectively. It was found that age (PR: 1.01; 95% CI: 1.00–1.03) and confirming HIV transmission through syringe sharing (PR: 0.47; 95% CI: 0.27–0.81) were statistically significant associated with STI positivity ($p \leq 0.05$).

Conclusion The dissemination of knowledge about STIs is an important tool for breaking the chain of transmission of sexual infections. In Brazil, access of the homeless population to health and sexuality education services can be improved using mobile health teams, known as Street Outreach Offices.

P3.145 GENITAL CHLAMYDIA TRACHOMATIS INFECTIONS AMONG PATIENTS ATTENDING THE UNIVERSIDAD DE BUENOS AIRES HOSPITAL DURING 2010–2015

Gallo Vaulet Maria Lucia, L Amaya, M Mestre, V Gualtieri, ML Gallo Vaulet, AC Entrocassi. Universidad de Buenos Aires, Facultad De Farmacia Y Bioquímica, Depto. Bioquímica Clínica, Buenos Aires, Argentina

10.1136/sextrans-2017-053264.380

Introduction: *Chlamydia trachomatis* is the most prevalent bacterial sexually transmissible infection (STI) worldwide. Since the majority of the cases remain asymptomatic, lack of treatment can lead to serious sequels, principally in women. Previous studies at the University Hospital (period 2001–2009) reported a *C. trachomatis* prevalence of 2.1%. The aim of

this study was to analyse the prevalence of *C. trachomatis* infections in patients attending the University Hospital in the 2010–2015 period.

Methods All genital samples (cervical swabs, male urethral swabs and first void urine) taken from patients attending the University Hospital laboratory between 1/1/2010 and 31/12/2015 with indication of *C. trachomatis* detection were included in this study. *C. trachomatis* was detected by ompA nested PCR. Genotyping of *C. trachomatis* was done by PCR-RFLP.

Results A total of 2664 samples (1998 feminine and 666 masculine) were included in this study. Global prevalence of *C. trachomatis* was 2.25% (60/2664), 5.55% in males (37/666) and 1.15% (23/1998) in females. Among *C. trachomatis* positive patients, 56.8% of male (21/37) and 26.1% of female (6/23) patients were symptomatic. *C. trachomatis* prevalence was 13% at the STI clinic, 3.4% at the OBGYN clinic and 0.5% at the fertility program. Among male infected patients, 63.6% were under 30 years, and 47.8% of infected women were under 25 years. The most frequently detected genotype was E (35%) followed by D (21%) and F (13%).

Conclusion Global prevalence of *C. trachomatis* did not change respect the 2001–2009 period. Comparing to 2001–2009 period, an increase in the frequency of infection in males was observed (5.55% vs 3.29% $p=0.07$). Difference was observed in prevalence of infection in the OBGYN clinic vs Fertility Program (3.4% vs 0.5% $p<0.05$). Genotype distribution is similar to previously reported in this population and does not differ from international distribution. According to our results vulnerable populations for *C. trachomatis* infection are young and male symptomatic patients.

P3.146 FIRST BRAZILIAN NATIONAL ANTIMICROBIAL SUSCEPTIBILITY SURVEILLANCE FOR NEISSERIA GONORRHOEAE

¹Maria Luiza Bazzo, ¹Lisléia Golfetto, ²Miriam Franchini, ²Pâmela Cristina Gaspar, ³Ana Flávia Pires, ⁴Ligia MariaBedeschi Costa, ⁵Mauro Cunha Ramos, ⁶Loeci Natalina Timm, ⁷William Antunes Ferreira, ⁸Purificação Pereira da Silva Maria Da, ⁹Carvalho da Silva Roberto José, ¹⁰Fonseca Andrade Lidiane da, ¹¹Fátima Mendes Pereira Lúcia de, ¹Felipe de Rocco, ¹Jéssica Martins, ¹Hanaldyia Machado, ¹Marcos André Schörner, ¹Thais Mattos Santos, ⁴Faria de Carvalho Simone Veloso, ⁴Luciane Guimarães Dias, ⁶Letícia Eidt, ⁶de Oliveira Amhold Guilherme Henrique, ⁶Souza Coelho Muniz Chayane Ariel, ⁷Waldemara de Souza Vasconcelos, ⁷Jairo de Souza Gomes, ⁷Pinto Da Silva Maria De Fátima, ⁸Rosan Barboza de Matos, Cláudio Campos do, ⁹Porto, ¹²Leonor Henriette de Lannoy, ²Adele Schwartz Benzaken. ¹Federal University of Santa Catarina, Florianópolis – SC, Brazil; ²Department of Surveillance, Prevention and Control of STI, HIV/AIDS and Viral Hepatitis, Brasília – DF, Brazil; ³Postgraduate Program In Collective Health Of University of Brasília, Brasília – DF, Brazil; ⁴Municipal Secretary of Health Belo Horizonte, Belo Horizonte – MG, Brazil; ⁵Brazilian Society of STD, Porto Alegre – RS, Brazil; ⁶State Department of Health of Rio Grande do Sul, Porto Alegre – RS, Brazil; ⁷Alfredo da Mata Foundation, Manaus – AM, Brazil; ⁸State Centre Specialised in Diagnosis, Assistance And Research, Salvador – BA, Brazil; ⁹STD Reference and Training Centre of São Paulo, São Paulo – SP, Brazil; ¹⁰Testing and Counselling Centre of Distrito Federal, Brasília – DF, Brazil; ¹¹Central Public Health Laboratory of Distrito Federal, Brasília – DF, Brazil; ¹²Day Hospital of Brasília, Brasília – DF, Brazil

10.1136/sextrans-2017-053264.381

Introduction The threat of multidrug resistant *Neisseria gonorrhoeae* (NG) is a concern worldwide, especially in settings with emerging resistance to the extended spectrum cephalosporins. Since 2009 WHO recommendation to the countries to perform Antimicrobial Surveillance has been reinforced. Brazilian's sexually transmitted infection guideline recommends

dual therapy to treat gonococcal infection (ciprofloxacin plus azithromycin). However, regional studies performed in three Brazilian states demonstrated quinolone resistance. For these states it was recommended to replace the quinolones by third generation cephalosporin. The aim of this study was to perform a Brazilian national gonococcal antimicrobial surveillance.

Methods The surveillance study included seven collection sites representing five geographic regions of Brazil. A total of 550 NG isolates from male urethral discharge was sent to the gonococcal national reference laboratory for analysis. The minimum inhibitory concentration (MIC) with the agar dilution method was performed for penicillin, tetracycline, ciprofloxacin, ceftriaxone, cefixime and azithromycin.

Results The sensitivity profile of the NG isolates was performed in 131 isolates from the Southeast, 104 from the Northeast, 100 from the North, 68 from the Center-West and 147 from the South. In addition to penicillin and tetracycline, a high level of resistance for ciprofloxacin (47% to 78% of isolates) was observed in the isolates of all the regions. All the isolates were sensitive to ceftriaxone and cefixime, although one isolate was found with elevated MIC. Regarding azithromycin sensitivity, the majority of the isolates were sensitive, but the emergence of an intermediate (5% to 15%) or resistant (4% to 10%) profile needs to be monitored.

Conclusion The national survey confirmed the high level of ciprofloxacin resistance already described worldwide. These results indicate the need to urgently change Brazilian recommendation for gonorrhoeae treatment and the importance of systematic gonococcal resistance surveillance.

P3.147 PREVALENCE OF SEXUALLY TRANSMITTED INFECTIONS AND ASSOCIATED FACTORS IN WOMEN WHO HAVE SEX WITH WOMEN

MAO Ignacio, J Andrade, APF Freitas, MG Silva, MTC Duarte. São Paulo State University, Brazil

10.1136/sextrans-2017-053264.382

Introduction The present research aims to fill a gap in the national and international literature concerning prevalences and factors related to STI (Sexually Transmitted Infections) in Women who Have Sex with Women (WSW).

Methods It is a cross sectional, analytical and non-randomized study with 100 WSW from Botucatu - SP and surrounding regions who answered the call from social media, mass communication means, health services and friends or acquaintances from January to November, 2015. Data was obtained by the researchers involved in the main study, through interviews, blood tests to diagnose HIV, syphilis and hepatitis B and cervical sample collection to investigate Chlamydia Trachomatis, Neisseria Gonorrhoeae, HPV through PCR and oncotic cytology to screen Trichomonas vaginalis. Associations were estimated by multiple regression.

Results Total prevalence of STI was 35%, chlamidia infection (3.3%), HIV infection (2.0%), Trichomonas vaginalis (1.1%) and syphilis (1.0%). Neisseria Gonorrhoeae and Hepatitis B were not found. The factors associated to STI were: to be single, [OR=3,76 (CI:95%: 1.14–12.43); p=0030], use of sex toys [OR=3,87(CI:95%: 1.14–13.16); p=0030], and number of male partners in the previous year [OR=7,99 (CI:95%: 1.51–42.44); p=0015].

Conclusion Considered as whole, these data lead to conclusion that the individuals of this study presented high vulnerability to STI/AIDS, as shown by the high prevalence of STI. This study clearly shows the need for a specific health assistance to these women, promoting prevention and education in a holistic approach.

P3.148 REPORTED SEXUALLY TRANSMITTED INFECTIONS IN BRAZIL: PREVALENCE AND RISK FACTORS

¹Mariana Jorge De Queiroz, ²Meritxell Sabidó, ¹Clarissa Habkost Dutra De Barros, ¹Ana Roberta Pati Pascom, ²Enrique Galbán, ¹Adele Schwartz Benzaken. ¹Department of STI, AIDS and Viral Hepatitis, Secretary for Health Surveillance, Ministry of Health, Brasília – DF, Brazil; ²University of Girona, Catalonia, Spain; ³Facultad de Medicina, Habana, Cuba

10.1136/sextrans-2017-053264.383

Introduction Population-based studies help provide more-robust prevalence estimates and related factors, which are needed to drive sexually transmitted infection (STI) prevention and control programmes. We use data from the Brazilian Survey of Knowledge, Attitudes, and Practices (PCAP) to examine the prevalence of self-reported STIs and its association with demographics, behaviour, and the uptake of interventions.

Methods A probability sample survey was conducted in 2013 (PCAP-2013; n=12,000) and in 2008 (PCAP-2008; n=8,000) among men and women aged 15–64. The participants completed the survey in their homes through computer-assisted face-to-face interviews and self-interviews. We describe the age-specific and sex-specific prevalence of self-reported STIs in relation to demographics, behavioural factors, and the uptake of interventions and explore associated changes since the PCAP-2008.

Results Overall, 9.4% of men and 5.3% of women reported having had an STI in 2013. This prevalence decreased from that in 2008 (15.9% in men and 8.8% in women). In 2013, self-reported STIs were mainly urethral discharge (5.8%) among men and genital ulcers (2.9%) among women. Men sought care and treatment more often than women did during their most recent experience of STI symptoms (care: 85.7% and 81.2%; treatment: 77.5% and 68.7%, respectively). Self-reported STI infections were associated with increasing age, decreasing socioeconomic status, current or previous drug use, sex with a casual partner in the last 12 months (for both sexes), sex with a same sex partner, and no condom use during the last sexual intercourse (for men), and non-indigenous status and one or more previous HIV tests (for women).

Conclusion This probability sample survey highlights a decrease in STIs for men and women in Brazil. Women carry a higher STI burden than men do. The increases in health care-seeking behaviour and treatment received when presenting symptoms are encouraging. Increasing age, casual sexual partners, and drug use are the main risk factors for STI acquisition.

P3.149 SPATIOTEMPORAL DYNAMICS OF THE HIV EPIDEMIC IN BRAZIL, 2009 TO 2014

Mariana Veloso Meireles, Ana Roberta Pati Pascom. Department of STIS, AIDS And Viral Hepatitis/Ministry of Health of Brazil, Brasília – DF, Brazil

10.1136/sextrans-2017-053264.384