Conclusion Acceptability and feasibility of STI screening among pregnant women in Vietnam was very high. The prevalence of CT was much higher than the mean prevalence of CT among low and middle income countries in Asia. Study findings support policy to incorporate routine screening for STIs during pregnancy to improve maternal and infant health.

P3.160 OPTIMISING EXISTING SEXUAL HEALTH CLINICS INCREASES HIV TESTING AMONG GAY AND BISEXUAL MEN AT HIGHER RISK OF INFECTION

¹Muhammad Shahid Jamil, ¹Hamish Mcmanus, ¹Denton Callander, ¹Garrett Prestage, ¹Hammad Ali, ²Vickie Knight, ³Tim Duck, ⁴Catherine C O'Connor, ⁵Marcus Chen, ²Anna M McNulty, ¹Phillip Keen, ⁵Nick Medland, ⁶Margaret Hellard, ⁷Andrew E Grulich, ⁸David A Lewis, ¹John M Kaldor, ⁵Christopher K Fairley, ¹Basil Donovan, ¹Rebecca J Guy. ¹The Kirby Institute, Unsw Australia, Sydney, Australia; ²Sydney Sexual Health Centre, Sydney Hospital, Sydney, Australia; ³Nsw Ministry of Health, Sydney, Australia; ⁴Sexual Health Service, Community Health, Sydney Local Health District, Sydney, Australia; ⁵Melbourne Sexual Health Centre, Melbourne, Australia; ⁶Centre for Population Health, Burnet Institute, Melbourne, Australia; ⁷The Kirby Institute, Unsw Australia, Sydney, Australia, Sydney, Australia; ⁸Western Sydney Sexual Health Centre, Sydney, Australia

10.1136/sextrans-2017-053264.395

Introduction Globally, community-based HIV testing models are recommended to improve access to testing with less focus on optimising existing clinical services. In the past 5 years, public-funded sexual health clinics (SHCs) in New South Wales (NSW, Australia) have taken a range of initiatives to improve efficiencies such as triage to divert low-risk heterosexuals, express clinics, online booking, self-registration and SMS reminders. We analysed temporal trends in HIV testing among gay and bisexual men (GBM) attending SHCs in this period and assessed if testing was targeted to high-risk GBM.

Methods We used retrospective data from 32 SHCs in NSW participating in a surveillance network. HIV-negative GBM were categorised based on client type (new or existing), risk status (using partner numbers and/or recent rectal sexually transmitted infection), and recent HIV testing (past 6 months for high-risk, past 12 months for low-risk GBM). We used repeated measures Poisson regression to assess trends in attendance, tests and contribution to total tests by GBM categories.

Results From 2009–2015, unique GBM attending increased by 82% (5,477 to 9,983), and HIV tests increased by 155% (4,779 to 12,173) with significant increase in all categories and greatest increase in existing high-risk clients. Of 58,377 HIV tests done, 74% were in existing and 35% in high-risk clients. Over time, existing high-risk clients with recent testing had an increasingly larger contribution to total tests (13% annual increase, 95% CI:8%–18%, $p<0.001$). There was a simultaneous annual decline in contribution by these low-risk categories: new clients (5% decline, 95% CI:2%–7%, $p<0.001$); existing clients with no recent testing (6% decline, 95% CI:5%–7%, $p<0.001$). There were no changes in contribution by other categories (new high-risk clients; existing high-risk clients with no recent testing; existing low-risk clients with recent testing).

Conclusion SHCs in NSW have successfully increased HIV testing among GBM, with greatest increase in high-risk men. The strategies adopted could be translated to other settings.

P3.161 FACTORS ASSOCIATED WITH POSITIVE TESTS FOR HIV AND SYPHILIS CENTRE AND ADVICE

¹Nara Borges Gonçalves Lima, ¹Heber José De Moura, ²Roumayne Fernandes Vieira Andrade, ¹Elk De Assis Araújo, ¹Maria Alix Leite Araújo. ¹Universidade de Fortaleza, Fortaleza – CE, Brazil; ²Faculdade de Ciências Médicas, Campina Grande – PB, Brazil

10.1136/sextrans-2017-053264.396

Introduction Faced with rapid evolution of sexually transmitted infections (STIs), especially HIV/AIDS, as well as the impacts on society, this study aimed to analyse the factors associated with seropositivity TR for HIV and syphilis in people served in the Testing Centre and counselling - CTA Fortaleza, Ceará.

Methods A cross-sectional study that analysed data from Forms Customer Information System of people who performed the rapid test (RT) in the months of June and July 2015. We analysed sociodemographic, behavioural and epidemiological variables. The data were entered into the Statistical Package for the Social Sciences and were analysed using frequency distributions, measures of central tendency and dispersion.

Results Were analysed 882 forms. The research result shows that 622 (70.5%) were males and 414 (46.9%) were aged between 19 and 29 years, with a predominance of non-white people, without a partner, with more than four years of study, coming from Fortaleza. Of the 875 tests for HIV and 648 tests for syphilis, 49 (5.3%) and 72 (8.1%) showed reagent test result, respectively. Cases of HIV reagent were associated with the identification of the service by friend/service user ($p=0.013$), male gender ($p=0.006$), non-Caucasians ($p=0.045$), lower education of four years of study ($p=0.045$) and working ($p=0.009$). There was statistical association in positive cases of syphilis and present IST ($p<0.001$) and heterosexual sexual partners and women ($p=0.012$).

Conclusion These findings suggest the need for effective measures for control and prevention of STIs aimed at young adults, male and more than one sexual partner.

P3.162 DEVELOPING A CLINICAL PREDICTION RULE TO TARGET STI TESTING AND CONTRACEPTION TO WOMEN IN COMMUNITY SETTINGS: IMPLICATIONS FOR SEXUAL HEALTH SURVEYS

¹Natalie Edelman, ²De Visser Ro, ³Mercer Cm, ¹J Cassell. ¹Brighton and Sussex Medical School, Brighton, UK; ²University of Sussex, Brighton, UK; ³University College London, London, UK

10.1136/sextrans-2017-053264.397

Introduction Applied sexual health surveys investigate associations between psychosocial and demographic factors, sexual risk behaviour, uptake of interventions and sexual morbidity. This work improves clinical and public health understandings of sexual morbidity but may not be easily translated into practice. We seek to identify key features of research surveys that would make them more relevant to the development and use of risk assessment and targeting tools.

Methods To support the development of clinically-embedded risk assessment tools we undertook a systematic review of population surveys reporting on sexual risk and morbidity. We identified aspects of sexual health research questionnaire items and their reporting which could be adapted to better serve public health and clinical services to identify patients for targeted intervention.