P4.35 REDUCING HIV RISK BEHAVIOUR AFTER INTERVENTION WITH YOUNG LEADERS OF KEY POPULATIONS IN BRAZIL

Diego Agostinho Calixto, Alicia Kruger, Carina Bernardes, Gilvane Casimiro da Silva, Paula Emilia Adamy, Adele Schwartz Benzaken. Ministry of Health of Brazil – Department of STI, HIV/AIDS and Viral Hepatitis, Brasília, Brazil

10.1136/sextrans-2017-053264.532

Introduction Peer dialogues influence the adoption of behavioural changes to reduce the risk of HIV infection (human immunodeficiency virus). By intervening experimentally in the community to change risk behaviour patterns, it may be possible to promote widespread reductions in HIV risk practices within a population.

Methods The intervention identified and trained young age range 18–26 people who are reliably identified as leaders among one of these key populations - gay men and MSM, transgender people (transvestites and transsexual women), drug users and harm reducers and sex workers - in all five regions of Brazil to act as multipliers of behavioural changes for their peers, in relation to HIV. We also include young people living with HIV, considering that it is important that these young people share the experience of living with HIV with other young people in greater vulnerability and risk.

Results 140 young people from the key populations were trained in the 5 Brazilian regions. The proportions of the key populations trained in this intervention were 41.9 per cent homosexuals and MSM, 14.5 per cent harm reduction or drug users, 8 per cent transgender people, 6 per cent sex workers and 15 per cent young people living With HIV. Approximately 70% of young trained in this intervention have already developed some activity to multiply the information about prevention and behavioural practices to reduce HIV infection in their respective territories and communities, promoting knowledge about combination prevention and changes related to sexual practices and behaviours.

Conclusion Interventions that empower young to endorse change can produce or accelerate changes in the behaviour and sexual practices of the young population to reduce the risk of HIV infection. These interventions have developed a network of multipliers in a successive chain of HIV information and combination prevention to reduce HIV infection and risk behaviours through peer-to-peer communication at the community level among young key populations in Brazil.

P4.36 EVALUATING THE COSTS, BENEFITS AND COST-EFFECTIVENESS OF MULTI-PATHOGEN POINT-OF-CARE TESTS FOR SEXUALLY TRANSMITTED INFECTIONS IN SYMPTOMATIC GENITOURINARY MEDICINE CLINIC ATTENDEES

¹Emma Harding-Esch, ²Susie E Huntington, ²Richeal M Burns, ²Mike Harvey, ³Rachel Hill-Tout, ¹Sebastian S Fuller, ²Elisabeth J Adams, ¹S Tariq Sadiq. ¹Applied Diagnostic Research and Evaluation Unit, St George's University of London, London, UK; ²Aquarius Population Health, London, UK; ³St George's University Hospitals NHS Foundation Trust, London, UK

10.1136/sextrans-2017-053264.533

Introduction In sexual health services, availability of rapid and accurate point-of-care tests (POCTs) may enable major improvements in care pathway efficiencies and outcomes. Previous economic evaluations of nucleic acid amplification test (NAAT) POCTs for *Chlamydia trachomatis* (CT) and *Neisseria*

gonorrhoeae (NG) indicate they may provide a cost-effective strategy for screening genitourinary medicine (GUM) attendees. We estimated costs, benefits and cost-effectiveness of three strategies using accurate, rapid NAAT POCTs that could detect different combinations of common multiple sexually transmitted infections (mSTIs) compared with standard care (SC; laboratory-based CT/NG NAAT).

Methods A decision tree was constructed to simulate a hypothetical cohort of 9 65 988 patients, representing annual numbers of GUM attendees in England, symptomatic for lower genitourinary tract infection. The model considered delivery costs (micro-costing) and reimbursement (tariff) to GUM services associated with diagnosing and managing STIs. POCT strategies compared to SC were: A) POCT for CT and NG; B) POCT for CT-NG and *Mycoplasma genitalium* (MG); C) POCT for CT-NG-MG and *Trichomonas vaginalis*. Data came from published literature and unpublished estimates.

Results SC was cheaper than all POCT strategies when micro-costing, but POCT C was the cheapest strategy for tariff costings. POCT C's incremental cost-effectiveness ratio (ICER) was £36 585 per quality-adjusted life year (QALY) gained compared to SC when micro-costing; it was cost-saving (by £26,451,382) when tariff costing was applied. POCT C also generated most benefits, with 2 40 467 fewer clinic attendances, 808 fewer onward STI transmissions, and 2 35 135 averted inappropriate treatments compared to SC.

Conclusion POCTs that detect STI diagnoses may be cost-effective, cost-saving and improve patient management. However, there is variation by costing strategy, patient population, clinical setting and patient pathways. Further evidence is needed to populate model parameters to reduce uncertainty in economic analyses.

P4.37 THE PERCEPTION AND DETERMINANTS OF SEXUAL BEHAVIOUR OF UNIVERSITY UNDERGRADUATES AT A TERTIARY INSTITUTION IN NIGERIA – WHAT ARE THE UNDERLYING FACTORS

¹Emmanuel Shobowale, ¹O Abiodun, ¹CJ Elikwu, ²KI Onyedibe, ¹J Sotunsa, ³O Iyoha. ¹Babcock University, Ogun, Nigeria; ²University of Jos, Plateau, Nigeria; ³University of Benin, Benin, Nigeria

10.1136/sextrans-2017-053264.534

Introduction Sexually Transmitted Infections are widespread, and have effects on the reproductive and sexual health of the general population particularly youths and adolescents. The study was done to assess the level of knowledge, perception and attitudes towards STIs by undergraduate students. The objective was to understand and describe the drivers of sexual behaviour amongst university undergraduates and gain an insight into their perception of sexually transmitted infections.

Methods This was a cross-sectional study. A structured self-administered questionnaire on risk factors for STIs was given to respondents to answer. The study was conducted in October 2016.

Results The mean age of the students (n=310) was 17.8 years $\pm$ 1.77 SD.: males were 0.39 times more likely to be sexually active than females [$p < 0.001$, $X^2 = 15.0$, $CI = 0.23-0.69$], females were 2.18 times more likely to join an abstinence club [$p = 0.005$, $X^2 = 7.6$, $CI = 1.24-3.81$]. Males were 0.3 times more likely to believe condoms protect against all STIs [$p < 0.001$, $X^2 = 10.93$, $CI = 0.14-0.63$] and were 2.61 times