



Highlights from this issue

doi:10.1136/sextrans-2018-053864

Jackie A Cassell, *Editor in Chief*

There is a growing move to think smarter about partner notification. As every clinician knows, patients' willingness to share information about exposure risk with partners or ex-partners, and their preferences as to how to do this, vary by relationship type and infection as well as more personal factors. Despite some raised eyebrows on social media for self-advertisement, Bellhouse and colleagues' paper on the implications of "fuckbuddies" for STI transmission, partner notification and behavioural surveillance makes an important contribution to opening up these questions.¹ Partner notification (PN) needs to be targeted and customised, and the better our understanding of sexual partnership typologies the better we can do this.² Partner notification is more explicitly the focus for Suzan-Monti's analysis of the IPERGAY study data, exploring use of street drugs and sexual contact outside main partnerships as barriers to partner notification.³ The interaction between drug use, chemsex and HIV risk behaviours described by Sewell *et al*⁴ addresses related issues, as does a Mmeje *et al*'s comparative study of diagnosed chlamydia incidence in US states with, and without, expedited partner notification.⁵

It is always good to see research that explores the needs of vulnerable and hard to study populations. This month we publish a study of female seasonal farm workers in Morocco, a study with low HIV prevalence yet significant potential for STI and HIV acquisition and transmission, by Bozicevic and colleagues.⁶

Testing is another key theme this month, with a range of articles spanning HIV self-testing,⁷ the acceptability of oral HIV testing,⁸ hepatitis B and C testing among MSM in China.⁹ At a population level, surveillance of gonococcal antimicrobial resistance testing, in the Caribbean, is addressed by Sawatzky and colleagues.¹⁰

The hinterland between gastrointestinal and sexual pathogens has recently re-emerged as an important topic. "Gay bowel syndrome" is now an archaic usage,

but the sexual transmission of organisms such as hepatitis A, salmonella and shigella has never disappeared as a public health problem. Different aspects of rectal STI in MSM are explored in two contrasting studies. Hassan *et al*¹¹ address the relationship between douching and gonococcal or chlamydial rectal infection in a population using PREP. In an elegant study using unlinked anonymous data to about classically gastrointestinal infections, Hughes *et al* explore their relationship with rectal chlamydia diagnoses.¹²

Further highlights include patient perspectives on 'intentional' HIV transmission,¹³ self perceived STI risk among Scandinavian women,¹⁴ the implications and opportunities of sexual health clinic attendance patterns for HPV vaccination,¹⁵ and trends in hospitalisation for pelvic inflammatory disease.¹⁶

Finally, do not forget to read this month's Clinical roundup.¹⁷

Handling editor Jackie A Cassell

Twitter @jackiecassell

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2018. No commercial re-use. See rights and permissions. Published by BMJ.

REFERENCES

- Bellhouse C, Walker S, Fairley CK, *et al*. Getting the terminology right in sexual health research: the importance of accurately classifying fuck buddies among men who have sex with men. *Sex Transm Infect* 2018;94:487–9.
- Mercer CH, Jones KG, Johnson AM, *et al*. How can we objectively categorise partnership type? A novel classification of population survey data to inform epidemiological research and clinical practice. *Sex Transm Infect* 2017;93:129–36.
- Suzan-Monti M, Cotte L, Fressard L, *et al*. Factors associated with partner notification of STIs in men who have sex with men on PrEP in France: a cross-sectional substudy of the ANRS-IPERGAY trial. *Sex Transm Infect* 2018;94:490–3.
- Sewell J, Cambiano V, Miltz A, *et al*. Changes in recreational drug use, drug use associated with chemsex, and HIV-related behaviours, among HIV-negative men who have sex with men in

London and Brighton, 2013–2016. *Sex Transm Infect* 2018;94:494–501.

- Mmeje O, Wallett S, Kolenic G, *et al*. Impact of expedited partner therapy (EPT) implementation on chlamydia incidence in the USA. *Sex Transm Infect* 2018;94:545–7.
- Bozicevic I, Guezzar F, Stulhofer A, *et al*. HIV prevalence and related risk behaviours in female seasonal farm workers in Souss Massa Draa, Morocco: results from a cross-sectional survey using cluster-based sampling. *Sex Transm Infect* 2018;94:515–7.
- Boukli N, Boyd A, Wendremaire N, *et al*. Sensitivity of the STAT-VIEW rapid self-test and implications for use during acute HIV infection. *Sex Transm Infect* 2018;94:475–8.
- Flores JA, Brown B, León SR, *et al*. Individual-level characteristics associated with oral HIV test acceptability among Peruvian men who have sex with men and transgender women: a cross-sectional study. *Sex Transm Infect* 2018;94:528–33.
- Fitzpatrick T, Pan SW, Tang W, *et al*. HBV and HCV test uptake and correlates among men who have sex with men in China: a nationwide cross-sectional online survey. *Sex Transm Infect* 2018;94:502–7.
- Sawatzky P, Martin I, Galarza P, *et al*. Quality assurance for antimicrobial susceptibility testing of *Neisseria gonorrhoeae* in Latin American and Caribbean countries, 2013–2015. *Sex Transm Infect* 2018;94:479–82.
- Hassan A, Blumenthal JS, Dube MP, *et al*. Effect of rectal douching/enema on rectal gonorrhoea and chlamydia among a cohort of men who have sex with men on HIV pre-exposure prophylaxis. *Sex Transm Infect* 2018;94:508–14.
- Hughes G, Silalang P, Were J, *et al*. Prevalence and characteristics of gastrointestinal infections in men who have sex with men diagnosed with rectal chlamydia infection in the UK: an 'unlinked anonymous' cross-sectional study. *Sex Transm Infect* 2018;94:508–14.
- Brown MJ, Serovich JM, Kimberly JA. Perceived intentional transmission of HIV infection, sustained viral suppression and psychosocial outcomes among men who have sex with men living with HIV: a cross-sectional assessment. *Sex Transm Infect* 2018;94:483–6.
- Guleria S, Faber MT, Hansen BT. Self-perceived risk of STIs in a population-based study of Scandinavian women. *Sex Transm Infect* 2018;94:522–7.
- Bayley J, Mesher D, Nadarzynski T. Attendance of MSM at Genitourinary Medicine services in England: implications for selective HPV vaccination programme (a short communication). *Sex Transm Infect* 2018;94:542–4.
- Goller JL, De Livera AM, Guy RJ, *et al*. Rates of pelvic inflammatory disease and ectopic pregnancy in Australia, 2009–2014: ecological analysis of hospital data. *Sex Transm Infect* 2018;94:534–41.
- Chung E, Herbert S. Clinical round-up. *Sex Transm Infect* 2018;94:548.