

	MSM n=51	Heterosexual men n=106	Women n=325
T1	% (n)/ median (IQR)	% (n)/ median (IQR)	% (n)/ median (IQR)
Age			
16-20 years	7.8 (4)	11.3 (12)	33.5 (109)
21-25 years	23.5 (12)	38.7 (41)	37.2 (121)
≥26 years	68.6 (35)	49.1 (52)	28.9 (94)
Western nationality	5.9 (3)	89.6 (95)	97.5 (317)
Number of sex partners in the past 6 months			
1	9.8 (5)	24.5 (26)	31.4 (102)
2	13.7 (7)	27.4 (29)	30.5 (99)
≥3	76.5 (39)	47.2 (50)	37.8 (123)
Antibiotic use <1 month	5.9 (3)	0.9 (1)	10.8 (35)
Any STI ever			
Yes	57.1 (28)	23.1 (24)	21.6 (70)
Among CT positives			
Urogenital symptoms	25.7 (9)	41.2 (42)	59.6 (186)
Proctitis	8.6 (3)	1.0 (1)	4.8 (15)
Among NG positives			
Urogenital symptoms	21.1 (4)	33.3 (1)	46.7 (7)
Proctitis	15.8 (3)	100 (3)	0
TPHA positivity	5.9 (3)	0	0
HIV	13.7 (7)	0	0
CT/NG co-infection	7.8 (4)	0.9 (1)	1.2 (4)
<b>T1-T2 interval<sup>1</sup></b>			
Antibiotic use	7.8 (4)	1.9 (2)	6.5 (21)
Among CT positives			
Urogenital symptoms	22.9 (8)	32.4 (33)	41.7 (130)
Proctitis	25.7 (9)	2.9 (3)	4.8 (15)
Among NG positives			
Urogenital symptoms	21.1 (4)	3 (100)	13.3 (2)
Proctitis	21.1 (4)	33.3 (1)	6.7 (1)
Unprotected vaginal intercourse	11.8 (6)	50.0 (53)	30.5 (99)
Unprotected anal intercourse	19.6 (10)	0.9 (1)	1.2 (4)

\* <sup>1</sup> 59 consultations had a missing questionnaire about the T1 - T2 interval.

Urogenital symptoms were defined if at least one of the following was reported: genital discharge, bleeding, itching, swelling, pain, burning sensation and more frequent urination. Anorectal symptoms were defined if at least one of the following was reported: rectal discharge, bleeding, pain, redness, burning sensation, or itching.