and C virus infection, genital herpes, amoebiasis, giardiasis, and condyloma, were recorded.

Results A total of 230 MSM completed the interviews from November 2017 through October 2018. The median age was 35 years (range: 18–70). MSM at the high risk was found in 185 (80.4%). MSM who knew PrEP was only 61.3%. Positive willingness of PrEP was observed in 64.8%. MSM who answered “No definitely” were 21 among 185 high risk MSM. Their main negative reasons were “I think my HIV acquisition risk is low” in 12 MSM and “I don’t like to take medicine” in 8. Histories of STI therapy within 1 year were found in 81 MSM (35.2%). Among them, 27 (33.3%) had still the negative willingness. There was no correlation of the positive willingness and history of STIs ($p=0.66$).

Conclusion Still one third of MSM in our cohort lacked the proper knowledge of PrEP. Especially it is important to allow themselves to assess their own risk of HIV infection.

Disclosure No significant relationships.

P557

“FLUX NZ”: AN ONLINE NATIONAL COHORT INVESTIGATING HIV, STI AND DRUG-RELATED PRACTICES AMONG NEW ZEALAND GAY AND BISEXUAL MEN

¹Peter Saxton*, ²Mohamed Hammoud, ³Samuel Andrews, ⁴David Newcombe, ⁵Anthony Walton, ⁶Seb Stewart, ⁷Ricky Te Akau, ⁸Mark Fisher, ⁹Kathryn Leafe, ⁸Carl Greenwood, ⁹J Green, ²Garrett Prestage. ¹University of Auckland, School of Population Health, Auckland, New Zealand; ²University of New South Wales, Kirby Centre, Sydney, Australia; ³New Zealand Drug Foundation, Wellington, New Zealand; ⁴University of Auckland, Centre for Addiction Research (CFAR), Auckland, New Zealand; ⁵New Zealand AIDS Foundation, Auckland, New Zealand; ⁶Body Positive, Auckland, New Zealand; ⁷New Zealand Needle Exchange Programme, Christchurch, New Zealand; ⁸Drugs Health and Development Project, Wellington, New Zealand; ⁹Te Whariki Takapou, Waikato, New Zealand

10.1136/sextrans-2019-sti.631

Background HIV pre-exposure prophylaxis (PrEP) and chemsex present both opportunities and challenges for sexual health improvement among gay and bisexual men (GBM). However, there is scant evidence to inform responses in New Zealand (NZ), notably prevalence, uptake and cessation of behaviours over time. We sought a better understanding by recruiting a diverse national cohort of GBM.

Methods Flux NZ is a national online prospective observational study investigating wellbeing among GBM in NZ. The protocol, questionnaire and data management system were based on a concurrent Australian study. Recruitment was predominantly via social media and community partners. Eligible participants were GBM living in NZ. We present baseline descriptions of PrEP, STI and illicit drug-related data among the cohort to date.

Results The 315 participants are broadly reflective of the target population (10.8% identified as indigenous Maori, 59% lived outside Auckland the largest city, mean age 33.6 years (SD 15.6), 8.3% diagnosed HIV-positive). Among HIV-negative participants, 50% had heard “a lot” and 44% “a little” about PrEP; 18.2% were currently using PrEP. 57% of all participants had tested for STIs in the past 6 months; 7%, 5.7% and 3.2% were diagnosed with gonorrhoea, chlamydia and syphilis respectively. Drug use in the past 6 months was 37.5% cannabis, 35.6% amyl, 17.5% MDMA, 5.1% methamphetamine, 4.8% LSD,

4.4% cocaine, 3.2% GHB, 2.5% amphetamine, 1.0% ketamine, 0.3% synthetic cannabis, 0% mephedrone, 0% heroin, and 1.9% other hallucinogens/psychedelics. “Any” drug use was 57.8% in the past 6 months and 78.7% ever. Of those using “party drugs” (MDMA, amphetamine, cocaine, methamphetamine, GHB, ketamine, LSD) recently, 29.2% had done so to enhance sex, the majority “once” (4.7%) or “a few times” (15.1%).

Conclusion The Flux online cohort design effectively recruits GBM in NZ. This will enable comparative research of the impact of HIV, STI and drug harm reduction policies in the two countries.

Disclosure No significant relationships.

P558

SEXUAL HEALTH, SYNDEMICS AND ASSETS AMONG MEN WHO HAVE SEX WITH MEN: SECONDARY ANALYSIS OF MULTI-NATIONAL SURVEYS

¹Lisa Mcdaid, ¹Paul Flowers, ²Olivier Ferlatte, ³Kareena Mcaloney-Kocaman, ⁴Mark Gilbert, ³Jamie Frankis. ¹University of Glasgow, MRC/CSO Social and Public Health Sciences Unit, Glasgow, UK; ²BC Centre for Substance Use, Vancouver, Canada; ³Glasgow Caledonian University, Glasgow, UK; ⁴BC Centre for Disease Control, Clinical Prevention Services, Vancouver, Canada

10.1136/sextrans-2019-sti.632

Background Globally, gay, bisexual and other men who have sex with men (GBMSM) experience an increased burden of poor sexual, mental and physical health. Syndemics theory provides a framework to understand these interrelated health problems and how to intervene.

Methods Comparative quantitative secondary analysis of syndemic-related data are presented from two international, online, cross-sectional surveys: SMMASH2 in Scotland, Wales, Northern Ireland and the Republic of Ireland (N=3220); and Sex Now in Canada (N=7872).

Results In both studies negative sexual, mental and physical health outcomes were clustered, providing evidence of the syndemic (SMMASH2 O/E Ratio=1.32, 95% CI 1.25–1.40; Sex Now O/E Ratio=1.59, 95% CI 1.45–1.73). There were differences between the studies in the variables that were associated with experience of the syndemic. In Sex Now we found experience of the syndemic was associated with worries about sexuality-related stigma (AOR 1.87, 95% CI 1.23–1.54) and experience of discrimination (AOR=1.83, 95% CI 1.60–2.10). Equally, some community assets appeared to have a protective effect on the experience of syndemics. In Sex Now, aspirations were significant; e.g., being unlikely to achieve quality of life significantly associated with increased odds of experiencing the syndemic (AOR=1.89, 95% CI 1.63–2.19), while measures of community engagement were not significant. In SMMASH2, sense of coherence – a measure of resilience – was significant (AOR=0.98, 95% CI 0.96–0.99), indicating that higher sense of coherence was associated with decreased odds of experiencing the syndemic.

Conclusion These results present an important step forward in our understanding of syndemics. They provide new insights into how to intervene to reduce the interrelated burden of poor sexual, mental and physical health among GBMSM and point to a theoretical mechanism through which assets-based approaches to health improvement could function.

Disclosure No significant relationships.