P717

USING ETHNOGRAPHY FOR PLANNING AND IMPROVING THE QUALITY OF HIV PREVENTION INTERVENTIONS FOR FEMALE SEX WORKERS IN NIGERIA

¹Baba Mari, ¹Kalada Green*, ²Shajy Isac, ¹Janet Halliday, ¹Judith Ariri-Edeafe, ³Kayode Ogungbemi, ⁴Robert Lorway, ⁵James Blanchard. ¹Centre for Global Public Health – Nigeria, Abuja, Nigeria; ²Centre for Global Public Health, University of Manitoba, Winnipeg, Canada; ³National Agency for the Control of AIDS, Abuja, Nigeria; ⁴Center for Global Public Health, Department of Community Health Sciences, University of Manitoba, Winnipeg, Canada; ⁵University of Manitoba, Centre for Global Public Health, Department of Community Health Sciences, Winnipeg, Canada

10.1136/sextrans-2019-sti.781

Background Female Sex Workers (FSW) and their clients account for half of HIV infections among key populations (KP) in Nigeria. The aim of this study was to better understand sexual behavior patterns and influences related to HIV vulnerability among men and women in venues that facilitates sexual networks in Nigeria as it relates to sex work.

Methods A qualitative approach was employed using participant observations and face-to-face in-depth key informant interviews. Community ethnographers collected information from FSWs, clients of FSWs and key venue staff at venues where FSWs solicited clients. A semi-structured tool guided the interviews and questions explored how connections occur for the sale of sex and probed the power dynamics within connections.

Results In all, 25 women and 25 men were interviewed. The in-depth interviews describe bar staff as vital players in the sex trade who provide financial and security services for FSWs. In exchange for these services FSWs are expected to pay with cash, commodities or sex. Majority of FSWs experienced violence from clients and law enforcement agents. Interviews indicate that there was a lack of governmental or non-governmental organizations involved in addressing cases of violence. Participants reported low condom use with intimate partners compared to paying clients. Top on the list of the problems associated with condom use was condom breakage, which was attributed to dryness, lack of lubricants and lengthy duration of sex. In addition, both male and female participants had power over decision making on condom use. Generally, as noted in transcripts, interactions between FSWs and clients switch between commercial sexual relationships and transactional liaisons.

Conclusion The study provides insight into the structure and operational dynamics occurring at venues facilitating sexual networks. Simultaneously exposing the barriers and challenges to HIV prevention service delivery. A combination of strategies-behavioural, biomedical, structural interventions are required in addressing the needs of FSWs.

Disclosure No significant relationships.

P718

RISK AND PLACE: THE ASSOCIATION BETWEEN HOTSPOT TYPOLOGY AND DETERMINANTS OF HIV RISK AMONG FEMALE SEX WORKERS IN UKRAINE

¹Eve Cheuk, ¹Stella Leung, ²Olga Balakireva, ³Daryna Pavlova, ¹Leigh Mcclarty, ⁴Shajy Isac, ¹Michael Pickles, ⁵Sharmistha Mishra, ⁶Evelyn Forget, ¹Robert Lorway, ⁷Paul Sandstrom, ¹James Blanchard, ¹Marissa Becker*. ¹University of Manitoba, Centre for Global Public Health, Department of Community Health Sciences, Winnipeg, Canada; ²Institute for Economics and Forecasting, Ukrainian National Academy of Sciences, Department for Monitoring-based Research of Social and Economic Transformations, Kyiv, Ukraine; ³NGO "Ukrainian Institute for Social Research after Oleksandr Yaremenko", Monitoring and Evaluation of Social Projects, Kyiv, Ukraine; ⁴India Health Action Trust, Delhi, India; ⁵St. Michael's Hospital, Li Ka Shing Knowledge Institute, Toronto, Canada; ⁶University of Manitoba, Department of Community Health Sciences, Winnipeg, Canada; ⁷Public Health Agency of Canada, National HIV and Retrovirology Laboratories, JC Wilt Infectious Diseases Research Centre, Winnipeg, Canada

10.1136/sextrans-2019-sti.782

Background The environmental contexts and interactions between people within that environment impact a female sex worker's (FSW) individual behaviour and likelihood of HIV acquisition. Understanding how the environment shapes HIV risk can provide important information for HIV prevention programs that go beyond behaviour-based interventions and address more distal factors contributing to HIV risk among FSWs.

Methods A cross-sectional survey was conducted among FSWs in Dnipro, Ukraine between September 2017 and January 2018. 560 FSWs aged ≥ 14 years were recruited from "hot-spots" (locations where FSWs meet clients). We compared determinants of HIV risk between six hotspot typologies (e.g., brothels, home-based, massage parlours, bars/restaurants, parks, highways) using chi-squared tests.

Results Age at start of sex work ($p < 0.001$), client volume ($p < 0.001$), and HIV testing in the past year ($p < 0.001$) all differed significantly across hotspot typologies. Mean age at start of sex work was oldest for FSWs recruited from highways (24.1 years) and youngest for FSWs recruited from massage parlours (19.8 years). FSWs from highways had the most clients in the past 30 days (mean 22.1 clients) while those from parks had the fewest (mean 17.5 clients). HIV testing in the past year was the most frequent among FSWs from brothels (mean 2.1 times), followed by home-based (1.8 times), massage parlours (1.1 times), highways (1.0 time), bars/restaurants (0.6 times) and parks (0.4 times). The prevalence of physical and sexual violence perpetrated by clients in the past 3 months was 4.3% and 1.6% overall. Compared to women from other hotspot typologies, the proportion of FSWs from highways who reported experience of physical and sexual violence by clients were highest (26.7% and 10.0%).

Conclusion Understanding the association between HIV risk factors and hotspot typology can help HIV prevention programs tailor interventions, and target linkage and delivery of services to the relevant subgroups of a key population.

Disclosure No significant relationships.