

consecutively with total of 127 samples. All steps in this research; history taking, physical examination, and blood tests were done blindly.

Results The results of this study using serum specimens were sensitivity of 91.30%, specificity of 97.53%, positive predictive value 95.45%, negative predictive value of 95.18%, and accuracy 95.28. Test results with fingerprick whole blood specimens gave sensitivity of 84.78%, specificity of 98.77%, positive predictive value of 97.50%, negative predictive value of 91.95%, and accuracy 93.70%. Compatibility of rapid test STANDARD™ Q Syphilis Ab results between serum and fingerprick whole blood specimens was very good ($\kappa=0.8223$).

Conclusion Rapid test STANDARD™ Q Syphilis Ab can be used as an option for treponemal test in supporting syphilis diagnosis, either as routine screening or confirmation of non-treponemal test result. The fingerprick whole blood specimen can be used as treponemal test alternative which is faster and easier to do.

Disclosure No significant relationships.

P750 A COMPARISON OF *TREPONEMA PALLIDUM* MOLECULAR TYPING SYSTEMS: MLST VS. ECDCT

¹Sharon Sahi*, ¹Lauren Tantaló, ²Christina Marra. ¹University of Washington, Neurology, Seattle, USA; ²University of Washington, USA

10.1136/sextrans-2019-sti.808

Background Several syphilis typing systems have been proposed. Recent work suggests that multilocus sequence typing (MLST) may be superior to enhanced CDC typing (ECDCT), particularly because ECDCT type may differ among organisms amplified from different anatomical sites in the same person. The goal of this study was to compare the two systems.

Methods DNA was extracted from 20 *Treponema pallidum* isolates propagated in rabbits, 10 oral and 10 genital or non-genital lesion swabs, and 10 blood samples from patients with syphilis. MLST type for *tp0136*, *tp0548* and *tp0705* and ECDCT type were determined according to published methods. Samples were chosen because they were completely typeable by ECDCT. ECDCT types were also determined for samples from different anatomical sites in 7 patients, and from blood and blood isolates (rabbit propagated) in 8 patients.

Results MLST type could be fully determined for 19 (95%) of 20 bacterial isolates, 8 (80%) of 10 bloods, 7 (70%) of 10 lesion swabs, and 5 (50%) of 10 oral swabs. 13 subtypes were identified by ECDCT, and 12 by MLST. While MLST was able to subdivide two common ECDCT types (1.1.1, 1.1.2, 1.1.9, and 1.37.1 within 14d/f; and 1.3.1, 1.38.1, and 6.3.1 within 14d/g), it failed to distinguish among less common ECDCT types. ECDCT type was identical in 5 paired lesion and oral swabs, 1 paired blood and oral swab, and 1 paired blood, lesion and oral swabs. In addition, ECDCT type was identical in 8 paired blood and blood isolates.

Conclusion Compared to ECDCT, determination of MLST was less often successful from isolates and from clinical samples, and it was not uniformly more discriminating. ECDCT was stable among anatomical sites and between direct patient-derived samples compared to rabbit propagated organisms.

Disclosure No significant relationships.

P751 EVALUATION OF A SYPHILIS AWARENESS CAMPAIGN FOR GAY, BISEXUAL AND OTHER MSM (GBMSM): DID WE REACH OUR TARGETED AUDIENCE?

¹Jason Wong*, ²Shenyi Pan, ¹Emma Cumming, ²Heather Armstrong, ²Nicanor Bacani, ¹Devon Haag, ³Venessa Ryan, ¹Jillian Arkles Schwandt, ¹Troy Grennan, ⁴Jody Jollimore, ⁵Nathan Lachowsky, ²David Moore. ¹BC Centre for Disease Control, Clinical Prevention Services, Vancouver, Canada; ²BC Centre for Excellence in HIV/AIDS, Epidemiology and Population Health, Vancouver, Canada; ³British Columbia Centre for Disease Control, Clinical Prevention Services, Vancouver, Canada; ⁴Community Based Research Centre, Vancouver, Canada; ⁵University of Victoria, School of Public Health and Social Policy, Victoria, Canada

10.1136/sextrans-2019-sti.809

Background In 2017, we launched a syphilis awareness campaign (“Syphistory”) targeted towards gay, bisexual, and other men who have sex with men (gbMSM). Using data from a study of gbMSM in Vancouver, we describe participants who reported seeing Syphistory and whether it reached gbMSM at higher risk of syphilis.

Methods Participants aged ≥ 16 years who reported having sex with another man in the previous six months were recruited through respondent-driven sampling. We analyzed data collected from September 17, 2017 to August 31, 2018. Characteristics of participants who reported seeing the campaign were compared using Wilcoxon rank-sum and chi-square/Fisher’s exact test. Multivariate logistic regression was used to examine the association between seeing Syphistory and recently being tested for syphilis, controlling for potential confounding factors, namely age, HIV status, place of residence, education level, and recent illicit drug use (IDU).

Results Of the 383 participants who responded, 103 (27%) reported seeing Syphistory. Participants who saw Syphistory tended to be younger (median 30 vs 32 years old, $p=0.03$), live in downtown Vancouver (58% vs 38%, $p=0.46$), had greater than a high school education (97% vs 89%, $p=0.08$), and were known to be HIV-negative (92% vs 81%, $p=0.15$). They reported more male sex partners (median 4 vs 3, $p=0.12$), condomless anal sex (92% vs 81%, $p=0.06$), and IDU (86% vs 72%, $p=0.04$) in the last 6 months. Almost 75% (52/86) of those who saw Syphistory reported a syphilis test within the last 3 months compared with 58% (97/244) who did not see Syphistory ($p=0.01$). Participants who saw Syphistory had a greater odds (aOR=3.63; 95% CI, 1.28–10.27) of a syphilis test within 3 months, versus no or unknown previous syphilis test.

Conclusion Participants who saw Syphistory tended to report behaviours that may increase the risk of syphilis infection and were more likely to have had a recent syphilis test.

Disclosure No significant relationships.

P752 UNDERSENSITIVE NON-TREPONEMAL TESTS: IMPLICATIONS FOR SYPHILIS MANAGEMENT

John Scythes*, Colman Jones. Community Initiative for AIDS Research, Toronto, Canada

10.1136/sextrans-2019-sti.810

Background Syphilis management has traditionally been based on non-treponemal tests (NTT). However, investigations over 3 decades have revealed serious issues with the sensitivity of NTT.

Methods Beginning in the late 1980s, our group in Toronto and Budapest investigated the sensitivity of NTT in over 5000 high-risk gay men, and in an additional 800 men where we could control for HIV status and AIDS. Besides the use of quantified treponemal tests (TT), IgM/IgG screening (in Mardx SDS-PAGE blots), and experimental PCR were employed.

Results The main findings of accelerated screening were undetected and untreated latent cases, not old treated cases as traditionally believed. NTT only detected about 30 percent of cases. Investigators in Houston detected TT IgM in most HIV cases with no NTT reactions, possibly representing latent and chronic cases. Investigators in Vienna showed that aggressive therapy reversed the IgM marker. In Toronto, 557 high risk men were screened with EIA TT and 27 possible latent cases were detected, with negative NTT. 24/27 of these patients had IgM or IgG directed against the main *T. pallidum* proteins. In the 800 men screened sequentially, TT often dropped in titre or reverted to negative *only* in HIV cases - titres in other routine tests did not drop. Syphilis PCR found that 13/183 gay men screened in Hungary had latent syphilis. The PCR found 9 cases negative in both NTT and TT, and identified the four TPHA (+) treated men.

Conclusion NTT is historically unreliable in relapse or reinfection. Syphilis management should always include TT. Syphilis and HIV have an overwhelming association, yet syphilis is never opportunistic. Latent syphilis may be chronically active in many HIV persons. Th-1 \rightarrow Th-2 immunoregulation is the norm in untreated syphilis - many longstanding syphilitics have cutaneous anergy to TB and mitogens. We suggest a syphilis IGRA be developed.

Disclosure No significant relationships.

P753

MONITORING THE EVALUATION OF CONGENITAL SYPHILIS CASES IN THE STATE OF SÃO PAULO, 2007 TO 2017

Carla Gianna Luppi, Solange Gomes, Angela Tayra, Mariza Tancredi, Carmen Silvia Domingues*. *STI/AIDS Reference Center, Sao Paulo State Program of STI/AIDS, Sao Paulo Department of Health, Sao Paulo, Brazil*

10.1136/sextrans-2019-sti.811

Background Congenital syphilis (CS) is still considered a public health problem in the state of São Paulo, Brazil: 4,039 cases were reported in 2017 -incidence rate = 6.6 per 1,000 live births. The objective of this study was to investigate the evaluation of CS cases in 2007 and 2017 in the state of São Paulo.

Methods A descriptive study was carried out with the data of national surveillance system in the period 2007 to 2017. Of the following evaluation tests performed the percentage of non-performed and positive results were calculated: - Venereal Disease Research Laboratory (VDRL) on the serum; - long-bone radiography; -cerebrospinal fluid (CSF) for cell count and protein; - CSF for VDRL.

Results From 2007 to 2017, 21,524 cases of CS born alive were reported: 743 in 2007 and 3,687 in 2017. The percentage of non-performed and positive tests in 2007 and 2017, and the percentage variation (PV) between the two years were: - **non-performed** - VDRL on the serum (4.0% and 1.8%, PV = -56.3%), CSF for VDRL (23.4% and 21.4%, PV = -7.1), CSF for cell count and protein (23% and

21,4%: PV = 7,1%), long-bone radiography (13,7% and 9.9%, PV = -27.9%); - **positive results** - VDRL on the serum (59.1% and 81.1%; PV = 37.3%), CSF for VDRL (1.3% and 3.5%; PV = 157.9%), CSF for cell count and protein (1.6% and 10.1%: PV = 523%), long-bone radiography (2.6% and 3.5%, PV = 36.8%).

Conclusion There was an improvement in adherence to the CS evaluation protocol. In addition, we highlight the increase in the percentage of altered tests of children born alive in the period, pointing to the greater severity of the cases, and the need of improving the monitoring of these children in order to reduce the impact of congenital syphilis.

Disclosure No significant relationships.

P754

QUANTITATION OF CYTOKINES IN RABBITS FOLLOWING TRI-ANTIGEN VACCINE COCKTAIL IMMUNIZATION AND *T. PALLIDUM* CHALLENGE

¹Charmie Godornes*, ¹Barbara Molini, ¹Lorenzo Giacani, ²Alloysius Gomez, ³Darrick Carter, ²Caroline Cameron, ³Sheila Lukehart. ¹University of Washington, Medicine, Seattle, USA; ²University of Victoria, Biochemistry and Microbiology, Victoria, Canada; ³University of Washington, Global Health, Seattle, USA

10.1136/sextrans-2019-sti.812

Background Immunological analysis of primary and secondary syphilis in rabbits and humans suggests that T helper cells mount a vigorous interferon- γ -dominated immune response (T_H1) to facilitate macrophage-mediated clearance of *T. pallidum*. In this study, we used quantitative reverse transcriptase (qRT)-PCR to evaluate post-challenge rabbit cytokine profiles of primary lesions in unimmunized rabbits and rabbits immunized with a tri-antigen vaccine cocktail.

Methods Groups of 8 male New Zealand White rabbits were immunized with a trivalent recombinant antigen cocktail, (N-term of TprK + N-term of Tpr Subfamily I + Tp 0751, emulsified in either of two custom adjuvants containing Natural or Synthetic TLR4 agonists + a natural Mincle agonist). Unimmunized control and immunized animals were intradermally challenged with 10^5 *T. pallidum* (Nichols) at each of 10 sites. Lesion biopsies were collected at days 2 and 21 post-challenge. Expression of IFN- γ , TGF- β , p40 IL-12/23, IL-4, IL-2, TNF- α , IL-10, IL-17A, IL-17F, and IL-22 was quantified by qRT-PCR using plasmids containing the target rabbit cytokines sequences, and expression levels were normalized to rabbit HPRT.

Results At day 2, transcripts for IFN- γ , IL-2, IL-17A, IL-17F, and TNF- α were significantly upregulated in both immunized groups ($P < 0.01$) compared to controls, a finding consistent with the development of clinical delayed type hypersensitivity and the induction of a T_H1 -type immune response at challenge sites. At day 21, the level of IFN- γ was lower in the Natural adjuvant group ($P < 0.02$), compared to controls, consistent with enhanced treponemal clearance in that group.

Conclusion In the development of a syphilis vaccine, it is important to determine correlates of protection to allow for assessment of the induction of a protective immune response. Our results demonstrate a robust immunization-induced T_H1 and T_H17 proinflammatory response in immunized rabbit groups, which suggests an effective level of resistance conferred by immunization with the trivalent protein vaccine.

Disclosure No significant relationships.