P842 HEALTHCARE ENGAGEMENT AND HPV VACCINATION AMONG GAY, BISEXUAL, AND OTHER MEN WHO HAVE SEX WITH MEN (GBMSM): A CIRN STUDY

¹Ramandip Grewal*, ²Anna Yeung, ³Marc Brisson, ⁴Troy Grennan, ⁵Alexandra De Pokomandy, ⁶Joseph Cox, ⁷Gilles Lambert, ⁸David Moore, ⁹François Coutlée, ¹⁰Shelley Deeks, ¹¹Sandra Gardner, ¹²Dane Griffiths, ¹³Wanrudee Isaranuwatthai, ¹⁴Jody Jollimore, ¹⁵James Murray, ¹⁶Gina Ogilvie, ¹⁷Chantal Sauvageau, ¹⁸Darrell Tan, ¹⁹Barry Adam, ²⁰Heather Armstrong, ²¹Mark Gaspar, ²²Clemon George, ²³Daniel Grace, ²⁴Trevor Hart, ²⁵Ann Burchell. ¹St. Michael's Hospital, Toronto, Canada; ²St. Michael's Hospital, Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute, Toronto, Canada; ³Université Laval, Quebec City, Canada; ⁴BC Centre for Disease Control, Clinical Prevention Services, Vancouver, Canada; ⁵McGill University, Montreal, Canada; ⁶University of Montreal, Montreal, Canada; ⁷BC Centre for Excellence in HIV/AIDS, Vancouver, Canada; ⁸Université de Montréal, Microbiologie Médicale et Infectiologie, Montreal, Canada; ⁹Public Health Ontario, Toronto, Canada; ¹⁰University of Toronto, Toronto, Canada; ¹¹Gay Men's Sexual Health Alliance, Toronto, Canada; ¹²Community Based Research Centre, Vancouver, Canada; ¹³Ontario Ministry of Health and Long-Term Care, Toronto, Canada; ¹⁴University of British Columbia, Vancouver, Canada; ¹⁵Institut National de Santé Publique du Québec, Montreal, Canada; ¹⁶University of Windsor, Windsor, Canada; ¹⁷University of West Indies – Cave Hill, Cave Hill, Barbados; ¹⁸University of Toronto, Dalla Lana School of Public Health, Toronto, Canada; ¹⁹Ryerson University, Psychology, Toronto, Canada

10.1136/sextrans-2019-sti.888

Background Canada was one of the first countries to offer publicly funded programs providing HPV vaccine free of charge to gbMSM. In 2015–2016, the provinces of British Columbia (BC), Ontario (ON), and Quebec (QC) implemented programs for gbMSM aged 9–26 years. We sought to explore where men received the HPV vaccine and the influence of healthcare engagement on vaccination.

Methods Engage is a sexual health study among gbMSM aged 16+ in the largest urban centres in each province: Vancouver, BC; Toronto, ON; and Montreal, QC. Men are recruited via respondent driven sampling (RDS). We compared proportions (non-RDS adjusted) to questionnaire responses on healthcare engagement among vaccinated (1+ doses) versus unvaccinated in the subset of men aged ≤26 years old at enrolment.

Results From 01/2017 to 31/12/2018, 477 men aged ≤26 enrolled (144 Vancouver, 84 Toronto, 249 Montreal). Their median age was 24 years (IQR 22–25). In Vancouver, Toronto and Montreal, respectively, 48.6%, 44.1%, 44.2% had initiated HPV vaccination with 58.6%, 56.8% and 48.2% of vaccinated men having received all 3 doses. Popular venues for vaccination included a: sexual health clinic (50.7%), medical clinic not specializing in sexual health (14.8%), community health centre (10.6%), and doctor's office (7.4%). Compared to unvaccinated men, more vaccinated men had a family doctor (65.9% vs 52.3%, $p=0.0047$), had a different provider/clinic for sexual health care or had a HIV care provider (66.8% vs 51.8%, $p=0.0022$), had a STI/HIV test in the past year (90.8% vs 68.6%, $p<0.0001$), and were diagnosed with a STI in the past year (39.6% vs 20.0%, $p<0.0001$).

Conclusion Compared to unvaccinated men, vaccinated men were more engaged in healthcare. It is unknown whether men requested the vaccine or providers offered it. Nevertheless, our findings suggest opportunistic HPV vaccination when men receive other services, particularly those related to sexual health.

Disclosure No significant relationships.

P843 THE TRANSMISSION OF HPV IN THE SEXUAL NETWORKS OF SWINGERS, A MULTILEVEL NETWORK APPROACH

¹Anne-Marie Niekamp*, ²Fleur Koene, ³Petra Wolffs, ⁴Christian Hoebe. ¹Public Health Service South Limburg, Maastricht University Medical Center (MUMC), Sexual Health, Infectious Diseases and Environmental Health, Medical Microbiology, Care and Public Health Research Institute (CAPHRI), Heerlen, Netherlands; ²Amsterdam UMC, Medical Microbiology, Amsterdam, Netherlands; ³Maastricht University Medical Center (MUMC), Medical Microbiology, Care and Public Health Research Institute (CAPHRI), Maastricht, Netherlands

10.1136/sextrans-2019-sti.888

Background Swingers, i.e. heterosexual couples who, as a couple, have sex with others within the swingers' subculture, are a subpopulation at risk for Sexually Transmitted Infections (STI). Human papillomavirus (HPV) has proven to be highly prevalent in female swingers (unpublished data). The aim of this study is to assess how the different HPV genotypes are distributed within the sexual networks of swingers and whether certain HPV genotypes cluster.

Methods We build a sexual network of swingers based on the network data given by 115 swingers who participated in our prospective cohort network study on swingers between April 2011 and March 2012. The vaginal samples of all female swingers in the network and known in our STI clinic were tested for HPV. HPV genotypes were identified by the highly sensitive SPF₁₀-LiPA₂₅ method detecting 25 different genital HPV genotypes. The multilevel transmission network consisted of the one-mode sexual network of the swingers and the two-mode network of swingers and the HPV genotypes. Distribution and clustering of HPV genotypes in the sexual network was assessed using multilevel network analysis.

Results The sexual network contained 1231 individual swingers of whom 556 female swingers. The sexual network consists of 38 separate components. Of 103 female swingers known in our STI clinic and tested for HPV, 96 (93%) were positive for at least one of the 25 HPV genotypes. All 25 HPV genotypes were present, but not equally distributed throughout the network. Some genotypes clustered together in specific components of the sexual network.

Conclusion All assessed genotypes of HPV were prevalent in the sexual networks of swingers. Despite the incomplete data in our study, multilevel network analysis turns out to be useful for studying transmission networks. Clustering of highly prevalent HPV could be used as a marker for sexual behaviour, potential STI risk and transmission in swingers.

Disclosure No significant relationships.

P844 THE BURDEN OF HPV AND HSV-RELATED ANOGENITAL DISEASES IN A SOUTHEASTERN US URBAN HIV CLINIC

¹Yuanfan Ye*, ²Greer Burkholder, ³Howard Wiener, ⁴Sadeep Shrestha. ¹University of Alabama at Birmingham, Epidemiology, Birmingham, USA; ²University of Alabama at Birmingham, Medicine, Birmingham, USA

10.1136/sextrans-2019-sti.889

Background Although HPV- and HSV-related conditions (HPVCs, HSVCs) have increased in PLWH, incidence, racial, and gender disparities of the conditions are unclear. We