antigens using immunoblot and immunoprecipitation experiments.

Results Although vaccination with WT dOMVs significantly enhanced gonococcal clearance relative to adjuvant-only controls, vaccination with Δ ABR dOMVs resulted in clearance of a higher percentage of mice relative to WT dOMV-vaccinated mice one week post-immunization. Higher levels of clearance in Δ ABR dOMV-immunized mice correlated with significantly increased vaginal IgA titers and enhanced immunogenicity of unique meningococcal protein antigens.

Conclusion Immunization with meningococcal dOMVs deleted for PorA, PorB, and RmpM promotes gonococcal clearance in a murine model. Deletion of the major porins likely enhances immunogenicity of proteins that are less abundant on the meningococcal surface but exhibit a high degree of homology with corresponding gonococcal proteins, suggesting the potential utility of these dOMVs as a broadly cross-protective *Neisseria* vaccine.

Disclosure No significant relationships.

002 – EXTRAGENITAL BACTERIAL STIS: EPIDEMIOLOGY, NATURAL HISTORY, TESTING AND ANTIMICROBIAL RESISTANCE

Monday, July 15, 2019

10:45 AM – 12:15 PM

002.1 WHAT IS THE OPTIMAL TESTING STRATEGY FOR OROPHARYNGEAL *NEISSERIA GONORRHOEA* IN WOMEN VISITING STI CLINICS?

^{1,2}Geneviève Van Liere, ^{1,2}Nicole Dukers-Muijers, ³Sophie Kuizenga-Wessel, ^{4,5,6}Hannelore Götz, ^{1,2}Christian Hoebe*. ¹Public Health Service South Limburg, Dept of Sexual Health, Infectious Diseases and Environmental Health, Heerlen, The Netherlands; ²Maastricht University Medical Center (MUMC+), Dept of Medical Microbiology, Care and Public Health Research Institute (CAPRI), Maastricht, The Netherlands; ³Public Health Service Haaglanden, Dept of Sexual health, Den Haag, The Netherlands; ⁴Public Health Service Rotterdam Rijnmond, Dept of Infectious Disease Control, Rotterdam, The Netherlands; ⁵Erasmus MC University Medical Center Rotterdam, Dept of Public Health, Rotterdam, The Netherlands; ⁶National Institute for Public Health and the Environment (RIVM), Centre for Infectious Disease Control, Epidemiology and Surveillance Unit, Bilthoven, The Netherlands

10.1136/sextrans-2019-sti.110

Background Oropharyngeal *Neisseria gonorrhoeae* (*N. gonorrhoeae*) is not routinely tested for in women visiting Dutch STI clinics. It is hypothesized that many oropharyngeal *N. gonorrhoeae* infections remain undetected due to its asymptomatic nature, creating a reservoir for ongoing transmission and drug resistance. It is yet unknown what the optimal testing policy is for women, as data on universal screening are missing.

Methods Surveillance data 2008–2017 from all Dutch STI clinics were used (n=546,246 consultations). Oropharyngeal testing policy was defined as (1) universal screening, that is >85% of consultations included oropharyngeal testing per clinic per year, (2) selective testing (<85% tested) or (3)

incidental testing (0.1–5% tested). The proportion infections missed using selective testing was calculated by extrapolating *N. gonorrhoeae* positivity found by routine universal screening. Independent risk factors for oropharyngeal *N. gonorrhoeae* were assessed among women routinely universally screened between 2016–2017 using backward multivariable logistic regression analyses.

Results Routine universal screening was used in 11% (n=57,359) of consultations, selective testing in 81% (n=444,283) and incidental testing in 8% (n=44,108). Oropharyngeal *N. gonorrhoeae* positivity was comparable between universal and selective; 1.4%(95%CI 1.3–1.5,n=703), 1.4%(95%CI 1.3–1.3,n=1858,P=0.68), and higher in incidental 2.8%(95%CI 1.9–3.9,n=30, P<0.01). Selective testing missed 89% (n=5,517) of oropharyngeal infections (95%CI 88%–90%). The proportion oropharyngeal-only was 47% in routine universal screening and 52% in selective testing. Independent risk factors were being notified for any STI (OR1.3,95%CI1.03–1.5), concurrent urogenital *N. gonorrhoeae* (OR80.0,95%CI59.0–108.4) and commercial sex work (OR4.1,95%CI2.8–5.9). When using the risk factors except urogenital *N. gonorrhoeae* as testing indicators, 27.8% (n=5,418) of all women would be tested, finding 55.6% (n=119) of infections.

Conclusion Selective testing potentially misses almost 90% of oropharyngeal *N. gonorrhoeae* in women, of which almost half were oropharyngeal-only infections. Using two risk factors as testing indicators, half of all oropharyngeal *N. gonorrhoeae* infections would be detected by testing almost one-third of women. This seems like a valid and minimal testing strategy for women, as is advocated in the Dutch STI-guidelines.

Disclosure No significant relationships.

002.2 OROPHARYNGEAL AND GENITAL GONORRHOEA AMONG HETEROSEXUALS WHO REPORT SEXUAL CONTACT WITH PARTNERS WITH GONORRHOEA

¹Eric Chow*, ¹Marcus Chen, ²Deborah Williamson, ¹Catriona Bradshaw, ¹Sabrina Trumpour, ²Benjamin Howden, ¹Christopher Fairley. ¹Alfred Health, Melbourne Sexual Health Centre, Carlton, Australia; ²The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, Microbiological Diagnostic Unit Public Health Laboratory, Parkville, Australia

10.1136/sextrans-2019-sti.111

Background Recent evidence has shown that the oropharynx may be the primary driver for gonorrhoea transmission among men who have sex with men, but there have been very limited studies on heterosexuals due to lack of routine screening of oropharyngeal gonorrhoea. The aim of this study was to examine oropharyngeal gonorrhoea positivity among heterosexuals reporting contact with sexual partners with gonorrhoea.

Methods At the Melbourne Sexual Health Centre, all heterosexual individuals reporting contact with sexual partners with gonorrhoea are tested for genital gonorrhoea. In May-2017, MSHC also included screening for oropharyngeal gonorrhoea in heterosexuals reporting sexual contact with partners with gonorrhoea. All contacts of gonorrhoea cases among heterosexuals between May-2017 and November-2018 were